



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

February 19, 2026

Quinn Peacock
Regulatory Affairs Manager
BASF Agricultural Solutions
US LLC, 26 Davis Drive
North Carolina 27713-2839

Subject: Label Amendment - Registration Review Mitigation for Pyraclostrobin
Product Name: Headline SC Fungicide
EPA Registration Number: 7969-289
Case Number: 476563
Application Dates: 12/7/2020

Dear Quinn Peacock:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Pyraclostrobin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Tiffany Green by phone at 919-541-2446, or via email at green.tiffany@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kevin Costello". The signature is written in a cursive style with a large initial "K".

Kevin Costello, Branch Chief
Risk Management and Implementation Branch 2
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label



We create chemistry



Headline® SC

Fungicide

For use in disease control and plant health in the following crops: alfalfa, barley, citrus fruits, corn (all types), cotton, dried shelled peas and beans, edible-podded legume vegetables, grass grown for seed, mint, oats, oilseed crops, peanut, pecan, rye, sorghum, soybean, succulent shelled peas and beans, sugar beet, sugarcane, tuberous and corm vegetables (includes potato), and wheat and triticale

Active Ingredient*:

pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester) 23.3%

Other Ingredients: 76.7%

Total: 100.0%

* Equivalent to 2.08 pounds of pyraclostrobin per gallon

EPA Reg. No. 7969-289

EPA Est. No.

U.S. Patent No. 7,816,392

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See full label for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

BASF Agricultural Solutions US LLC
2 TW Alexander Drive
Research Triangle Park, NC 27713

FIRST AID

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after first 5 minutes; then continue rinsing. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Agricultural Solutions US LLC (hereafter "BASF") for emergency medical treatment information: 1-800-832-HELP (4357).

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING. May be fatal if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

This pesticide is toxic to mammals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Protective eyewear (goggles, face shield, or safety glasses)
- Shoes plus socks
- Chemical-resistant gloves made of waterproof material (barrier laminate, butyl rubber ≥ 14 mils, natural rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils)
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, and loading

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product may contaminate water through drift of spray in wind.

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DO NOT apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. Pyraclostrobin is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of pyraclostrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of waterproof material (barrier laminate, butyl rubber \geq 14 mils, natural rubber \geq 14 mils, neoprene rubber \geq 14 mils, nitrile rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils)
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses)

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake

(capacity \leq 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake

(capacity $>$ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(continued)

STORAGE AND DISPOSAL *(continued)*

Container Handling *(continued)*

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with

Headline® SC fungicide or **Headline® fungicide**.

Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF: 1-800-832-HELP (4357)

Steps to take if material is released or spilled:

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

Product Information

Headline SC, a suspension concentrate (SC), contains the active ingredient pyraclostrobin, a member of the **strobilurin class of chemistry** derived from a natural antifungal substance. Preventive applications optimize disease control, resulting in improved plant health. The increase in plant health comes from the combined effect of disease control (including fungal diseases listed in Crop-specific directions), improved growth efficiency and improved stress tolerance. Overall increased plant health may result in an improvement in crop growth and crop quality as well as increased crop yields.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>.

To maximize disease control, apply **Headline SC** in a regularly scheduled protective spray program and use in a rotation program with other fungicides.

Because of its high specific activity, **Headline SC** has good residual activity against target fungi.

Headline SC is not for use in greenhouse or transplant production.

Mode of Action

Pyraclostrobin, the active ingredient of **Headline SC**, belongs to the group of respiration inhibitors classified by the US EPA and Canada PMRA as quinone outside inhibitors (Qol) or target site of action **Group 11** fungicides.

Resistance Management

Headline SC is effective against pathogens resistant to fungicides with modes of action different from those of Qol fungicides (target site **Group 11**), such as dicarboximides, sterol inhibitors, benzimidazoles, or phenylamides.

Fungal isolates resistant to **Group 11** fungicides, such as pyraclostrobin, azoxystrobin, fluoxastrobin, trifloxystrobin, and kresoxim-methyl, can eventually dominate the fungal population if **Group 11** fungicides are used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species. This can result in reduction of disease control by **Headline SC** or other **Group 11** fungicides.

DO NOT exceed the maximum annual use rate or the total number of **Headline SC** applications per year and the maximum number of **Headline SC** applications stated in **Table 1. Headline® SC fungicide Restrictions and Limitations Overview** and **Table 2. Headline® SC fungicide Crop-specific Requirements**. Follow the label instructions for use of **Headline SC** or other target site of action **Group 11** fungicides that have a similar site of action on the same pathogens.

When using a **Group 11** fungicide as a solo product, the number of applications should be no more than 1/3 of the total number of fungicide applications per year.

In programs applying or using tank mixes or pre-mixes of a **Group 11** fungicide with a fungicide of another group, the number of **Group 11** fungicide (QoI)-containing applications must not be more than 1/2 of the total number of fungicide applications per year. In programs applying or using **Group 11** fungicides with both solo products and mixtures, the number of **Group 11** fungicide (QoI)-containing applications must not be more than 1/2 of the total number of fungicide applications per year.

In fungicide alternation programs of **Group 11** (QoI)-containing fungicides with **non-Group 11** fungicides of different modes of action, the maximum number of sequential applications stated in **Table 1. Headline® SC fungicide Restrictions and Limitations Overview** and **Table 2. Headline® SC fungicide Crop-specific Requirements** must be alternated with at least an equal number of applications of a **non-Group 11**-containing fungicide prior to using the **Group 11** (QoI)-containing fungicide again. If two sequential applications of a **Group 11** (QoI)-containing fungicide are made, follow this block of applications with 2 or more applications of a **non-Group 11**-containing fungicide prior to using the **Group 11** (QoI)-containing fungicide again.

For resistance management, **Headline SC** contains a **Group 11** fungicide. Any fungal population may contain individuals naturally resistant to **Headline SC** and other **Group 11** fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Headline SC** or other **Group 11** fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from different target site of action groups that are registered/permited for the same use and that are effective against the pathogens of concern.
- Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Consult your local extension specialist, certified crop advisor and/or BASF representative for additional IPM strategies established for your area. **Headline SC** can be used in agricultural extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.

- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.
- If a **Group 11** target site fungicide, such as **Headline SC**, appears to be less effective against a pathogen that it previously controlled or suppressed, contact a BASF representative, local extension specialist, or certified crop advisor for further investigation.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

Application Instructions

Apply **Headline SC** rates as instructed in **Table 2. Headline® SC fungicide Crop-specific Requirements**. Apply **Headline SC** with ground sprayer, aerial equipment, or through sprinkler irrigation equipment. Check equipment frequently for calibration.

For containers 5 gallons or less, shake well prior to use. For containers greater than 5 gallons, recirculate prior to use. Consult BASF Representatives for additional information regarding agitation and recirculation.

Under low-level disease conditions, use minimum application rates. For severe or threatening disease conditions, use maximum application rates and shortened spray intervals.

Cleaning Spray Equipment

Spray equipment must be cleaned thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used prior to **Headline SC**.

Ground Application

Apply **Headline SC** in sufficient water to ensure thorough coverage of foliage, blooms, and fruit for optimum disease control. Refer to **Additives and Tank Mixing Information** section for adjuvant or crop oil restrictions for ground applications in corn. See **Table 2. Headline® SC fungicide Crop-specific Requirements** for in-furrow instructions.

Aerial Application

For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Unless otherwise specified in this label or in supplemental labeling, use no less than 5 gallons of spray solution per acre. For aerial application to citrus orchards, use no less than 10 gallons of spray solution per acre. **DO NOT** apply when conditions favor drift from target area.

Aerial Application to Alfalfa, Barley, Corn, Oats, Rye, Soybean, Wheat and Triticale

Aerial applications of **Headline® SC fungicide** can be made to corn, soybean, wheat and triticale in water volumes of 1 or more gallons of spray solution per acre (gpa). Aerial applications of **Headline SC** can be made to alfalfa, barley, oats and rye in water volumes of 2 or more gallons of spray solution per acre (gpa). The use of a crop oil or adjuvant can be used to improve spray coverage (see **Additives and Tank Mixing Information** section). Refer to the adjuvant product label for specific use directions and restrictions.

For optimum results in high disease pressure, use a minimum spray volume of 4 gpa. Select spray nozzles, pumping pressure, and sprayer height to provide spray droplets that penetrate throughout the crop canopy. Spray calibration must be conducted to confirm spray droplet sizes. Continue to monitor spray application (including weather conditions) to ensure proper droplet size and canopy penetration.

Spray volumes of 5 gallons or more per acre. For applications with spray volumes of 5 gallons or more per acre, an adjuvant can be used to improve spray coverage (see **Additives and Tank Mixing Information**). Refer to the adjuvant product label for specific use directions and restrictions.

Select spray nozzles, pumping pressure, and sprayer height to provide spray droplets that penetrate throughout the crop canopy. Calibrate spray to confirm spray droplet sizes. Monitor spray application including weather conditions to ensure proper droplet size and canopy penetration.

Aerial Application Methods and Equipment

The interaction of many equipment-related and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

DO NOT apply under circumstances where possible drift to unprotected persons, to food, forage, or other plantings that might be damaged, or crops thereof rendered unfit for sale, use or consumption can occur.

The following drift management requirements must be followed to avoid off-target drift movement to agricultural field crops. These requirements **DO NOT** apply to forestry applications, public health uses, or to applications using dry formulations.

Mandatory Spray Drift Management

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzle and pressure that deliver a medium or coarser droplet size (ASABE S641).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 to 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 ft above the ground or crop canopy.
- Applicators are required to select nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NONTARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.

- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

Boom Height – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Release Height - Aircraft

Higher release heights increase the potential for spray drift.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing. Avoid applications during temperature inversions.

Wind

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

Directions For Use Through Sprinkler Irrigation Systems

Sprayer Preparation

Chemical tank and injector system must be thoroughly cleaned. Flush system with clean water.

Application Instructions

Apply **Headline® SC fungicide** at rates and timings specified in this label.

Sprinkler Irrigation Applications Use Precautions

- Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Add this product to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product/water mixture continuously, applying the labeled rate per acre for that crop. **DO NOT** exceed 1/2 inch (13,577 gallons) per acre. In stationary or non-continuous moving systems, inject the product/water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Thorough coverage of foliage is required for good control. Maintain agitation during the entire application period.
- Contact state extension service specialists, equipment manufacturers, or other experts for calibration questions.
- The system must contain a functional check valve, vacuum-relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide-injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the

responsible person, shall shut the system down and make necessary adjustments should the need arise.

- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Additives and Tank Mixing Information

Headline® SC fungicide can be tank mixed with most recommended fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives as specified in **Table 2. Headline® SC fungicide Crop-specific Requirements.**

Under some conditions, the use of additives or adjuvants may improve the performance of **Headline SC**. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury can result from

mixing **Headline SC** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Follow the most restrictive label.

Adjuvant or Crop Oil Use Limitations on Corn (ground and aerial applications)

Adjuvant crop damage can occur when an adjuvant or crop oil is used after the V8 stage and before the VT stage (the VT stage is defined as when the tassel's last branch is completely visible outside the whorl). If an adjuvant or crop oil is used after the V8 stage and before the VT stage, the grower and user are responsible for contacting the adjuvant source (adjuvant distributor, retailer, or manufacturer) for advice and confirmation that the adjuvant has been tested and proven to be safe for application from V8 to VT corn stage. Refer to adjuvant and/or crop oil labels for specific use directions and restrictions. Always follow the most restrictive label.

Another fungicide or an insecticide may be included in the tank mix if needed and labeled for use on corn. Refer to the tank mix pesticide product labels for specific use directions and restrictions. Always follow the most restrictive label.

Mixing Order

1. **Water** - Agitate a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (such as **Headline SC**, dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) - **For containers 5 gallons or less, shake well prior to use. For containers greater than 5 gallons, recirculate prior to use.** Consult BASF Representatives for additional information regarding agitation and recirculation.
6. **Water-soluble products**
7. **Emulsifiable concentrates** (such as oil concentrates when applicable)
8. **Water-soluble additives** (such as ammonium sulfate [AMS] or urea ammonium nitrate [UAN] when applicable)
9. **Remaining quantity of water**

Make sure each component is thoroughly mixed and suspended before adding tank mix partners. Maintain constant agitation during application. See **Table 2. Headline® SC fungicide Crop-specific Requirements** for more details.

Restrictions and Limitations - All Crops

- **DO NOT** exceed the maximum product rate (fl ozs/A) per year, the maximum rate per application, or the total number of applications of **Headline® SC fungicide** per year as stated in **Table 1. Headline® SC fungicide Restrictions and Limitations Overview** and **Table 2. Headline® SC fungicide Crop-specific Requirements**. Preharvest interval (PHI) restrictions are also included in these tables.
- **DO NOT** use **Headline SC** in greenhouse or transplant production.
- **For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).**

Crop Rotation Restriction

Crops listed on the **Headline SC, Cabrio® EG fungicide** and **Pristine® fungicide** labels may be planted immediately following the last application. For all other crops, **DO NOT** plant sooner than 14 days after the last application.

Ground Application Directed or Banded Sprays

The application rates shown in the following tables pertain to both aerial and ground (broadcast) methods of application. **Headline SC** may also be applied as a directed or banded spray over the rows or plant beds with alleys or row middles left unsprayed. For such uses, reduce the **Headline SC** rate in proportion to the area actually sprayed. This adjustment is necessary to prevent applying the product at use rates higher than permitted on this label.

Use the following formula to determine the broadcast equivalent rate for directed or banded sprays:

sprayed bed width + unsprayed row middles = total row width

$$\frac{\text{Sprayed Bed Width in Inches}}{\text{Total Row Width in Inches}} \times \frac{\text{Broadcast Rate}}{\text{Treated Acre}} = \frac{\text{Band Rate}}{\text{Field Acre}}$$

EXAMPLE: Directed spray application to 45-inch plant beds separated by 15-inch unsprayed row-middles at a 12 fl ozs/A label broadcast rate:

45 inches sprayed bed width + 15 inches unsprayed row middles = 60 inches total row width

$$\frac{45 \text{ Inches Sprayed Bed Width}}{60 \text{ Inches Total Row Width}} \times \frac{12 \text{ fl ozs Headline SC}}{\text{Treated Acre}} = \frac{9 \text{ fl ozs Headline SC}}{\text{Field Acre}}$$

Table 1. Headline® SC fungicide Restrictions and Limitations Overview¹

Crop/Crop Group²	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A) (lbs ai pyraclostrobin)
Alfalfa³	14	9	3	27 (0.45)
Barley	See Table 2. Headline® SC fungicide Crop-specific Requirements	9	2	18 (0.29)
Citrus Fruits Group⁴	0	15	2	54 (0.88)
Corn^{5,7} (all types)	7	12	2	72 (1.18)
Cotton⁵	30	12	2	36 (0.58)
Dried Shelled Peas and Beans^{5,7} (except soybeans)	21	9	2	18 (0.29)
Edible-podded Legume Vegetables	7	9	2	18 (0.29)
Grass Grown for Seed	14	12	2	24 (0.39)
Mint	14	12	2	48 (0.78)
Oats	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59)	9	2	18 (0.29)
Oilseed Crops⁵	21	12	2	24 (0.39)
Peanut^{5,7}	14	15	2	45 (0.73)
Pecan	14	7	2	28 (0.46)
Rye	Apply no later than 50% head emergence (Feekes 10.3, Zadok's 55)	9	2	18 (0.29)
Sorghum	Apply no later than 25% flowering	12	1	12 (0.20)
Soybean^{5,7}	21	12	2	24 (0.39)
Succulent Shelled Peas and Beans	7	9	2	18 (0.29)
Sugar Beet⁵ (roots and tops)	7	12	2	48 (0.78)
Sugarcane⁶	14	12	2	48 (0.78)
Tuberous and Corm Vegetables Subgroup⁵ (includes potato)	3	12	1	72 (1.18)

(continued)

Table 1. Headline® SC fungicide Restrictions and Limitations Overview¹ *(continued)*

Crop/Crop Group ²	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A) <i>(lb ai pyraclostrobin)</i>
Wheat and Triticale	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59)	9	2	18 <i>(0.29)</i>

¹ See **Table 2. Headline® SC fungicide Crop-specific Requirements** for complete directions and exceptions.

² For a complete list of crops within a crop group, see **Table 2. Headline® SC fungicide Crop-specific Requirements**.

³ **DO NOT** apply more than 27 fl ozs/A (0.45 lb ai/acre) of **Headline SC** in alfalfa per year.

⁴ Maximum product rate per acre per application may vary for citrus fruits depending on target disease. Refer to **Table 2. Headline® SC fungicide Crop-specific Requirements, Citrus Fruits** for maximum rates per application by target disease.

⁵ The maximum product rate per year includes the combination in-furrow and foliar uses.

⁶ Not for use on sugarcane in California.

⁷ Not for in-furrow use in California.

Aerial application is permitted for all labeled crops. **For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).**

Table 2. Headline® SC fungicide Crop-specific Requirements

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Alfalfa	Anthracnose <i>Colletotrichum trifolii</i> Common leaf spot <i>Pseudopeziza medicaginis</i> Downy mildew <i>Peronospora trifoliorum</i> Leaf spot <i>Leptosphaerulina briosiani</i> Powdery mildew <i>Erysiphe pisi</i> Rhizoctonia blight/black patch <i>Rhizoctonia</i> spp. Rust <i>Uromyces</i> spp. Spring black stem and leaf spot <i>Phoma medicaginis</i> Stagnospora leaf spot <i>Stagnospora melliloti</i> Stemphyllium leaf spot <i>Stemphyllium</i> spp. Summer black stem and leaf spot <i>Cercospora medicaginis</i> Yellow leaf blotch <i>Leptotrichila medicaginis</i>	6 to 9	3	27 (0.45 lb ai/acre)	14

Application Directions. For optimal disease control, begin **Headline SC** applications prior to disease development.

Resistance Management. DO NOT make more than three (3) **Headline SC** applications per year.

Repeat applications on a 14 to 21 day interval if conditions are conducive for disease development. **DO NOT** make more than two (2) **Headline SC** applications per cutting or three (3) **Headline SC** applications per year. Use the high-rate and shorter interval when disease pressure is high.

* **DO NOT** apply more than 27 fl ozs/A (0.45 lb ai/acre) of **Headline SC** in alfalfa per year.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Barley	Black point Kernel blight or Head mold <i>Cochliobolus sativus</i> , <i>Alternaria</i> spp. Leaf rust <i>Puccinia hordei</i> , <i>P. recondita</i> Net blotch <i>Pyrenophora teres</i> Powdery mildew <i>Erysiphe graminis</i> f. sp., <i>hordei</i> Scald <i>Rhynchosporium secalis</i> Septoria leaf and glume blotch <i>Septoria</i> spp., <i>Stagonospora</i> spp. Spot blotch <i>Cochliobolus sativus</i> Stem rust <i>Puccinia graminis</i> f. sp., <i>tritici</i> Stripe rust <i>Puccinia striiformis</i> Tan spot Yellow leaf spot <i>Pyrenophora trichostoma</i>	6 to 9*	2	18 (0.29 lb ai/acre)	Apply no later than 50% head emergence (Feekes 10.3, Zadok's 55); 14 days in selected states (see map).

Application Directions. Begin **Headline SC** applications prior to disease development. To maximize yields in cereals, protect the flag leaf. Apply **Headline SC** immediately after flag-leaf emergence for optimum results.

Headline SC does not control Fusarium head blight (head scab) or prevent the reductions in grain quality that can result from this disease. When head blight is a concern, manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.29 lb ai pyraclostrobin (18 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

DO NOT harvest barley hay or feed green-chopped barley within 14 days of last application.

* For early season control of net blotch, Septoria leaf and glume blotch, spot blotch, and tan spot when conditions favor disease development, apply 3 to 6 fl ozs per acre of **Headline SC** either in combination with a herbicide application or when conditions favor disease development. When the 3 to 6 fl ozs early season application rate is used, a second application of **Headline SC** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Headline SC** rate for the second application. For high disease pressure, use the higher rate of **Headline SC**. Early season control is not registered for use in California.

Table 2. Headline® SC fungicide Crop-specific Requirements *(continued)*

Application Directions *(continued)*

Barley can be harvested 14 days after the last application in the following states: **AZ** (north of I-10), **CO, ID, MT** (west of Rt 87/I-15), **NV, NM, OR, TX** (west of Rt 283/377), **UT, WA**, and **WY** (west of I-25/I-90), as shown in the **Headline SC Use Area Map - Barley, 14-day PHI**.

**Headline SC Use Area Map - Barley
14-day PHI**




Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Citrus Fruits Group Australian desert lime Australian finger lime Australian round lime	Greasy spot <i>Mycosphaerella citri</i>	9 to 12	2	54 (0.88 lb ai/acre)	0
	Scab <i>Elsinoe</i> spp.				
Brown River finger lime Calamondin Citron Citrus hybrids Chironja Grapefruit Japanese summer grapefruit Kumquat Lemon Lime Mediterranean mandarin Mount white lime New guinea wild lime Orange, sour Orange, sweet Pummelo Russell River lime Satsuma mandarin Sweet lime Tachibana orange Tahiti lime Tangelo Tangerine (mandarin) Tangor Trifoliate orange Uniq fruit Cultivars, varieties and/or hybrids of these	Alternaria brown spot <i>Alternaria citri</i> Anthracnose <i>Colletotrichum acutatum</i> , <i>C. gloeosporioides</i> Black spot <i>Guignardia citricarpa</i> Melanose <i>Diaporthe citri</i> Post bloom fruit drop <i>Colletotrichum acutatum</i>	12 to 15			

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 10- to 21-day interval.

Use the higher rate when disease pressure is high.

For control of diseases other than greasy spot, integrate 1 to 2 **Headline SC** applications early in the spray program. **For greasy spot control,** integrate 1 to 2 **Headline SC** applications into the fungicide program during the mid-to-late season.

For aerial application to citrus orchards, use no less than 10 gallons of spray solution per acre.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.88 lb ai pyraclostrobin (54 fl ozs of **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Corn Field corn Popcorn Sweet corn Seed production corn	Anthracnose** <i>Colletotrichum graminicola</i> Eyespot <i>Kabatiella zeae</i> Gray leaf spot <i>Cercospora zea-maydis</i> Northern corn leaf blight** <i>Exserohilum turcicum</i> Northern corn leaf spot** <i>Cochliobolus carbonum</i> Physoderma brown spot** <i>Physoderma maydis</i> Rust, common <i>Puccinia sorghi</i> Rust, Southern <i>Puccinia polyspora</i> Southern corn leaf blight** <i>Bipolaris maydis</i> Yellow leaf blight** <i>Phyllosticta maydis</i>	6 to 12	2	72 (1.18 lbs ai/acre)	7

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high. Under high disease pressure for Northern corn leaf blight and Southern corn leaf blight, apply 9 to 12 fl ozs per acre.

Headline SC can be used with adjuvants in corn. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 1.18 lbs ai pyraclostrobin (72 fl ozs of **Headline SC**) per acre per year.

In field corn, DO NOT make more than two (2) **Headline SC** applications per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action. If more than two (2) **Headline SC** applications are made in a multiple spray program, alternate each subsequent **Headline SC** application with at least one (1) application of a **non-Group 11** fungicide.

* The maximum product rate per year includes the combination of in-furrow and foliar uses.

** The use rate in California is 9 to 12 fl ozs per acre.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne Rhizoctonia and the Suppression of Soilborne Fusarium Species in Corn*									
Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)								
	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.1	3.5								
0.2	7.0	5.2	4.7	3.5	3.3	3.2	3.0		
0.3	10.5	7.8	7.1	5.2	5.0	4.8	4.5	4.3	4.0
0.4	see footnote ¹	10.4	9.5	6.9	6.7	6.4	6.0	5.7	5.4
0.5	see footnote ¹	see footnote ¹	11.8	8.7	8.4	8.0	7.5	7.1	6.7
0.6	see footnote ¹	see footnote ¹	see footnote ¹	10.4	10.0	9.6	9.0	8.5	8.1
0.7	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	11.7	11.2	10.5	10.0	9.4
0.8	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	12.0	11.4	10.8

Application Directions. Use 0.1 to 0.8 fl oz of **Headline SC** per 1000 row feet. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered. Use a minimum application volume of 2.5 gallons of water per acre.

When Rhizoctonia or Fusarium seedling disease pressure conditions are expected to be severe or if the field has a history of seedling diseases, use **Headline SC** at a product rate per acre equivalent to 9 to 12 fl ozs and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre in-furrow of **Headline SC**.

* Not for in-furrow use in California.

¹ For 32- to 34-inch rows, use a maximum of 0.7 fl oz per 1000 row feet.
 For 30-inch rows, use a maximum of 0.6 fl oz per 1000 row feet.
 For 22-inch rows, use a maximum of 0.5 fl oz per 1000 row feet.
 For 20-inch rows, use a maximum of 0.4 fl oz per 1000 row feet.
 For 15-inch rows, use a maximum of 0.3 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Cotton	Alternaria leaf spot, boll rot <i>Alternaria</i> spp. Anthracnose, boll rot <i>Glomerella</i> spp. Ascochyta blight, boll rot <i>Ascochyta</i> spp. Cercospora blight and leaf spot <i>Cercospora</i> spp. Diplodia boll rot <i>Diplodia</i> spp. Hard lock, boll rot <i>Fusarium</i> spp. Phoma blight, boll rot <i>Phoma</i> spp. Rust <i>Puccinia</i> spp., <i>Phykopsora</i> spp. Stemphyllium leaf spot <i>Stemphyllium</i> spp.	6 to 12	2	36 (0.58 lb ai/acre)	30

Application Directions. For foliar and boll rot disease control, begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high. For seedling disease control, see in-furrow application instructions following.

Headline SC can be used with adjuvants in cotton. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

No livestock grazing or feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.58 lb ai pyraclostrobin (36 fl ozs of **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

*The maximum product rate per year includes the combination of in-furrow and foliar uses.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne Rhizoctonia in Cotton									
Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)								
	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.1	3.5								
0.2	7.0	5.2	4.7	3.5	3.3	3.2	3.0		
0.3	10.5	7.8	7.1	5.2	5.0	4.8	4.5	4.3	4.0
0.4	see footnote ¹	10.4	9.5	6.9	6.7	6.4	6.0	5.7	5.4
0.5	see footnote ¹	see footnote ¹	11.8	8.7	8.4	8.0	7.5	7.1	6.7
0.6	see footnote ¹	see footnote ¹	see footnote ¹	10.4	10.0	9.6	9.0	8.5	8.1
0.7	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	11.7	11.2	10.5	10.0	9.4
0.8	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	12.0	11.4	10.8

Application Directions. Use 0.1 to 0.8 fl oz of **Headline SC** per 1000 row feet. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered. Use a minimum application volume of 2.5 gallons of water per acre.

When Rhizoctonia seedling disease pressure conditions are expected to be severe or if the field has a history of seedling diseases, use **Headline SC** at a product rate per acre equivalent to 9 to 12 fl ozs and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre of **Headline SC**.

¹ For 32- to 34-inch rows, use a maximum of 0.7 fl oz per 1000 row feet.
 For 30-inch rows, use a maximum of 0.6 fl oz per 1000 row feet.
 For 22-inch rows, use a maximum of 0.5 fl oz per 1000 row feet.
 For 20-inch rows, use a maximum of 0.4 fl oz per 1000 row feet.
 For 15-inch rows, use a maximum of 0.3 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Dried Shelled Peas and Beans (except soybeans)</p> <p>Broad bean Chickpea Guar Lablab bean Lentil Pigeon pea</p> <p>Lupinus spp.</p> <p>Grain lupin Sweet lupin White lupin</p> <p>Phaseolus spp.</p> <p>Field bean Kidney bean Lima bean Navy bean Pink bean Pinto bean Tepary bean</p> <p>Vigna spp.</p> <p>Adzuki bean Black-eyed pea Catjang Cowpea Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean</p> <p>Pisum spp.</p> <p>Field pea</p>	<p>Anthracnose <i>Colletotrichum</i> spp.</p> <p>Alternaria leaf and pod spot <i>Alternaria</i> spp.</p> <p>Ascochyta blight <i>Phoma exigua</i>, <i>Ascochyta</i> spp.</p> <p>Asian soybean rust <i>Phakopsora pachyrhizi</i></p> <p>Cercospora leaf spot <i>Cercospora</i> spp.</p> <p>Downy mildew <i>Phytophthora nicotianae</i></p> <p>Mycosphaerella blight <i>Mycosphaerella</i> spp.</p> <p>Powdery mildew <i>Erysiphe polygoni</i></p> <p>Rust <i>Uromyces appendiculatus</i></p>	<p>6 to 9</p>	<p>2</p>	<p>18 (0.29 lb ai/acre)</p>	<p>21</p>

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Bean forage, bean hay, pea vines, and pea hay may be fed no sooner than 14 days after last application.

Headline SC can be used with adjuvants in dried shelled peas and beans (except soybean). See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.29 lb ai pyraclostrobin (18 fl ozs of **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

* The maximum product rate per year includes the combination of in-furrow and foliar uses.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne Rhizoctonia and the Suppression of Soilborne Fusarium Species in Dried Shelled Peas and Beans** (except soybeans)									
Rate Per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)								
	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.1	3.5								
0.2	7.0	5.2	4.7	3.5	3.3	3.2	3.0		
0.3	see footnote ¹	7.8	7.1	5.2	5.0	4.8	4.5	4.3	4.0
0.4	see footnote ¹	see footnote ¹	see footnote ¹	6.9	6.7	6.4	6.0	5.7	5.4
0.5	see footnote ¹	see footnote ¹	see footnote ¹	8.7	8.4	8.0	7.5	7.1	6.7
0.6	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	9.0	8.5	8.1

Application Directions. Use 0.1 to 0.6 fl oz of **Headline SC** per 1000 row feet. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered. Use a minimum application volume of 2.5 gallons of water per acre.

When Rhizoctonia or Fusarium seedling disease pressure conditions are expected to be severe or if the field has a history of seedling diseases, use **Headline SC** at a product rate per acre equivalent to 9 fl ozs and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 9 fl ozs per acre in-furrow of **Headline SC**.

* Adzuki bean, black-eyed pea, broad bean, catjang, chickpea, crowder pea, field bean, grain lupin, guar, kidney bean, lablab bean, lima bean, moth bean, mung bean, navy bean, pink bean, pinto bean, rice bean, Southern pea, sweet lupin, tepary bean, urd bean, and white lupin

** Not for in-furrow use in California.

¹ For 30- to 34-inch rows, use a maximum of 0.5 fl oz per 1000 row feet.
 For 20- to 22-inch rows, use a maximum of 0.3 fl oz per 1000 row feet.
 For 15-inch rows, use a maximum of 0.2 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Edible-podded Legume Vegetables Jack bean Pigeon pea Soybean (immature seed) Sword bean Phaseolus spp. Runner bean Snap bean Wax bean Vigna spp. Asparagus bean Chinese longbean Moth bean Yardlong bean Pisum spp. Dwarf pea Edible-podded pea Snowpea Sugar snap pea	Alternaria leaf and pod spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> spp. Ascochyta blight <i>Phoma exigua</i> , <i>Ascochyta</i> spp. Asian soybean rust <i>Phakopsora pachyrhizi</i> Cercospora leaf spot <i>Cercospora</i> spp. Downy mildew <i>Phytophthora nicotianae</i> Mycosphaerella blight <i>Mycosphaerella</i> spp. Powdery mildew <i>Erysiphe Polygoni</i> Rust <i>Uromyces appendiculatus</i>	6 to 9	2	18 (0.29 lb ai/acre)	7

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Bean forage, bean hay, pea vines, and pea hay maybe fed no sooner than 14 days after last application.

Headline SC can be used with adjuvants in edible-podded legume vegetables. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.29 lb ai pyraclostrobin (18 fl ozs of **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Grass Grown for Seed	Rust <i>Puccinia recondita</i> , <i>P. graminis</i> Suppression Only: Powdery mildew <i>Erysiphe graminis</i>	6 to 12	2	24 (0.39 lb ai/acre)	14

Application Directions. Begin **Headline SC** applications prior to disease development. Apply again 14 to 21 days later.

Use the higher rate and shorter interval when disease pressure is high.

DO NOT graze or feed forage or hay to livestock within 27 days of last application.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.39 lb ai pyraclostrobin (24 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Mint	Leaf spot <i>Ramularia</i> spp., <i>Alternaria</i> spp., <i>Phoma</i> spp. Powdery mildew <i>Erysiphe</i> spp. Rust <i>Puccinia</i> spp.	9 to 12	2	48 (0.78 lb ai/acre)	14

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Headline SC can be used with adjuvants in mint. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.78 lb ai pyraclostrobin (48 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oats	Crown rust <i>Puccinia coronata</i> Helminthosporium leaf spot <i>Drechslera avenae</i> Leaf blotch <i>Pyrenophora avenae</i> Leaf rust <i>Puccinia</i> spp. Septoria blotch and stem rot <i>Septoria avenae</i> , <i>Phaeosphaeria avenaria</i> , <i>Stagnospora avenae</i> Spot blotch <i>Bipolaris</i> spp. Stem rust <i>Puccinia graminis</i>	6 to 9*	2	18 (0.29 lb ai/acre)	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59)

Application Directions. Begin **Headline SC** applications prior to disease development. To maximize yields in cereals, protect the flag leaf. Apply **Headline SC** immediately after flag-leaf emergence for optimum results.

Headline SC does not control Fusarium head blight (head scab) or prevent reductions in grain quality that can result from this disease. When head blight is a concern, manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

DO NOT harvest oat hay or feed green-chopped oats within 14 days of last application.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.29 lb ai pyraclostrobin (18 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

* For early season control of leaf blotch, Septoria blotch and stem rot, and spot blotch when conditions favor disease development, apply 3 to 6 fl ozs per acre of **Headline SC** either in combination with a herbicide application or when conditions favor disease development. When the 3 to 6 fl ozs early season application rate is used, a second application of **Headline SC** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Headline SC** rate for the second application. For high disease pressure, use the higher rate of **Headline SC**. Early season control is not registered for use in California.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oilseed Crops Flax seed	Pasmo <i>Septoria linicola</i>	6 to 12	2	24 (0.39 lb ai/acre)	21
Rapeseed	Blackleg <i>Leptosphaeria maculans</i>				
	Blackspot <i>Alternaria</i> spp.				
Sunflower	Alternaria leaf spot <i>Alternaria</i> spp. Cercospora leaf spot <i>Cercospora helianthi</i> Downy mildew <i>Plasmopara halstedii</i> Powdery mildew <i>Erysiphe cichoracearum</i> Rust <i>Puccinia helianthi</i> , <i>Uromyces</i> spp. Septoria leaf spot <i>Septoria</i> spp. White rust <i>Albugo tragopogonis</i>				

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Flax seed. Apply **Headline SC** at mid-flowering (7 to 10 days after flower initiation). Make a second application 7 to 10 days later if disease persists or if weather conditions are favorable for disease development.

Rapeseed. For control of blackleg, apply **Headline SC** at 2- to 4-leaf stage. **For control of blackspot,** apply **Headline SC** at early pod development. A second application 7 to 10 days later can be made if disease persists or if weather conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Headline SC can be used with adjuvants in oilseed crops. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.39 lb ai pyraclostrobin (24 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a **non-Group 11** fungicide with a different mode of action.

*The maximum product rate per year for **sunflower** includes the combination of in-furrow and foliar uses.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oilseed Crops (continued) Borage Calendula Castor oil plant Chinese tallowtree Crambe Cuphea Echium Euphorbia Evening primrose Gold of pleasure (Camelina) Hare's ear mustard Jojoba Lesquerella Lunaria Meadowfoam Milkweed Mustard seed Niger seed Oil radish Poppy seed Rose hip Safflower Sesame Stokes aster Sweet rocket Tallowwood Tea oil plant Vernonia	<i>Alternaria</i> spp. <i>Septoria</i> spp.	6 to 12	2	24 (0.39 lb ai/acre)	21

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Headline SC can be used with adjuvants in oilseed crops. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.39 lb ai pyraclostrobin (24 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a **non-Group 11** fungicide with a different mode of action.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne Rhizoctonia in Sunflower									
Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)								
	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.1	3.5								
0.2	7.0	5.2	4.7	3.5	3.3	3.2	3.0		
0.3	10.5	7.8	7.1	5.2	5.0	4.8	4.5	4.3	4.0
0.4	see footnote ¹	10.4	9.5	6.9	6.7	6.4	6.0	5.7	5.4
0.5	see footnote ¹	see footnote ¹	11.8	8.7	8.4	8.0	7.5	7.1	6.7
0.6	see footnote ¹	see footnote ¹	see footnote ¹	10.4	10.0	9.6	9.0	8.5	8.1
0.7	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	11.7	11.2	10.5	10.0	9.4
0.8	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	12.0	11.4	10.8

Application Directions. Use 0.1 to 0.8 fl oz of **Headline SC** per 1000 row feet. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered. Use a minimum application volume of 2.5 gallons of water per acre.

When Rhizoctonia seedling disease pressure conditions are expected to be severe or if the field has a history of seedling diseases, use **Headline SC** at a product rate per acre equivalent to 9 to 12 fl ozs and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre of **Headline SC**.

¹ For 32- to 34-inch rows, use a maximum of 0.7 fl oz per 1000 row feet.
 For 30-inch rows, use a maximum of 0.6 fl oz per 1000 row feet.
 For 22-inch rows, use a maximum of 0.5 fl oz per 1000 row feet.
 For 20-inch rows, use a maximum of 0.4 fl oz per 1000 row feet.
 For 15-inch rows, use a maximum of 0.3 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Peanut	Early leaf spot <i>Cercospora arachidicola</i>	6 to 15 (see details below)	2	45 (0.73 lbs ai/acre)	14
	Late leaf spot <i>Cercosporidium personatum</i>				
	Pepperspot <i>Leptosphaerulina crassiasca</i>				
Rust <i>Puccinia arachidis</i>					
Web blotch <i>Phoma arachidicola</i>					
Rhizoctonia limb rot, Peg rot, Pod rot <i>Rhizoctonia solani</i>	9 to 15				
Sclerotium rot, Southern stem rot, Southern blight, and White mold <i>Sclerotium rolfsii</i>					
Suppression Only: Sclerotinia blight <i>Sclerotinia minor</i>					
Cylindrocladium black rot <i>Cylindrocladium crotalariae</i>	12 to 15				

Application Directions. For control of early and late leaf spot, pepperspot, rust, and web blotch, begin **Headline SC** applications prior to disease development and continue on a 14- to 21-day interval. When using a 14-day spray interval, apply **Headline SC** at 6 to 12 fluid ounces per acre. At spray intervals between 14 and 21 days, apply **Headline SC** at 9 to 15 fluid ounces per acre.

For control of Rhizoctonia and Sclerotium, begin **Headline SC** applications prior to disease development and continue on a 14- to 28-day interval. For intervals greater than 14 days, use 15 fluid ounces per acre.

Use the higher rate and/or shorter spray interval when disease pressure is high or in fields with a history of disease.

Headline SC can be used with adjuvants in peanut; however, mixes with silicone-containing adjuvants may cause crop injury under certain conditions. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Peanut meal can be fed. **DO NOT** graze or harvest for forage use.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.73 lbs ai pyraclostrobin (45 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action. In spray programs where four (4) or less fungicide applications are made in a year, **Headline SC** should be alternated with at least one (1) application of a labeled **non-Group 11** fungicide with a different mode of action.

*The maximum product rate per year includes the combination of in-furrow and foliar uses.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne Rhizoctonia and the Suppression of Soilborne Fusarium Species in Peanut*									
Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)								
	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.1	3.5								
0.2	7.0	5.2	4.7	3.5	3.3	3.2	3.0		
0.3	10.5	7.8	7.1	5.2	5.0	4.8	4.5	4.3	4.0
0.4	see footnote ¹	10.4	9.5	6.9	6.7	6.4	6.0	5.7	5.4
0.5	see footnote ¹	see footnote ¹	11.8	8.7	8.4	8.0	7.5	7.1	6.7
0.6	see footnote ¹	see footnote ¹	see footnote ¹	10.4	10.0	9.6	9.0	8.5	8.1
0.7	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	11.7	11.2	10.5	10.0	9.4
0.8	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	12.0	11.4	10.8

Application Directions. Use 0.1 to 0.8 fl oz of **Headline SC** per 1000 row feet. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered. Use a minimum application volume of 2.5 gallons of water per acre.

When Rhizoctonia or Fusarium seedling disease pressure conditions are expected to be severe or if the field has a history of seedling diseases, use **Headline SC** at a product rate per acre equivalent to 9 to 12 fl ozs and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre of **Headline SC** for peanut in-furrow use.

* Not for in-furrow use in California.

¹ For 32- to 34-inch rows, use a maximum of 0.7 fl oz per 1000 row feet.
 For 30-inch rows, use a maximum of 0.6 fl oz per 1000 row feet.
 For 22-inch rows, use a maximum of 0.5 fl oz per 1000 row feet.
 For 20-inch rows, use a maximum of 0.4 fl oz per 1000 row feet.
 For 15-inch rows, use a maximum of 0.3 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements *(continued)*

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Pecan	Pecan scab <i>Cladosporium caryigenum</i>	6 to 7	2	28 <i>(0.46 lb ai/acre)</i>	14

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 14-day interval. For optimum performance, apply **Headline SC** early in the spray program (e.g. prepollination and first cover).

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.46 lb ai pyraclostrobin (28 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Potato	Black dot <i>Colletotrichum coccodes</i>	6 to 9	1	72 (1.18 lbs ai/acre)	3
	Early blight <i>Alternaria solani</i>				
	Late blight <i>Phytophthora infestans</i>	6 to 12			
	Powdery mildew <i>Erysiphe</i> spp., <i>Leveillula taurica</i>				
	Suppression Only: White mold <i>Sclerotinia sclerotiorum</i>				

Application Directions. Begin **Headline SC** applications at 7- to 14-day intervals prior to disease development. The low rate and longer interval can be used early season prior to the observance of symptoms and when disease pressure is low. **For control of late blight**, follow **Headline SC** application with a labeled **non-Group 11** fungicide with a different mode of action 5 to 7 days later.

Use the higher rates and shorter intervals once disease has been confirmed in your area or weather conditions are favorable to disease development.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 1.18 lbs ai pyraclostrobin (72 fl ozs **Headline SC**) per acre per year.

DO NOT make more than one (1) **Headline SC** application before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

*The maximum product rate per year includes the combination of in-furrow and foliar uses.

In-furrow Use to Aid in the Control of Soilborne Rhizoctonia in Potatoes

Use 0.4 to 0.8 fl oz of **Headline SC** per 1000 row feet. For applications on 32-inch or 34-inch rows, the maximum application rate is 0.73 fl oz/1000 row feet. Apply at planting as an in-furrow spray by directing spray pattern to uniformly cover seed pieces and surrounding soil. The spray pattern must be a 4- to 8-inch band applied to the seed piece prior to being covered with soil.

When Rhizoctonia disease pressure conditions are expected to be severe or if the field has a history of Rhizoctonia infestations, use **Headline SC** at 0.6 to 0.8 fl oz per 1000 row feet and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre of **Headline SC**.

Use a minimum volume of application of 5 gallons of water per acre.

Headline SC Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)				
	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.4	6.7	6.4	6.0	5.7	5.4
0.6	10.0	9.6	9.0	8.6	8.1
0.8	see footnote ¹	see footnote ¹	12.0	11.4	10.8

¹ For 32-inch or 34-inch rows, use a maximum of 0.73 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Rye	Leaf rust <i>Puccinia recondita</i> Leaf spot <i>Pyrenophora</i> spp. Powdery mildew <i>Erysiphe graminis</i> Septoria leaf and glume blotch <i>Septoria</i> spp., <i>Stagonospora</i> spp. Stem rust <i>Puccinia graminis</i> Stripe rust <i>Puccinia striiformis</i>	6 to 9*	2	18 (0.29 lb ai/acre)	Apply no later than 50% head emergence (Feekes 10.3, Zadok's 55)

Application Directions. Begin **Headline SC** applications prior to disease development. To maximize yields in cereals, protect the flag leaf. Apply **Headline SC** immediately after flag-leaf emergence for optimum results.

Headline SC does not control Fusarium head blight (head scab) or prevent the reductions in grain quality that can result from this disease. When head blight is a concern, manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.29 lb ai pyraclostrobin (18 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

* For early season control of leaf spot and Septoria leaf and glume blotch when conditions favor disease development, apply 3 to 6 fl ozs per acre of **Headline SC** either in combination with a herbicide application or when conditions favor disease development. When the 3 to 6 fl ozs early season application rate is used, a second application of **Headline SC** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Headline SC** rate for the second application. For high disease pressure, use the higher rate of **Headline SC**. Early season control is not registered for use in California.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sorghum	Anthracnose <i>Colletotrichum graminicola</i> Gray leaf spot <i>Cercospora</i> spp. Northern leaf blight <i>Exserohilum turcicum</i> Rust <i>Puccinia</i> spp. Southern leaf blight <i>Bipolaris</i> spp.	6 to 12	1	12 (0.20 lb ai/acre)	Apply no later than 25% flowering

Application Directions. Begin **Headline SC** applications prior to disease development. Use the higher rate when disease pressure is high. Under high disease pressure for Northern leaf blight and Southern leaf blight, apply 9 to 12 fl ozs per acre.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.20 lb ai/acre pyraclostrobin (12 fl ozs **Headline SC**) per acre per year.

DO NOT make more than one (1) **Headline SC** application per year. If additional fungicide applications are needed, use a labeled **non-Group 11** fungicide with a different mode of action.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Soybean	Alternaria leaf spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum truncatum</i> Asian soybean rust <i>Phakopsora pachyrhizi</i> Brown spot <i>Septoria glycines</i> Cercospora blight <i>Cercospora kikuchii</i> Frogeye leaf spot <i>Cercospora sojina</i> Pod and stem blight <i>Diaporthe phaseolorum</i> Rhizoctonia aerial blight <i>Rhizoctonia solani</i>	6 to 12	2	24 (0.39 lb ai/acre)	21
	Suppression Only: Southern blight <i>Sclerotium rolfsii</i>	12			

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high.

For control of soybean rust, apply **Headline SC** prior to infection.

Headline SC can be used with adjuvants in soybeans. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Soybean forage can be fed no sooner than 14 days after last application.

Soybean hay can be fed no sooner than 21 days after last treatment.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.39 lb ai pyraclostrobin (24 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

*The maximum product rate per year includes the combination of in-furrow and foliar uses.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne Rhizoctonia and the Suppression of Soilborne Fusarium Species in Soybean*									
Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)								
	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.1	3.5								
0.2	7.0	5.2	4.7	3.5	3.3	3.2	3.0		
0.3	10.5	7.8	7.1	5.2	5.0	4.8	4.5	4.3	4.0
0.4	see footnote ¹	10.4	9.5	6.9	6.7	6.4	6.0	5.7	5.4
0.5	see footnote ¹	see footnote ¹	11.8	8.7	8.4	8.0	7.5	7.1	6.7
0.6	see footnote ¹	see footnote ¹	see footnote ¹	10.4	10.0	9.6	9.0	8.5	8.1
0.7	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	11.7	11.2	10.5	10.0	9.4
0.8	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	12.0	11.4	10.8

Application Directions. Use 0.1 to 0.8 fl oz of **Headline SC** per 1000 row feet. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered. Use a minimum application volume of 2.5 gallons of water per acre.

When Rhizoctonia or Fusarium seedling disease pressure conditions are expected to be severe or if the field has a history of seedling diseases, use **Headline SC** at a product rate per acre equivalent to 9 to 12 fl ozs and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre in-furrow of **Headline SC**.

* Not for in-furrow use in California.

¹ For 32- to 34-inch rows, use a maximum of 0.7 fl oz per 1000 row feet.
 For 30-inch rows, use a maximum of 0.6 fl oz per 1000 row feet.
 For 22-inch rows, use a maximum of 0.5 fl oz per 1000 row feet.
 For 20-inch rows, use a maximum of 0.4 fl oz per 1000 row feet.
 For 15-inch rows, use a maximum of 0.3 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Succulent Shelled Peas and Beans</p> <p>Pigeon pea</p> <p>Vigna spp.</p> <p>Black-eyed pea</p> <p>Cowpea</p> <p>Southern pea</p> <p>Pisum spp.</p> <p>Broad bean</p> <p>English pea</p> <p>Garden pea</p> <p>Green pea</p> <p>Phaseolus spp.</p> <p>Lima bean, green</p>	<p>Alternaria leaf and pod spot <i>Alternaria</i> spp.</p> <p>Anthracnose <i>Colletotrichum</i> spp.</p> <p>Ascochyta blight <i>Phoma exigua</i>, <i>Ascochyta</i> spp.</p> <p>Asian soybean rust <i>Phakopsora pachyrhizi</i></p> <p>Cercospora leaf spot <i>Cercospora</i> spp.</p> <p>Downy mildew <i>Phytophthora nicotianae</i>, <i>P. phaseoli</i></p> <p>Mycosphaerella blight <i>Mycosphaerella</i> spp.</p> <p>Powdery mildew <i>Erysiphe polygoni</i></p> <p>Rust <i>Uromyces appendiculatus</i></p>	<p>6 to 9</p>	<p>2</p>	<p>18 (0.29 lb ai/acre)</p>	<p>7</p>

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 7- to 14-day interval if conditions are favorable for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Bean forage, bean hay, pea vines, and pea hay may be fed no sooner than 14 days after last application.

Headline SC can be used with adjuvants in succulent shelled peas and beans. See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.29 lb ai pyraclostrobin (18 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sugar Beet (roots and tops)	Cercospora leaf spot <i>Cercospora beticola</i> Powdery mildew <i>Erysiphe betae</i>	9 to 12	2	48 (0.78 lb ai/acre)	7

Application Directions. Begin **Headline SC** applications prior to disease development and continue on a 14-day interval.

Use the higher rate when disease pressure is high.

Headline SC applications will aid in the control of Rhizoctonia stem canker and crown rot.

In sugar beet, **Headline SC** can be combined with low rates of crop oil concentrate (COC), methylated seed oil (MSO), and nonionic surfactant (NIS) adjuvants. **DO NOT** use silicone-containing adjuvants. Some combinations and rates may result in temporary crop injury.

Headline SC Tank Mixes. **Headline SC** can be tank mixed with herbicides such as **Poast® herbicide, Select® herbicide, Assure® II herbicide** or **Prism® herbicide** for postemergence control of grasses in sugar beet. **DO NOT** use silicone-based adjuvants in such combinations. **Headline SC** tank mix combinations can include COC or MSO; however, crop injury may result. The level of injury tends to increase with increasing rates of COC or MSO.

See **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.78 lb ai pyraclostrobin (48 fl ozs **Headline SC**) per acre per year.

DO NOT make more than one (1) **Headline SC** application before the 4-leaf stage of plant growth. After the 4-leaf stage of plant growth, **DO NOT** make more than (1) **Headline SC** application before alternating to a **non-Group 11** fungicide with a different mode of action.

*The maximum product rate per year includes the combination of in-furrow and foliar uses.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne Rhizoctonia in Sugar Beet									
Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)								
	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.1	3.5								
0.2	7.0	5.2	4.7	3.5	3.3	3.2	3.0		
0.3	10.5	7.8	7.1	5.2	5.0	4.8	4.5	4.3	4.0
0.4	see footnote ¹	10.4	9.5	6.9	6.7	6.4	6.0	5.7	5.4
0.5	see footnote ¹	see footnote ¹	11.8	8.7	8.4	8.0	7.5	7.1	6.7
0.6	see footnote ¹	see footnote ¹	see footnote ¹	10.4	10.0	9.6	9.0	8.5	8.1
0.7	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	11.7	11.2	10.5	10.0	9.4
0.8	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	see footnote ¹	12.0	11.4	10.8

Application Directions. Use 0.1 to 0.8 fl oz of **Headline SC** per 1000 row feet. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered. Use a minimum application volume of 2.5 gallons of water per acre.

When Rhizoctonia seedling disease pressure conditions are expected to be severe or if the field has a history of seedling diseases, use **Headline SC** at a product rate per acre equivalent to 9 to 12 fl ozs and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre of **Headline SC**.

¹ For 32- to 34-inch rows, use a maximum of 0.7 fl oz per 1000 row feet.
 For 30-inch rows, use a maximum of 0.6 fl oz per 1000 row feet.
 For 22-inch rows, use a maximum of 0.5 fl oz per 1000 row feet.
 For 20-inch rows, use a maximum of 0.4 fl oz per 1000 row feet.
 For 15-inch rows, use a maximum of 0.3 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sugarcane*	Brown Rust <i>Puccinia melanocephala</i> Orange Rust <i>Puccinia kuehnii</i>	9 to 12	2	48 (0.78)	14

Application Directions. For optimal disease control, begin applications of **Headline SC** prior to disease development and continue on a 14 to 28 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 48 ozs of **Headline SC** per acre per year. **DO NOT** make more than two (2) sequential applications of **Headline SC** before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

* Not for use in California.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year* (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Tuberous and Corm Vegetables Subgroup Arracacha Arrowroot Chinese artichoke Jerusalem artichoke Cassava (bitter and sweet) Chayote (root) Chufa Dasheen Edible canna Ginger Leren Sweet potato Tanier True yam Turmeric Yam bean	Downy mildew <i>Plasmopara</i> spp. Leaf spot <i>Cercospora</i> spp., <i>Alternaria</i> spp. Powdery mildew <i>Erysiphe</i> spp., <i>Leveillula taurica</i> Rust <i>Uromyces</i> spp., <i>Puccinia</i> spp.	6 to 12	1	72 (1.18 lbs ai/acre)	3
Potato	Black dot <i>Colletotrichum coccodes</i>	6 to 9			
	Early blight <i>Alternaria solani</i>				
	Late blight <i>Phytophthora infestans</i> Powdery mildew <i>Erysiphe</i> spp., <i>Leveillula taurica</i> Suppression Only: White mold <i>Sclerotinia sclerotiorum</i>	6 to 12			

Application Directions. Begin **Headline SC** applications at 7- to 14-day intervals prior to disease development. The low rate and longer interval can be used early season prior to the observance of symptoms and when disease pressure is low. **For control of late blight**, follow **Headline SC** application with a labeled fungicide with a different mode of action 5 to 7 days later.

Use the higher rates and shorter intervals once disease has been confirmed in your area or if weather conditions are favorable for disease development.

No livestock feeding restrictions.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 1.18 lbs ai pyraclostrobin (72 fl ozs **Headline SC**) per acre per year.

DO NOT make more than one (1) **Headline SC** application before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

*The maximum product rate per year includes the combination of in-furrow and foliar uses. **(For above-listed crops, in-furrow use is permitted in potato only.)**

In-furrow Use to Aid in the Control of Soilborne Rhizoctonia in Potatoes

Use 0.4 to 0.8 fl oz of **Headline® SC fungicide** per 1000 row feet. For applications on 32-inch or 34-inch rows, the maximum application rate is 0.73 fl oz/1000 row feet. Apply at planting as an in-furrow spray by directing spray pattern to uniformly cover seed pieces and surrounding soil. The spray pattern must be a 4- to 8-inch band applied to the seed piece prior to being covered with soil.

When Rhizoctonia disease pressure conditions are expected to be severe or if the field has a history of Rhizoctonia infestations, use **Headline SC** at 0.6 to 0.8 fl oz per 1000 row feet and/or tank mix with a fungicide having a different mode of action.

DO NOT apply more than 12 fl ozs per acre of **Headline SC**.

Use a minimum volume of application of 5 gallons of water per acre.

Headline SC Rate per 1000 row feet (fl oz product)	Headline SC Rate (fl ozs/A)				
	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.4	6.7	6.4	6.0	5.7	5.4
0.6	10.0	9.6	9.0	8.6	8.1
0.8	see footnote ¹	see footnote ¹	12.0	11.4	10.8

¹ For 32-inch or 34-inch rows, use a maximum of 0.73 fl oz per 1000 row feet.

Table 2. Headline® SC fungicide Crop-specific Requirements (continued)

Crop	Target Disease	Product Rate per Application (fl ozs/A)	Maximum Number of Sequential Foliar Applications	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Wheat and Triticale	Black point Kernel smudge <i>Alternaria</i> spp., <i>Helminthosporium</i> spp. Leaf rust <i>Puccinia triticina</i> Powdery mildew <i>Erysiphe graminis</i> f. sp., <i>tritici</i> Septoria leaf and glume blotch <i>Septoria</i> spp., <i>Stagonospora</i> spp. Spot blotch <i>Cochliobolus sativus</i> Stem rust <i>Puccinia graminis</i> f. sp., <i>tritici</i> Stripe rust <i>Puccinia striiformis</i> f. sp., <i>tritici</i> Tan spot Yellow leaf spot <i>Pyrenophora</i> spp.	6 to 9*	2	18 (0.29 lb ai/acre)	Apply no later than the beginning of flowering (Feekes 10.5, Zadok's 59)

Application Directions. Begin **Headline SC** applications prior to disease development. To maximize yields in cereals, protect the flag leaf. Apply **Headline SC** immediately after flag-leaf emergence for optimum results.

Headline SC does not control Fusarium head blight (head scab) or prevent the reductions in grain quality that can result from this disease. When head blight is a concern, manage this disease with fungicides that are labeled for and effective in managing this disease, and with cultural practices like crop rotation and plowing to reduce crop residues that serve as an inoculum source.

DO NOT harvest wheat hay or feed green-chopped wheat within 14 days after last application.

Resistance Management. To limit development of resistance, **DO NOT** apply more than 0.29 lb ai pyraclostrobin (18 fl ozs **Headline SC**) per acre per year.

DO NOT make more than two (2) sequential **Headline SC** applications before alternating to a labeled **non-Group 11** fungicide with a different mode of action.

* For early season control of tan spot, Septoria leaf and glume blotch, and spot blotch when conditions favor disease development, apply 3 to 6 fl ozs per acre of **Headline SC** either in combination with a herbicide application or when conditions favor disease development. When the 3 to 6 fl ozs early season application rate is used, a second application of **Headline SC** may be required to protect the emerged flag leaf. Environmental conditions for disease or current disease pressure at the time of flag-leaf emergence should be used to determine the **Headline SC** rate for the second application. For high disease pressure, use the higher rate of **Headline SC**. Early season control is not registered for use in California.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Agricultural Solutions US LLC ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

***Cabrio, Headline, Poast, and Pristine** are registered trademarks of BASF.*

***Assure and Prism** are registered trademarks of E.I. duPont de Nemours and Company.*

***Select** is a registered trademark of Arysta Life Science North America Corporation.*

© 2026 BASF Agricultural Solutions US LLC
All rights reserved.

007969-00289.20201214e.**NVA 2020-04-340-0211**
Supersedes: NVA 2021-04-340-0062

BASF Agricultural Solutions US LLC
2 TW Alexander Drive
Research Triangle Park, NC 27713


We create chemistry