

7969-309

2/14/2014

1/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

FEB 14 2014

Khalid H. Akkari, Ph.D.
BASF Corporation
26 Davis Drive
P.O. Box 13528
Research Triangle Park, NC 27709-3528

Subject: Label Amendment – New Food Crop Uses
EPA Registration No.: 7969-309
Sercadis™ Xemium® Brand Fungicide
D#: 466683
Your submission dated June 15, 2012

Dear Dr. Akkari:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable.

At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate this supplemental labeling into the main product labeling. Two (2) copies of the final printed label must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

If you have any questions, please contact Olga Odiott at (703) 308-9369.

Sincerely,

A handwritten signature in black ink, appearing to read "Michael Walsh".

Michael Walsh
Acting Product Manager 13
Insecticide Branch
Registration Division (7505P)

2/19

Group 7 Fungicide

Supplemental Label

Sercadis™

XEMIUM® BRAND FUNGICIDE

For disease control in the following crops: berries and small fruits, Brassica leafy vegetables, bulb vegetables, cucurbit vegetables, grapes, leafy vegetables, rice, root vegetables, sorghum and millet, strawberries, sugarcane, and tree nuts

This supplemental label expires on September 30, 2016 and must not be used or distributed after this date.


Powered by Xemium® fungicide

Active Ingredient:
fluxapyroxad*: 1H-Pyrazole-4-carboxamide, 3-(difluoromethyl)-
1-methyl-N-(3',4',5'-trifluoro[1,1'-biphenyl]-2-yl)-

Other Ingredients:.....

Total:

* Equivalent to 2.47 pounds of active ingredient per gallon



26.55%
73.45%
100.00%

EPA Reg. No. 7969-309

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater.

This product is classified as having high potential for reaching aquatic sediment via runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this active ingredient or its degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Sercadis™ Xemium® brand fungicide** container label, EPA Reg. No. 7969-309, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Sercadis** before applying.
- Use of **Sercadis** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Sercadis**.

Restrictions and Limitations

Crop Rotation Restriction - Barley, berries and small fruits, Brassica leafy vegetables, bulb vegetables, corn (all types), cotton, cucurbit vegetables, dried shelled peas and beans, edible-podded legume vegetables, fruiting vegetables, grapes, leafy vegetables, millet, oat, oilseed crops (including flax seed, rapeseed and sunflower), peanut, pome fruits, rice, root vegetables, rye, sorghum, soybean, stone fruits, strawberries, succulent shelled peas and beans, sugar beet, sugarcane, tree nuts, tuberous and corm vegetables (including potato), and wheat and triticale may be

planted immediately following the last application. For all other crops, **DO NOT** plant sooner than 365 days after the last application.

Application Instructions

Apply **Sercadis™ Xemium®** brand fungicide according to the rate, timing, and resistance management use instructions in the **Crop-specific Directions** in this label. Observe the additional instructions on application methods, additive use and mixing order on the **Sercadis** main label.

Table 1. Sercadis™ Xemium® brand fungicide Restrictions and Limitations Overview*

Crop**	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Number of Sequential Applications	Maximum Product Rate per Season (fl ozs)
Berries and small fruits subgroups	0, 14	9.1	3	2	27.3
Brassica leafy vegetables group	3	4.6	3	2	13.8
Bulb vegetables group	7	9.1	3	2	27.3
Cucurbit vegetables group	0	4.6	3	2	13.8
Grapes – Botrytis disease	14	9.1	3	2	27.3
Grapes – all other diseases	14	4.55	6	2	27.3
Leafy vegetables group (except Brassica)	1	9.1	3	2	27.3
Rice	28	6.8	2	2	13.6
Root vegetables (except sugar beet) subgroup	7	4.6	3	2	13.8
Sorghum and millet	21	4.6	2	2	9.2
Strawberries	0	9.1	3	2	27.3
Sugarcane	14	5.7	2	2	11.4
Tree nuts group	14	5.7	3	2	17.1

*See Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions for additional directions.

**For a complete list of crops within a crop group, see Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions.

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Berries and small fruits subgroups* Bushberry subgroup Blueberry* (highbush and lowbush) Currant Elderberry Gooseberry Huckleberry Caneberry subgroup Blackberry (all varieties) Loganberry Raspberry (black and red) Wild raspberry Low growing berry subgroup Bearberry Bilberry Cloudberry Cranberry Lingonberry Muntries Partridgeberry	Alternaria leaf spot and fruit rot (<i>Alternaria</i> spp.) Botrytis gray mold (<i>Botrytis cinerea</i>) Leaf spot and blotch (<i>Mycosphaerella</i> spp., <i>Septoria</i> spp.) Monilinia blight and mummy berry (<i>Monilinia</i> spp.) Phomopsis leaf spot, twig blight, and fruit rot (<i>Phomopsis</i> spp.) Powdery mildew (<i>Sphaerotheca</i> spp., <i>Microshaera</i> spp., <i>Oidium</i> spp.) Spur blight (<i>Didymella</i> spp., <i>Phoma</i> spp.) Suppression Only: Rust (<i>Pucciniastrum</i> spp., <i>Arthuriomyces</i> spp., <i>Phragmidium</i> spp., <i>Kuehneola</i> spp.)	3.4 to 9.1	3	27.3	0 (bushberry, caneberry, and low growing berry)
Small fruit vine climbing subgroup, (except fuzzy kiwifruit) Amur river grape Gooseberry Kiwifruit, hardy Maypop Schisandra berry					14 (small fruit vine climbing)

Application Directions. Begin applications of **Sercadis** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

DO NOT apply **Sercadis** to any crops in the Berries and Small Fruits subgroups, including blueberries (highbush and lowbush), as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Sercadis** with water only for applications to crops listed in the Berries and Small Fruits subgroups.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 27.3 fl ozs of **Sercadis** per acre per season. **DO NOT** apply more than three (3) applications of **Sercadis** per season.

DO NOT make more than two (2) sequential applications of **Sercadis** before alternating to a labeled non-Group 7 fungicide.

*Not registered for use in California.

5/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)	
Brassica leafy vegetables group* Head and stem Broccoli Broccoli, Chinese Brussels sprouts Cabbage Cabbage, Chinese Cabbage, Chinese mustard Cauliflower Cavalo broccolo Kohlrabi Leafy greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens	Alternaria leaf spot (<i>Alternaria</i> spp.) Black leg (<i>Phoma lingam</i>) Cercospora leaf spot (<i>Cercospora brassicicola</i>) Powdery mildew (<i>Erysiphe polygoni</i>) Rhizoctonia blight (<i>Rhizoctonia solani</i>) Ring spot (<i>Mycosphaerella brassicicola</i>) White leaf spot (<i>Pseudocercospora capsellae</i>)	3.4 to 4.6	3	13.8	3	
	Suppression Only:					
	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)					
	Southern blight (<i>Sclerotium rolfsii</i>)					

Application Directions. Begin applications of **Sercadis** prior to onset of disease development and continue on a 7 to 14 day interval.

Use the shorter interval and/or the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 13.8 fl ozs of **Sercadis** per acre per season. **DO NOT** apply more than three (3) applications of **Sercadis** per season.

DO NOT make more than two (2) sequential applications of **Sercadis** before alternating to a labeled **non-Group 7** fungicide.

*Not registered for use in California.

4/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Bulb vegetables group* Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves	Powdery mildew (<i>Leveillula taurica</i>) Purple blotch and leaf blight (<i>Alternaria porri</i>) Rust (<i>Puccinia porri</i>) Stemphylium leaf blight and stalk rot (<i>Stemphylium vesicarium</i>)	3.4 to 9.1	3	27.3	7
	Botrytis leaf blight (<i>Botrytis</i> spp.) Botrytis neck rot (<i>Botrytis</i> spp.)	4.6 to 9.1			

Application Directions. Begin applications of **Sercadis** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 27.3 fl ozs of **Sercadis** per acre per season. **DO NOT** apply more than three (3) applications of **Sercadis** per season.

DO NOT make more than two (2) sequential applications of **Sercadis** before alternating to a labeled **non-Group 7** fungicide.

*Not registered for use in California.

7/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Cucurbit vegetables group*	Alternaria leaf blight (<i>Alternaria cucumerina</i>)	3.4 to 4.6	3	13.8	0
Chayote Chinese waxgourd Citron melon	Powdery mildew (<i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp., <i>Erysiphe</i> spp.)		.		
Cucumber Gherkin Pumpkin Watermelon	Cercospora leaf spot (<i>Cercospora citrulina</i>)	4.6	.		
Edible gourd Hyotan Cucuzza Chinese okra	Gummy stem blight (<i>Didymella bryoniae</i>) Microdochium blight (<i>Plectosporium tabacinum</i>)		.		
Momordica spp. Balsam apple Balsam pear Bitter melon Chinese cucumber	Target leaf spot (<i>Corynespora cassicola</i>)		.		
Muskmelon Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persian melon Pineapple melon Santaclaus melon Snake melon			.		
Summer squash Crookneck squash Scallop squash Straightneck squash Vegetable marrow Zucchini			.		
Winter squash Butternut squash Calabaza Hubbard squash Acorn squash Spaghetti squash			.		

(continued)

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Cucurbit vegetables group, continued

Application Directions. For optimal disease control, begin applications of **Sercadis** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

DO NOT apply **Sercadis** to any crops in the Cucurbit Vegetable Group as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Sercadis** with water only for applications to crops listed in the Cucurbit Vegetable Group.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 13.8 fl ozs of **Sercadis** per acre per season. **DO NOT** make more than two (2) sequential applications of **Sercadis** before alternating to a labeled **non-Group 7** fungicide. In areas where gummy stem blight resistance to **Group 7** fungicides have been confirmed, tank mix with chlorothalonil at full label rates and adhering to all label precautions.

*Not registered for use in California.

9/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Grapes	Powdery mildew (<i>Erysiphe necator</i>)	3.4 to 4.55	6	27.3	14
	Black rot (<i>Guignardia bidwellii</i>)	4.55			
	Botrytis gray mold (<i>Botrytis cinerea</i>) Aids in control: Summer bunch rot (<i>Aspergillus</i> spp., <i>Alternaria</i> spp., <i>Botrytis</i> spp., <i>Cladosporium</i> spp., <i>Penicillium</i> spp., <i>Rhizopus</i> spp.)	4.5 to 9.1	3		

Application Directions. For powdery mildew control, begin applications of **Sercadis** as of bud break or prior to onset of disease. When using 3.4 fl ozs per acre, continue on a 10 to 14 day interval. Longer spray intervals for powdery mildew control may be possible with higher rates of **Sercadis**. The effectiveness of longer spray intervals will depend on the current powdery mildew infection level in the field, the amount of disease pressure after application and factors such as crop growth stage and rate of growth.

For black rot control, begin applications of **Sercadis** before pre-bloom and continue on a 10 to 14 day interval.

Begin applications targeted against **Botrytis gray mold** prior to disease development when conditions favor disease development during early bloom, bunch preclosure and veraison. Use the higher rates from bunch preclosure to veraison. Use shorter intervals when disease pressure is high.

For all other diseases listed, begin applications of **Sercadis** prior to onset of disease development and continue on a 10 to 14 day interval. Use the higher rate and shorter interval when disease pressure is high.

DO NOT apply **Sercadis** to Grapes as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Sercadis** with water only for application to Grapes.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 27.3 fl ozs of **Sercadis** per acre per season. **DO NOT** make more than two (2) sequential applications of **Sercadis** before alternating to a labeled **non-Group 7** fungicide.

10/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Leafy vegetables group* (except Brassica)	Alternaria leaf spot (<i>Alternaria</i> spp.)	3.4 to 9.1	3	27.3	1
	Ascochyta leaf spot (<i>Ascochyta</i> spp.)				
Amaranth	Cercospora leaf spot (<i>Cercospora</i> spp.)	4.6 to 9.1			
Arugula	Phoma (<i>Phoma</i> spp.)				
Cardoon	Powdery mildew (<i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.)				
Celery	Rust (<i>Puccinia</i> spp., <i>Uromyces</i> spp.)				
Celery, Chinese	Septoria leaf spot (<i>Septoria</i> spp.)				
Celtuce	Lettuce drop caused by <i>Sclerotinia minor</i>				
Chervil	Botrytis rot (<i>Botrytis</i> spp.)				
Chrysanthemum (edible-leaved and garland)	Suppression Only:				
Corn salad	Lettuce drop caused by <i>Sclerotinia sclerotiorum</i>				
Cress (garden and Upland)					
Dandelion					
Dock					
Endive					
Fennel, Florence					
Lettuce (head and leaf)					
Orach					
Parsley					
Purslane (garden and winter)					
Radicchio (red chicory)					
Rhubarb					
Spinach					
Spinach (New Zealand and vine)					
Swiss chard					

(continued)

11/
19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Leafy vegetables group, continued

Application Directions. Begin applications of **Sercadis** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Tank Mix Restrictions

Spinach (all varieties). **DO NOT** apply **Sercadis** to spinach as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Sercadis** with water only for applications to Spinach (all varieties).

Leafy Vegetables (except spinach). It is impossible for BASF to test all varieties of leafy vegetables for sensitivity to **Sercadis** under all environments and all potential product mixture combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Sercadis** use, particularly in tank mixes and/or adjuvant combinations on leafy vegetables. To reduce the risk of leafy vegetable injury, BASF recommends testing **Sercadis** or **Sercadis** tank mixtures on a small portion of the crop before broadscale use.

To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Sercadis** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 27.3 fl ozs of **Sercadis** per acre per season. **DO NOT** apply more than three (3) applications of **Sercadis** per season.

DO NOT make more than two (2) sequential applications of **Sercadis** before alternating to a labeled **non-Group 7** fungicide.

*Not registered for use in California.

12/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Rice* Wild Rice*	Aggregate sheath spot (<i>Rhizoctonia oryzae-sativae</i>) Sheath blight (<i>Rhizoctonia solani</i>)	3.4 to 6.8	2	13.6	28

Application Directions. Begin applications of **Sercadis** at the first sign of disease. Repeat applications on 7 to 14 day intervals as needed if conditions for disease infection continue. Use the shorter interval and/or the higher rate when disease pressure is high.

Sercadis can be applied by ground sprayer or aerial equipment. For ground application, apply a minimum of 15 gpa (gallons per acre) spray solution. For aerial application, apply 7 to 10 gpa spray solution. **DO NOT** apply **Sercadis** through any type of irrigation system. Apply **Sercadis** with a non-ionic spray (NIS) adjuvant at 0.25% v/v. Under hot weather conditions apply **Sercadis** with 1% COC. BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 13.6 fl ozs of **Sercadis** per acre per season. **DO NOT** make more than two (2) sequential applications of **Sercadis** before alternating to a labeled non-Group 7 fungicide.

*Not Registered for use in California.

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl.ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Root vegetables (except sugar beet) subgroup* Beet, garden Burdock, edible Carrot Celeriac Chervil, turnip-rooted Chicory Ginseng Horseradish Parsley, turnip-rooted Parsnip Radish Radish, Oriental Rutabaga Salsify Salsify, black Salsify, Spanish Skirret Turnip	Alternaria leaf spot/blight (<i>Alternaria</i> spp.) Powdery mildew (<i>Erysiphe</i> spp., <i>Leveillula</i> spp.)	3.4 to 4.6	3	13.8	7
	Cercospora leaf spot/blight (<i>Cercospora</i> spp.) Suppression Only: Sclerotinia white mold/cottony rot (<i>Sclerotinia sclerotiorum</i>) Southern blight (<i>Sclerotium rolfsii</i>)	4.6			
<p>Application Directions. For optimal disease control, begin applications of Sercadis prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.</p>					
<p>Resistance Management. To limit the potential for development of resistance, DO NOT apply more than 13.8 fl ozs of Sercadis per acre per season. DO NOT make more than two (2) sequential applications of Sercadis before alternating to a labeled non-Group 7 fungicide.</p>					
<p>*Not registered for use in California.</p>					

14/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sorghum* Milo* Millet* (pearl and proso)	Gray leaf spot and Cercospora leaf spot (<i>Cercospora</i> spp.) Northern leaf blight (<i>Exserohilum turcicum</i>) Rust (<i>Puccinia</i> spp.) Southern leaf blight and Bipolaris leaf spot (<i>Bipolaris</i> spp.)	3.4 to 4.6	2	9.2	21

Application Directions. For optimal disease control, begin applications of **Sercadis** prior to disease development. Use the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 9.2 fl ozs of **Sercadis** per acre per season. **DO NOT** apply more than two (2) sequential applications of **Sercadis** before alternating to a non-Group 7 fungicide.

*Not registered for use in California.

15/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Strawberries*	Leaf spot (<i>Mycosphaerella fragariae</i> , <i>Ramularia tulasnei</i>)	3.4 to 5.7	3	27.3	0
	Powdery mildew (<i>Sphaerotheca macularis</i>)				
	Botrytis gray mold (<i>Botrytis cinerea</i>)	6.9 to 9.1			

Application Directions. Begin applications of **Sercadis** no later than 10% bloom, or prior to disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

The restricted entry interval (REI) for treated strawberries is **12 hours**. Refer to the **Agricultural Use Requirements** on the **Sercadis** main label for PPE required for early entry to treated areas as permitted under the Worker Protection Standard.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 27.3 fl ozs of **Sercadis** per acre per season. **DO NOT** make more than two (2) sequential applications of **Sercadis** before alternating to a labeled **non-Group 7** fungicide.

*Not registered for use in California.

16/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Sugarcane*	Brown rust (<i>Puccinia melanocephala</i>) Orange rust (<i>Puccinia kuehnii</i>)	3.4 to 5.7	2	11.4	14

Application Directions. Begin applications of **Sercadis** at the first sign of disease. Repeat applications on 14 to 28 day intervals as needed if conditions for rust infection continue. Use the shorter interval and/or the higher rate when disease pressure is high.

Sercadis can be applied by ground or air. When applying by air, **DO NOT** use less than 5 gallons of spray solution per acre.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 11.4 fl ozs of **Sercadis** per acre per season. **DO NOT** apply more than two (2) sequential applications of **Sercadis** before alternating to a labeled **non-Group 7** fungicide.

*Not registered for use in California.

17/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Season	Maximum Product Rate per Season (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Tree Nuts Group* Almond Beech nut Brazil nut Butternut Cashew Chestnut Chiquapin Filbert Hickory nut Macadamia nut Pecan Pistachio Walnut (black and English)	Alternaria late blight (<i>Alternaria</i> spp.) Botrytis blossom and shoot blight (<i>Botrytis cinerea</i>) Brown rot/Blossom blight (<i>Monilinia</i> spp.) Eastern filbert blight (<i>Anisogramma anomala</i>) Green fruit rot/ Jacket rot (<i>Botrytis cinerea</i> , <i>Sclerotinia sclerotiorum</i> , <i>Monilinia laxa</i>) Leaf rust (<i>Tranzschelia discolor</i>) Panicle and shoot blight (<i>Botryosphaeria dothidea</i>) Scab (<i>Cladosporium carpophilum</i> , <i>C. caryigenum</i>) Shothole (<i>Wilsonomyces carpophilus</i>) <u>Suppression Only:</u> Hull rot (<i>Rhizopus stolonifer</i> and <i>Monilinia</i> spp.)	3.4 to 5.7	3	17.1	14

(continued)

18/19

Table 2. Sercadis™ Xemium® brand fungicide Crop-specific Directions (continued)

Tree nuts group, continued

Application Directions. For almond, begin applications of Sercadis prior to onset of disease development and continue on a 7 to 14 day interval up to 14 days before harvest.

For filbert, begin applications at budswell to budbreak, or prior to infection and onset of disease development. Continue on a 7 to 14 day interval to cover and protect new growth.

For pecan, begin applications of Sercadis prior to onset of disease development and continue on a 7 to 21 day interval for the control of scab. Use the shorter interval and/or the higher rate when disease pressure is high.

For pistachio, apply Sercadis prior to onset of disease development and continue on a 10 to 30 day interval. Use the higher rate and shorter intervals when disease pressure is high.

For all other crops listed, apply Sercadis prior to disease development and continue on a 7 to 28 day interval. In all cases use the shorter interval when shoot growth is rapid.

No restriction on livestock feeding of almond hulls.

For aerial application to tree nuts, **DO NOT** use less than 10 gallons of spray solution per acre.

Resistance Management. To limit the potential for development of resistance, **DO NOT** apply more than 17.1 fl ozs of Sercadis per acre per season. **DO NOT** make more than two (2) sequential applications of Sercadis before alternating to a labeled non-Group 7 fungicide.

* Not registered for use in California.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

19/19

Sercadis™ Xemium® brand fungicide is specially formulated and sold by BASF for the control of pests according to the directions on this label. The purchase price of **Sercadis** includes a royalty whereby the purchaser acquires a prepaid license under U.S. Patent No. 5,438,070 and (pending No. 2008/0153707 A1) under which purchaser agrees to employ the purchased quantity of **Sercadis** only for the above-specified uses under BASF's United States patent rights and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of **Sercadis** other than those specified on this label are not licensed through the purchase of this product and the use of this product for other purposes may violate this license and patent rights of BASF.

Xemium is registered trademark of BASF.

Sercadis is a trademark of BASF.

©2014 BASF Corporation
All rights reserved.

007969-00309.20120613c.NVA 2012-04-370-0099

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



The Chemical Company