



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

**OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION**

January 18, 2024

Elizabeth Kneller
Product Registration Manager
BASF Corporation, Agricultural Products
P.O. Box 13528, 26 Davis Drive
Research Triangle Park, NC 27709

Subject: PRIA Label Amendment – Addition of avocado use
Product Name: Merivon Xemium Brand Fungicide
EPA Registration Number: 7969-310
Application Date: October 25, 2021; October 17, 2022
Case Number: 00482406; 00476883

Dear Ms. Kneller:

The application referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable under FIFRA section 3(c)(7)(B), subject to the following conditions:

1. You must submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.
2. Be aware that proposed data requirements have been identified in a Work Plan. For more information on these proposed data requirements, you may contact the Chemical Review Manager in the Pesticide Re-Evaluation Division:
<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements

Page 2 of 2
EPA Reg. No. 7969-310
Case No. 00482406; 00476883

EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e).

If you have any questions, please contact Lindsay DeMers at demers.lindsay@epa.gov.

Sincerely,




Nathan Mellor, Acting Branch Chief
Fungicide and Herbicide Branch
Registration Division (7505T)
Office of Pesticide Programs

Enclosure: stamped "accepted" label

[NOTE to PM: This text to appear at the top of page 1 of the container booklet.] The label accessed from the QR code is legally valid only for use in registered states and only with a container of **Merivon® Xemium® brand fungicide**, EPA Reg. No. 7969-310. For pesticide applications following the label accessed from the QR code, you must possess a copy of the QR code label at the time of application. If the QR code label conflicts with the labeling found on the pesticide product, use only one set of the labeling instructions in its entirety. **DO NOT** mix and match labeling directions. The QR code label is the most current version of the **Merivon**, EPA Reg. No. 7969-310 label approved by US EPA and state lead agencies. Currently the user is responsible for complying with the requirements of product labeling obtained when the user took possession of the product's container. The user must comply with all instructions and requirements in the labeling and must keep the labeling as long as the user has the container. Similarly, the user who relies upon the QR code label may meet this requirement by maintaining a printed or electronic copy of the labeling to meet federal and state record keeping requirements. In the situations described above where the QR code label conflicts with labeling on the pesticide container, the user must be able to identify which labeling was used and followed at the time of the pesticide's application.



Pyraclostrobin	Group	11	Fungicide
Fluxapyroxad	Group	7	Fungicide



[NOTE to PM: (1) This is a placeholder. The final printed label/container label will bear a unique QR code that when scanned takes the user to the PDF version of the most current label approved by US EPA and state lead agencies where the product is sold. The container label and QR code PDF label both bear EPA required web distributed labeling text per PR-Notice 2014-1. (2) The user must choose a single, valid label version to follow in its entirety. (3) The QR code locations include (based on package type) carton, front panel of booklet, container decal/stay-behind.]



Merivon®

Xemium® Brand Fungicide

For disease control and plant health in the following crops: avocado, berries and small fruits, Brassica leafy vegetables, bulb vegetables, cucurbit vegetables, fruiting vegetables (including tomato), grapes, leafy vegetables, oilseed crops (including flax seed, rapeseed and sunflower), pome fruits, pomegranate, root vegetables, soybean, stone fruits, strawberries, tree nuts, and tuberous and corm vegetables (including potato)

Powered by **Xemium®** and **F500®** fungicides

Active Ingredients:

fluxapyroxad*: 1 <i>H</i> -Pyrazole-4-carboxamide, 3-(difluoromethyl)-1-methyl-N-(3',4',5'-trifluoro[1,1'-biphenyl]-2-yl)-	21.26%
pyraclostrobin**: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1 <i>H</i> -pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester)	21.26%
Other Ingredients:	57.48%
Total:	100.00%

* Equivalent to 2.09 pounds of fluxapyroxad per gallon.
 ** Equivalent to 2.09 pounds of pyraclostrobin per gallon.

EPA Reg. No. 7969-310 **EPA Est. No.**

KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See full label for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Shake Well Before Using

Net Contents:

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).	

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING. May be fatal if swallowed. Harmful if inhaled. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils)
- Shoes plus socks

User Safety Requirements

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

This product may impact surface water quality because of runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this active ingredient or its degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding application when rainfall is forecast to occur within 48 hours. Sound erosion

control practices will reduce this product's potential to reach aquatic sediment via runoff.

Groundwater Advisory

Pyraclostrobin is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Directions For Use

Read the entire **Directions For Use** and **Conditions of Sale and Warranty** before using this product.

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

You may obtain labeling from the QR code on the front page of this label. If using the additional labeling to apply the product, you must possess a copy of this additional labeling at the time of application. It is a violation of Federal law to use this product in a manner inconsistent with its attached label or any additional labeling, including any web-distributed labeling. In instances where the additional web-distributed labeling conflicts with the container label, the user may choose a single, valid version of the labeling to follow. However, for areas of overlap or conflict, the user must use only one set of labeling instructions, either the attached container label or the web-distributed labeling. **DO NOT** mix and match labeling directions.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

Exception is a 5 day REI for:

- cane tying and leaf pulling tasks in grapes. Notify workers of this prohibition.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves (barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils)
- Shoes plus socks

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

(continued)

STORAGE AND DISPOSAL *(continued)*

Container Handling *(continued)*

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity > 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

(continued)

STORAGE AND DISPOSAL *(continued)*

Container Handling *(continued)*

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to be taken in case material is released or spilled:

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

Product Information

This package contains **Merivon® Xemium® brand fungicide**, a suspension concentrate (SC) containing fluxapyroxad and pyraclostrobin. The active ingredients in **Merivon** belong to two classes of fungicides, the strobilurins or Quinone Outside Inhibitors (QoI) and succinate-dehydrogenase (SDH) inhibitor classes. To maximize disease control, apply **Merivon** in a regularly scheduled protective spray program and use in a rotation program with other fungicides.

Preventive applications optimize disease control, resulting in improved plant health. The increase in plant health comes from the combined effect of disease control (including fungal diseases listed in Crop-specific directions), improved growth efficiency and improved stress tolerance. Overall increased plant health may result in an improvement in crop growth and crop quality as well as increased crop yields.

Because of its high specific activity, **Merivon® Xemium® brand fungicide** has residual activity against target fungi.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Merivon is not for use in greenhouse or transplant production.

Modes of Action

Fluxapyroxad and pyraclostrobin, the active ingredients of **Merivon**, belong to the groups of respiration inhibitors classified as target-site-of-action **Group 7** and **Group 11** fungicides, respectively.

Resistance Management

For resistance management, please note that **Merivon** contains a premix of **Group 7** (fluxapyroxad) and **Group 11** (pyraclostrobin) fungicides. Any fungal population may contain individuals naturally resistant to **Merivon** and other **Group 7** or **Group 11** fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Merivon** or other **Group 7** or **Group 11** fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such a use is permitted. Use at least the minimum labeled rates of each fungicide in the tank mix.
- Adopt an integrated disease and pest management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treatment area for lack of efficacy that might indicate possible resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide

resistance-management and/or IPM recommendations for specific crops and pathogens.

- For further information or to report suspected resistance contact your local BASF representative, extension specialist or certified crop advisor.

Application Instructions

Apply specified rates of **Merivon** as instructed in **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions**. **Merivon** can be applied by ground and aerial application. For best results, thorough coverage of plant materials is required. **Merivon** can also be applied through sprinkler irrigation equipment. Application using drip irrigation equipment is permitted in select crops as instructed in **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions**. Check equipment frequently for calibration.

Under low-level disease conditions, the minimum application rates can be used while maximum application rates and shortened spray schedules are advised for severe or threatening disease conditions.

Ground Application

Apply **Merivon** in sufficient water to ensure thorough coverage of foliage, bloom, and fruit. Thorough coverage is required for optimum disease control. Complete coverage of the stem, all the way down to the soil, is required for suppression of soilborne diseases of the stem.

Instructions for Directed or Banded Crop Sprays

The application rates shown in **Table 1. Merivon® Xemium® brand fungicide Restrictions and Limitations Overview** and **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions** on this label reflect the amount of product to be applied uniformly over an acre of ground on a broadcast basis. In some crops, **Merivon** may be used as a directed or banded spray over the rows or plant beds with the alleys or row middles left unsprayed. For such uses, reduce the rate of **Merivon** in proportion to the area actually sprayed. Make this adjustment to avoid applying the product at use rates higher than permitted on this label.

The following formula may be used to determine the broadcast equivalent rate for doing directed or banded sprays:

$$\text{sprayed bed width} + \text{unsprayed row middle width} = \text{total row width}$$

$$\frac{\text{sprayed bed width in inches}}{\text{total row width in inches}} \times \frac{\text{broadcast rate}}{\text{treated acre}} = \frac{\text{band rate}}{\text{field acre}}$$

Example: A directed spray application will be made to 45-inch plant beds that are separated by 15 inches of unsprayed row middles.

45 inches sprayed bed width + 15 inches unsprayed row middles = 60 inches total row width

The calculation to determine the appropriate equivalent rate of product to use for this situation based on a label broadcast rate of 4 fluid ounces product/acre follows:

$$\frac{45 \text{ inches sprayed bed width}}{60 \text{ inches total row width}} \times \frac{4 \text{ fl ozs Merivon® Xemium® brand fungicide}}{\text{treated acre}} = \frac{3 \text{ fl ozs Merivon}}{\text{field acre}}$$

Mandatory Spray Drift Management

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to select nozzle and pressure that deliver a medium or coarser droplet size (ASABE S641).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 to 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 ft above the ground or crop canopy.
- Applicators are required to select nozzle and pressure that deliver uniform application and desired spray quality. A medium or coarser droplet size is preferred (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Boomless Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

Importance Of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

Boom Height – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Release Height - Aircraft

Higher release heights increase the potential for spray drift.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly

dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Aerial Application

For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (including, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fishponds).

For all crops listed in this label, aerial application can be made and thorough coverage is required to obtain optimum disease control. Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. **DO NOT** use less than 2 gallons of spray solution per acre. For aerial applications to tree crops, **DO NOT** use less than 10 gallons of spray solution per acre. For aerial applications to avocado, **DO NOT** use less than 20 gallons of spray solution per acre. For all other crops, thorough coverage is required for optimum disease control. The reduced spray volumes used in aerial applications may result in physical incompatibility, reduced disease control, or crop injury from **Merivon® Xemium® brand fungicide** applications, particularly when tank mixed with other products. Therefore, before making aerial applications test the spray on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

DO NOT apply when possible drift can occur to: unprotected persons; to food, forage, or other plantings that might be damaged; or crops that would then be rendered unfit for sale, use or consumption.

Where states have more stringent regulations, they must be observed.

Sensitive Areas

The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g. bodies of water or non-target crops) is minimal and when wind is blowing away from the sensitive areas.

Directions for Use Through Irrigation Systems

Sprayer Preparation

Clean chemical tank and injector system thoroughly. Flush system with clean water.

Application Instructions

Apply **Merivon** at rates and timings as required in this label.

Use Precautions for Sprinkler Irrigation and Drip Irrigation Applications

- This product can be applied through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems equipment. Application using drip irrigation is permitted in select crops as instructed in **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions. DO NOT** apply this product through any other type of irrigation system.
- Add **Merivon** to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop. **DO NOT** exceed 1/2 inch (13,577 gallons) of water per acre. In stationary or noncontinuous moving systems, inject the product-water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Thorough coverage of foliage is required for optimal control. Maintain agitation during the entire application period.
- Contact a state extension service specialist, equipment manufacturers or other experts for calibration questions.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of

the responsible person, must shut the system down and make necessary adjustments should the need arise.

- **DO NOT** connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There must be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Additives and Tank Mixing Information

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Merivon® Xemium® brand fungicide can be tank mixed with fungicides, insecticides, herbicides, liquid fertilizers,

biological control products, adjuvants, and additives as specified in **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions**. See avocado, berries and small fruits subgroups, cucurbit vegetables group, fruiting vegetables group, grapes, and leafy vegetables group in **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions** for exceptions.

Under some conditions, the use of additives or adjuvants may improve the performance of **Merivon**. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Merivon** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

When an adjuvant is to be used with this product, BASF advises the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

If tank mixes are used, follow restrictions for rates, label instructions, and precautions on all labels.

Compatibility Test for Tank Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of label rate per acre:

1. **Water** - For 100 gallons per acre spray volume, use 16 cups (1 gallon) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. **Water-dispersible products** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) - Cap the jar and invert 10 cycles.
3. **Water-soluble products** - Cap the jar and invert 10 cycles.
4. **Emulsifiable concentrates** (oil concentrate or methylated seed oil when applicable) - Cap the jar and invert 10 cycles.
5. **Water-soluble additives** - Cap the jar and invert 10 cycles.
6. Let the solution stand for 15 minutes.
7. **Evaluate** the solution for uniformity and stability. The spray solution must not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

Mixing Order

Maintain constant agitation throughout mixing and application. Make sure that each component is thoroughly mixed and suspended before adding tank mix partners. See **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions** for more details and applicable restrictions.

1. **Water** - Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
3. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. **Water-dispersible products** (including dry flowables, wettable powders, suspension concentrates including **Merivon® Xemium® brand fungicide**, or suspo-emulsions):
 - **Containers 5 gallons or less:** shake well before use.
 - **Containers larger than 5 gallons:** recirculate before use.
 - Consult BASF Representatives for additional information regarding agitation and recirculation.
5. **Water-soluble products**
6. **Emulsifiable concentrates** (including oil concentrates when applicable)
7. **Water-soluble additives** (including ammonium sulfate [AMS] or urea ammonium nitrate [UAN] when applicable)
8. **Remaining quantity of water**

Cleaning Spray Equipment

Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with potential to injure crops was used prior to **Merivon**.

Restrictions and Limitations

- **DO NOT** exceed the maximum product rate (fl ozs/A) per year, the maximum product rate per application, or the maximum number of applications of **Merivon** per year as stated in **Table 1. Merivon® Xemium® brand fungicide Restrictions and Limitations Overview** and **Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions**. Retreatment and preharvest interval (PHI) restrictions are also included in these tables.

- **DO NOT** use less than the minimum gallons per acre listed in the crop-specific directions (**Table 3**).
- **DO NOT** apply more than the maximum annual use rate of ai/acre for each specific crop from any combination of products containing pyraclostrobin or fluxapyroxad. See **Table 2. Merivon® Xemium® brand fungicide Rate Conversions** for corresponding pounds active ingredient per acre.
- **DO NOT** use **Merivon** in greenhouse or transplant production.
- **Merivon** is not for sale, distribution, or use in Nassau and Suffolk counties in New York State.
- **DO NOT** use **Merivon** on Concord, Noiret and NY73.0136.17 grape varieties due to possible foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related varieties. Not all varieties have been thoroughly tested.
- **Crop Rotation Restriction** - The following crops may be planted immediately following the last application: alfalfa, barley, berries and small fruits, Brassica leafy vegetables, bulb vegetables, corn (all types), cotton, cucurbit vegetables, dried shelled peas and beans, edible-podded legume vegetables, fruiting vegetables (including tomato), grapes, leafy vegetables, millet, mint (spearmint and peppermint), oat, oilseed crops (including flax seed, rapeseed and sunflower), peanut, pome fruits, pomegranate, root vegetables, rye, sorghum, soybean, stone fruits, strawberries, succulent shelled peas and beans, sugar beet, sugarcane, tree nuts, tuberous and corm vegetables (including potato), and wheat and triticale, and any other crop labeled for direct application of this product.

For rice, **DO NOT** plant sooner than 14 days after the last application.

For all other crops not listed above, **DO NOT** plant sooner than 365 days after the last application.

Table 1. Merivon® Xemium® brand fungicide Restrictions and Limitations Overview*

Crop**	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Retreatment Interval (days)	Maximum Number of Applications per Year	Maximum Number of Sequential Applications	Maximum Product Rate per Year (fl ozs/A)
Avocado	0	11	14	3	2	33
Berries and small fruits subgroups	0	11	7	3	2	33
Bushberry crop subgroup 13-07B						
Caneberry crop subgroup 13-07A						
Low growing berry (except cranberry) crop subgroup 13-07G						
Small fruit vine climbing (except fuzzy kiwifruit) crop subgroup 13-07F	14					
Brassica leafy vegetables crop subgroups 5A and 5B	3	5.5	7	3	2	16.5
Bulb vegetables crop group 3-07	7	11	7	3	2	33
Cucurbit vegetables crop group 9	0	5.5	7	3	1	16.5
Fruiting vegetables crop group 8-10	0	5.5	7	3	2	16.5
Grapes	14	5.5	28	6	2	33
Leafy vegetables (except Brassica) crop group 4	1	11	7	3	2	33
Oilseed crop group 20	21	5.5	7	2	2	11
Pome fruits crop group 11-10	0	5.5	7	4	2	22
Pomegranate	14	7	7	2	2	14
Root vegetables (except sugar beet) crop subgroup 1B	7	5.5	7	3	2	16.5
Soybean	21	5.5	7	2	2	11
Stone fruits crop group 12	0	6.7	7	3	2	20.1
Strawberries	0	11	7	3	2	33
Tree nuts crop group 14	14	6.5	7	3	2	19.5

(continued)

Table 1. Merivon® Xemium® brand fungicide Restrictions and Limitations Overview*

(continued)

Crop**	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Retreatment Interval (days)	Maximum Number of Applications per Year	Maximum Number of Sequential Applications	Maximum Product Rate per Year (fl ozs/A)
Tuberous and corm vegetables subgroup 1C	7	5.5	7	3	2	16.5

* See Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions for additional directions.
 ** For a complete list of crops within a crop group, see Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions.

Table 2. Merivon® Xemium® brand fungicide Rate Conversions*

Rate (fl ozs/A)	Fluxapyroxad (lb ai/A)	Pyraclostrobin (lb ai/A)
3	0.049	0.049
4	0.065	0.065
4.8	0.078	0.078
5	0.082	0.082
5.1	0.083	0.083
5.4	0.088	0.088
5.5	0.090	0.090
6	0.098	0.098
6.5	0.106	0.106
6.7	0.109	0.109
7	0.114	0.114
8	0.131	0.131
11	0.180	0.180
14	0.228	0.228
16.5	0.269	0.269
19.5	0.318	0.318
20.1	0.328	0.328
22	0.359	0.359
33	0.539	0.539

* Corresponding pounds active ingredient per acre for Product Use Rates (oz/A) in Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions. Also see Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions for the maximum product use rate per year in ozs/A and lb ai.

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Avocado*	Anthracnose (<i>Colletotrichum</i> spp.)	5 to 11	3	33	0

[Optional Text: * State-specific Restrictions - Not registered for use in California.]

Application Directions. Begin applications of **Merivon** prior to onset of disease development and continue on a 14-day interval.

For ground applications to avocado, use a minimum of 100 gallons of spray solution per acre.

Mechanically pressurized backpack sprayer applications may be used.

For aerial applications to avocado, use a minimum of 20 gallons of spray solution per acre.

Apply **Merivon** in sufficient spray volume to provide complete plant coverage without excessive runoff. Thorough plant coverage is required to maximize the effectiveness of **Merivon**.

DO NOT tank mix with other pesticide products, additives, nutrients, or any adjuvants other than NIS. When mixing with NIS, **DO NOT** exceed 1 pint/100 gallons (0.125% v/v).

All varieties and cultivars have not been tested with NIS tank mix combinations. Local conditions can also influence crop tolerance. Therefore, before tank mixing **Merivon** with NIS adjuvants, test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not result.

Restrictions

The minimum retreatment interval is 14 days.

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

DO NOT apply more than 11 fl ozs (0.180 lb fluxapyroxad 0.180 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs (0.539 lb fluxapyroxad, 0.539 lb pyraclostrobin) per acre per year.

DO NOT apply more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Berries and small fruits subgroups*</p> <p>Bushberry crop subgroup 13-07B Blueberry* (highbush and lowbush) Currant Elderberry Gooseberry Huckleberry Cultivars, varieties, and/or hybrids of these</p> <p>Caneberry crop subgroup 13-07A Blackberry (all varieties) Loganberry Raspberry (black and red) Wild raspberry Cultivars, varieties, and/or hybrids of these</p> <p>Low growing berry (except cranberry) crop subgroup 13-07G Bearberry Bilberry Cloudberry Lingonberry Muntries Partridgeberry Cultivars, varieties, and/or hybrids of these</p>	<p>Alternaria leaf spot and fruit rot (<i>Alternaria</i> spp.)</p> <p>Anthrachnose (<i>Colletotrichum</i> spp., <i>Elsinoe</i> spp.)</p> <p>Leaf spot and blotch (<i>Mycosphaerella</i> spp., <i>Septoria</i> spp.)</p> <p>Phomopsis leaf spot, twig blight, and fruit rot (<i>Phomopsis</i> spp.)</p> <p>Powdery mildew (<i>Sphaerotheca</i> spp., <i>Microshaera</i> spp., <i>Oidium</i> spp.)</p> <p>Spur blight (<i>Didymella</i> spp., <i>Phoma</i> spp.)</p> <p>Suppression only</p> <p>Rust (<i>Pucciniastrum</i> spp., <i>Arthuriomyces</i> spp., <i>Phragmidium</i> spp., <i>Kuehneola</i> spp.)</p>	<p>4 to 11</p>	<p>3</p>	<p>33</p>	<p>0 (bushberry, caneberry, and low growing berry)</p>
<p>Small fruit vine climbing (except fuzzy kiwifruit) crop subgroup 13-07F Amur river grape Gooseberry Kiwifruit, hardy Maypop Schisandra berry Cultivars, varieties, and/or hybrids of these</p>	<p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> <p>Monilinia blight and mummy berry (<i>Monilinia</i> spp.)</p>	<p>8 to 11</p>			<p>14 (small fruit vine climbing)</p>

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Berries and small fruits subgroups* *(continued)*

Application Directions. Begin applications of **Merivon** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Mix **Merivon** with water only for applications to crops listed in the berries and small fruits subgroups.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

DO NOT apply **Merivon** to any crops in the berries and small fruits subgroups, including blueberries (highbush and lowbush), as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

The minimum retreatment interval is 7 days.

DO NOT apply more than 11 fl ozs of **Merivon** (0.180 lb fluxapyroxad, 0.180 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs of **Merivon** (0.539 lb/acre fluxapyroxad, 0.539 lb/acre pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Brassica leafy vegetables crop subgroups 5A and 5B</p> <p>Head and stem crop subgroup 5A</p> <p>Broccoli Broccoli, Chinese Brussels sprouts Cabbage Cabbage, Chinese Cabbage, Chinese mustard Cauliflower Cavalo broccolo Kohlrabi</p> <p>Leafy greens crop subgroup 5B</p> <p>Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens</p>	<p>Alternaria leaf spot (<i>Alternaria</i> spp.)</p> <p>Anthrachnose (<i>Colletotrichum</i> spp.)</p> <p>Black leg (<i>Phoma lingam</i>)</p> <p>Cercospora leaf spot (<i>Cercospora brassicicola</i>)</p> <p>Powdery mildew (<i>Erysiphe polygoni</i>)</p> <p>Rhizoctonia blight (<i>Rhizoctonia solani</i>)</p> <p>Ring spot (<i>Mycosphaerella brassicicola</i>)</p> <p>White leaf spot (<i>Pseudocercospora capsellae</i>)</p> <p>White rust (<i>Albugo candida</i>)</p> <p>Suppression only</p> <p>Downy mildew (<i>Peronospora parasitica</i>)</p>	<p>4 to 5.5</p>	<p>3</p>	<p>16.5</p>	<p>3</p>

Application Directions. Begin applications of **Merivon** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 16.5 fl ozs of **Merivon** (0.269 lb fluxapyroxad, 0.269 lb pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Bulb vegetables crop group 3-07 Chive, Chinese, fresh leaves Chive, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties, and/or hybrids of these	Powdery mildew (<i>Leveillula taurica</i>) Purple blotch and leaf blight (<i>Alternaria porri</i>) Stemphylium leaf blight and stalk rot (<i>Stemphylium vesicarium</i>)	4 to 11	3	33	7
	Botrytis leaf blight (<i>Botrytis</i> spp.) Botrytis neck rot (<i>Botrytis</i> spp.) Suppression only Downy mildew (<i>Peronospora destructor</i>)	8 to 11			

Application Directions. Begin applications of **Merivon** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 11 fl ozs of **Merivon** (0.180 lb fluxapyroxad, 0.180 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs of **Merivon** (0.539 lb/acre fluxapyroxad, 0.539 lb/acre pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Cucurbit vegetables crop group 9 Chayote Chinese waxgourd Citron melon Cucumber Gherkin Pumpkin Watermelon</p> <p>Edible gourd Chinese okra Cucuzza Hyotan</p> <p>Momordica spp. Balsam apple Balsam pear Bitter melon Chinese cucumber</p> <p>Muskmelon Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persian melon Pineapple melon Santaclaus melon Snake melon</p> <p>Summer squash Crookneck squash Scallop squash Straightneck squash Vegetable marrow Zucchini</p> <p>Winter squash Acorn squash Butternut squash Calabaza Hubbard squash Spaghetti squash</p>	<p>Alternaria leaf blight (<i>Alternaria cucumerina</i>)</p> <p>Powdery mildew (<i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp., <i>Erysiphe</i> spp.)</p>	<p>4 to 5.5</p>	<p>3</p>	<p>16.5</p>	<p>0</p>
	<p>Anthracnose (<i>Colletotrichum orbiculare</i>)</p> <p>Cercospora leaf spot (<i>Cercospora citrulina</i>)</p> <p>Gummy stem blight (<i>Didymella bryoniae</i>)</p> <p>Microdochium blight (<i>Plectosporium tabacinum</i>)</p> <p>Target leaf spot (<i>Corynespora cassiicola</i>)</p> <p>Suppression only</p> <p>Downy mildew (<i>Pseudoperonospora cubensis</i>)</p>	<p>5.5</p>			

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Cucurbit vegetables crop group 9 *(continued)*

Application Directions. For optimal disease control, begin applications of **Merivon** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Mix **Merivon** with water only for applications to crops listed in the cucurbit vegetables group.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above. See the restriction below regarding gummy stem blight resistance.

Restrictions

DO NOT apply **Merivon** to any crops in the cucurbit vegetables group as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 16.5 fl ozs of **Merivon** (0.269 lb fluxapyroxad, 0.269 lb pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than one (1) application of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide. In areas where gummy stem blight resistance to **Group 7** or **Group 11** fungicides have been confirmed, tank mix with chlorothalonil at full label rates and adhering to all label precautions.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Fruiting vegetables crop group 8-10 African eggplant Bush tomato Cocona Currant tomato Eggplant Garden huckleberry Goji berry Groundcherry Martynia Naranjilla Okra Pea eggplant Pepino	Anthracnose (<i>Colletotrichum coccodes</i>)	4 to 5.5 or 4 to 5.5 fl ozs per 100 gallons of spray volume (dilute)*	3	16.5	0
	Black mold (<i>Alternaria alternata</i>)				
	Early blight (<i>Alternaria solani</i>)				
	Powdery mildew (<i>Leveillula taurica</i>)				
	Septoria leaf spot (<i>Septoria lycopersici</i>)				
	Target spot (<i>Corynespora cassiicola</i>)				
	Suppression only	4 to 5.5 or 4 to 5.5 fl ozs per 100 gallons of spray volume (dilute)*			
	Botrytis gray mold (<i>Botrytis cinerea</i>)				
	Rhizoctonia stem rot** (<i>Rhizoctonia solani</i>)				
	Sclerotinia stem rot, white mold (<i>Sclerotinia sclerotiorum</i>)				
	Southern blight** (<i>Sclerotium rolfsii</i>)				
	Suppression only	5.5			
	Late blight (<i>Phytophthora infestans</i>)				
Pepper (all varieties) Pepper, bell Pepper, nonbell Roselle Scarlet eggplant Sunberry Tomatillo Tomato Tree tomato Cultivars, varieties and/or hybrids of these					

Application Directions. For optimal disease control, begin applications of **Merivon** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Applications using drip irrigation systems may provide disease suppression. The level and consistency of suppression from drip line applications varies with the soil type, level of inoculum, irrigation volumes, environment and other factors. See your local BASF representative for details on drip irrigation use in your area.

* For applications based on dilute volume, plants must be sprayed to runoff. Apply a minimum of 20 gallons of spray volume per acre, and increase the spray volume as the plants grow during the season. Spray volume must be proportional to the amount of plant tissue to be covered such that 100 gallons of spray per acre is used on mature plants.

Use of Adjuvants and Other Products as Mixes with Merivon on fruiting vegetables.

Merivon can be used with nonionic surfactants at their lowest label rate up to 0.125%. When **Merivon** is mixed with buffering agents and foliar nutrients, the pH of the final spray solution must be greater than 5.5.

DO NOT mix **Merivon** with the following products on fruiting vegetables:

- Emulsifiable concentrate (EC) formulation or solvent-based formulation products.
- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS) or MSO/OS blended adjuvant products.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Fruiting vegetables crop group 8-10 *(continued)*

For **Merivon** applications to **fresh market tomatoes** at less than 20 gallons per acre, **DO NOT** mix **Merivon** with any other products, adjuvants, additives, nutrients or anything other than water.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Merivon** may result from mixing **Merivon** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF advises testing **Merivon** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 16.5 fl ozs of **Merivon** (0.269 lb fluxapyroxad, 0.269 lb pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

** Not registered for use in California.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Grapes</p> <p>Table</p> <p>Wine and raisin</p> <p>See Crop Safety Precaution below</p>	<p>Angular leaf spot (<i>Mycosphaerella angulata</i>)</p> <p>Anthracnose (<i>Elsinoe ampelina</i>)</p> <p>Black rot (<i>Guignardia bidwellii</i>)</p> <p>Leaf blight (<i>Pseudocercospora vitis</i>)</p> <p>Phomopsis cane and Leaf spot (<i>Phomopsis viticola</i>)</p> <p>Powdery mildew (<i>Erysiphe necator</i>)</p> <p>Ripe rot (<i>Colletotrichum gloeosporioides</i>)</p> <p>Suppression only</p> <p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> <p>Downy mildew (<i>Plasmopara viticola</i>)</p>	4 to 5.5	6	33	14
<p>Application Directions. DO NOT allow workers to perform cane tying and leaf pulling tasks for 5 days after application. Notify workers of this prohibition.</p> <p>For powdery mildew control, begin applications of Merivon at bud break or prior to onset of disease and continue on 28 day intervals. The minimum retreatment interval is 28 days for powdery mildew control.</p> <p>For black rot control and downy mildew suppression, begin applications of Merivon before pre-bloom and continue on 28 day intervals. The minimum retreatment interval is 28 days for black rot control and downy mildew suppression.</p> <p>For all other diseases listed, begin applications of Merivon prior to onset of disease development and continue on 28 day intervals.</p>					

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Grapes *(continued)*

Use of Adjuvants and Other Products as Mixes with Merivon.

The use of adjuvants or additives may improve the performance of **Merivon** on **wine and raisin** grapes. However, under certain conditions, mixtures of **Merivon** with adjuvants, additives and/or other products may cause crop injury. Caution must be exercised if **Merivon** is tank mixed with products containing high amounts of solvents as crop response to actively growing foliage and/or fruit can occur under certain conditions.

The use of citrus oils or adjuvants containing methylated seed oil (MSO) may increase the likelihood of necrotic spots and/or marginal necrosis on active growing leaves of certain sensitive varieties.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Merivon** may result from mixing **Merivon** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF advises testing **Merivon** in combination with other products for crop safety on a small portion of young foliage before application to the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult your local BASF representative for more information specific to your area and specific conditions that may cause crop response.

Crop Safety Precaution. DO NOT use Merivon on Concord, Noiret and NY73.0136.17 varieties due to possible foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related varieties. Not all varieties have been thoroughly tested.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

The minimum retreatment intervals are listed above with their corresponding diseases.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs of **Merivon** (0.539 lb/acre fluxapyroxad, 0.539 lb/acre pyraclostrobin) per acre per year.

DO NOT make more than 6 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

Mix **Merivon** with water only for applications to **table grapes**.

DO NOT mix with EC formulation pesticides.

DO NOT mix with petroleum oil-based products such as mineral and paraffinic oil (including phytobland oil) or with blended adjuvants containing petroleum oil-based products, e.g., stilet oil, crop oil concentrate and crop oil (emulsifiable).

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Leafy vegetables (except Brassica) crop group 4 Amaranth Arugula Cardoon Celery Celery, Chinese Celtuce Chervil Chrysanthemum (edible-leaved and garland) Corn salad Cress (garden and Upland) Dandelion Dock Endive Fennel, Florence Lettuce (head and leaf) Orach Parsley Purslane (garden and winter)	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Ascochyta leaf spot (<i>Ascochyta</i> spp.) Cercospora leaf spot (<i>Cercospora</i> spp.) Phoma (<i>Phoma</i> spp.) Powdery mildew (<i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.) Rust (<i>Puccinia</i> spp., <i>Uromyces</i> spp.) Septoria leaf spot (<i>Septoria</i> spp.) White rust (<i>Albugo</i> spp.)	4 to 11	3	33	1
Radicchio (red chicory) Rhubarb Spinach	Suppression only Downy mildew (<i>Peronospora</i> spp., <i>Pseudoperonospora</i> spp.)	6 to 11			
Spinach (New Zealand and vine) Swiss chard	Botrytis rot (<i>Botrytis</i> spp.) Lettuce downy mildew (<i>Bremia</i> spp.) Lettuce drop caused by <i>Sclerotinia minor</i>	8 to 11			

Application Directions. Begin applications of **Merivon** prior to onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Tank Mix Restrictions

Spinach (all varieties). **DO NOT** apply **Merivon** to spinach as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives, or anything other than water.

Mix **Merivon** with water only for applications to spinach (all varieties).

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Leafy vegetables (except Brassica) crop group 4 *(continued)*

Leafy vegetables (except spinach). It is impossible for BASF to test all varieties of leafy vegetables for sensitivity to **Merivon** under all environments and all potential product mixture combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Merivon** use, particularly in tank mixes and/or adjuvant combinations on leafy vegetables. To reduce the risk of leafy vegetable injury, BASF advises testing **Merivon** or **Merivon** tank mixtures on a small portion of the crop before broadscale use.

To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Merivon** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 11 fl ozs of **Merivon** (0.180 lb fluxapyroxad, 0.180 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs of **Merivon** (0.539 lb/acre fluxapyroxad, 0.539 lb/acre pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oilseed crop group 20 Flax seed	PasmO <i>(Septoria linicola)</i>	4 to 5.5	2	11	21
Rapeseed (cultivars, varieties, and/or hybrids, including canola and crambe)	Blackleg <i>(Leptosphaeria maculans)</i> Blackspot <i>(Alternaria spp.)</i> Suppression only White mold/Sclerotinia stem rot <i>(Sclerotinia sclerotiorum)</i>				
Sunflower	Alternaria leaf spot <i>(Alternaria spp.)</i> Cercospora leaf spot <i>(Cercospora helianthi)</i> Powdery mildew <i>(Erysiphe cichoracearum)</i> Rust <i>(Puccinia helianthi, Uromyces spp.)</i> Septoria leaf spot <i>(Septoria spp.)</i> White rust <i>(Albugo tragopogonis)</i> Suppression only Sclerotinia head blight <i>(Sclerotinia sclerotiorum)</i>				

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Oilseed crop group 20</p> <p>Borage Castor oil plant Calendula Chinese tallowtree Cottonseed* Crambe Cuphea Echium Euphorbia Evening primrose Gold of pleasure (camelina) Hare’s ear mustard Jojoba Lesquerella Lunaria Meadowfoam Milkweed Mustard seed Niger seed Oil radish Poppy seed Rose hip Safflower Sesame Stokes aster Sweet rocket Tallowwood Tea oil plant Vernonia</p>	<p><i>Alternaria</i> spp. <i>Septoria</i> spp. Suppression only <i>Sclerotinia</i> spp.</p>	<p>4 to 5.5</p>	<p>2</p>	<p>11</p>	<p>21</p>

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Oilseedcrop group 20 *(continued)*

Application Directions for Rapeseed. For the control of blackleg, apply **Merivon** at the 2 to 4 leaf stage. For optimal control of blackspot, apply **Merivon** at early pod development. For suppression of Sclerotinia, apply **Merivon** at 20% to 50% flowering or prior to the onset of disease. A second application may be made 14 days later if weather conditions are favorable for disease development.

Application Directions for Other Oilseed Crops. Begin applications of **Merivon** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

No livestock feeding restrictions.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 11 fl ozs of **Merivon** (0.180 lb fluxapyroxad, 0.180 lb pyraclostrobin) per acre per year.

DO NOT make more than 2 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Pome fruits crop group 11-10</p> <p>Apple Azarole Crabapple Loquat Mayhaw Medlar Pear Pear, Asian Pear, Oriental Quince Quince, Chinese Quince, Japanese Tejocote Cultivars, varieties and/or hybrids of these</p>	<p>Alternaria blotch (<i>Alternaria mali</i>)</p> <p>Apple scab (<i>Venturia inaequalis</i>)</p> <p>Bitter rot (<i>Colletotrichum</i> spp.)</p> <p>Black rot/Frogeye leaf spot (<i>Botryosphaeria obtusa</i>)</p> <p>Blue mold* (<i>Penicillium</i> spp.)</p> <p>Brooks spot (<i>Mycosphaerella pomi</i>)</p> <p>Flyspeck (<i>Zygophiala jamaicensis</i>)</p> <p>Gray mold rot* (<i>Botrytis cinerea</i>)</p> <p>Pear scab (<i>Venturia pirina</i>)</p> <p>Powdery mildew (<i>Podosphaera leucotricha</i>)</p> <p>Sooty blotch (disease complex)</p> <p>White rot (<i>Botryosphaeria dothidea</i>)</p> <p>Suppression only</p> <p>Cedar apple rust (<i>Gymnosporangium juniperi-virginianae</i>)</p> <p>Quince rust (<i>Gymnosporangium clavipes</i>)</p>	4 to 5.5	4	22	0

Application Directions. For scab, powdery mildew, frogeye leafspot and rust, begin applications of **Merivon** prior to disease development and continue on a 7 to 10 day interval.

For sooty blotch, flyspeck, white rot, black rot, bitter rot and Alternaria blotch, begin applications of **Merivon** prior to disease development and continue on a 7 to 14 day interval.

Use of Adjuvants and Other Products as Mixes with Merivon.

The use of adjuvants or additives may improve the performance of **Merivon** on pome fruits. However, under certain conditions, mixtures of **Merivon** with adjuvants, additives and/or other products may cause crop injury. Caution must be exercised if **Merivon** is tank mixed with products formulated as emulsifiable concentrates (EC) or containing high amounts of solvents since injury may occur. Consult your local BASF representative for more information specific to your area and specific conditions that may cause crop injury.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Pome fruits crop group 11-10 *(continued)*

DO NOT use **Merivon** with:

- Crop oil concentrate (COC), methylated seed oil (MSO) adjuvants

For **pears**, **DO NOT** use **Merivon** with horticultural mineral oil as crop response to foliage and/or fruit can occur under certain conditions.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Merivon** may result from mixing **Merivon** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF advises testing **Merivon** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

No restriction on livestock grazing or feeding.

For aerial application to pome fruit trees, use a minimum of 10 gallons of spray solution per acre.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 22 fl ozs of **Merivon** (0.359 lb fluxapyroxad, 0.359 lb pyraclostrobin) per acre per year.

DO NOT make more than 4 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Pomegranate	Alternaria black heart <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i>	7	2	14	14

Application Directions. Begin applications of **Merivon** prior to onset of disease development and continue on a 14 day interval.

Apply **Merivon** in sufficient spray volume to provide complete plant coverage without excessive runoff. Thorough plant coverage is required to maximize the effectiveness of **Merivon**.

East of the Rocky Mountains: To avoid crop injury, **DO NOT** apply **Merivon** to pomegranates as a tank mix with any adjuvants, liquid fertilizers, nutrients, or anything other than water.

West of the Rocky Mountains: Merivon may be tank mixed with NIS adjuvants at a rate not to exceed 8 fl ozs/100 gallons (0.0625% v/v). **DO NOT** tank mix with other pesticide products, additives, nutrients, or any adjuvants other than NIS. All varieties and cultivars have not been tested with NIS tank mix combinations. Local conditions can also influence crop tolerance. Therefore, before tank mixing **Merivon** with NIS adjuvants, test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not result.

Restrictions

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

The minimum retreatment interval is 7 days.

DO NOT apply more than 7 fl ozs (0.114 lb fluxapyroxad 0.114 lb pyraclostrobin) per acre per application.

DO NOT apply more than 14 fl ozs (0.228 lb fluxapyroxad, 0.228 lb pyraclostrobin) per acre per year.

DO NOT apply more than 2 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Root vegetables (except sugar beet) crop subgroup 1B* Beet, garden Burdock, edible Carrot Celeriac Chervil, turnip-rooted Chicory Ginseng Horseradish Parsley, turnip-rooted Parsnip Radish Radish, Oriental Rutabaga Salsify Salsify, black Salsify, Spanish Skirret Turnip	Alternaria leaf spot/blight (<i>Alternaria</i> spp.) Powdery mildew (<i>Erysiphe</i> spp., <i>Leveillula</i> spp.)	4 to 5.5	3	16.5	7
	Cercospora leaf spot/blight (<i>Cercospora</i> spp.)	5.5			

Application Directions. For optimal disease control, begin applications of **Merivon** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 16.5 fl ozs of **Merivon** (0.269 lb fluxapyroxad, 0.269 lb pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California, except on carrot.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Soybean*	Alternaria leaf spot (<i>Alternaria</i> spp.)	4 to 5.5	2	11	21
	Anthracnose (<i>Colletotrichum truncatum</i>)				
	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)				
	Brown spot (<i>Septoria glycines</i>)				
	Cercospora blight (<i>Cercospora kikuchii</i>)				
	Frogeye leaf spot (<i>Cercospora sojina</i>)				
	Pod and stem blight (<i>Diaporthe phaseolorum</i>)				
	Rhizoctonia aerial blight (<i>Rhizoctonia solani</i>)				
	Suppression only	5.5			
	Sclerotinia blight (white mold) (<i>Sclerotinia sclerotiorum</i>)				
	Southern blight (<i>Sclerotium rolfsii</i>)				

Application Directions. For optimal disease control, begin applications of **Merivon** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

For adequate control of soybean rust, apply **Merivon** prior to infection.

Merivon may be used with adjuvants for application to soybeans. See the **Additives and Tank Mixing Information** and **Mixing Order** sections for more details.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 11 fl ozs of **Merivon** (0.180 lb fluxapyroxad, 0.180 lb pyraclostrobin) per acre per year.

DO NOT make more than 2 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

Soybean forage may be fed no sooner than 14 days after last application. Soybean hay may be fed no sooner than 21 days after last treatment.

* Not registered for use in California.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Stone fruits crop group 12 Apricot Cherry (sweet and tart) Nectarine Peach Plum (all varieties) Plumcot Prune	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Blossom blight (<i>Monilinia</i> spp.) Blue mold (<i>Penicillium</i> spp.) Brown rot (<i>Monilinia</i> spp.) Gray mold (<i>Botrytis</i> spp.) Leaf spot (<i>Blumeriella jaapii</i>) Powdery mildew (<i>Sphaerotheca</i> spp., <i>Podosphaera</i> spp.) Ripe fruit rot (<i>Monilinia fruticola</i> , <i>Monilinia laxa</i> , <i>Botrytis cinerea</i> , <i>Rhizopus</i> spp.) Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) Shothole (<i>Wilsonomyces carpophilus</i>)	4 to 6.7	3	20.1	0

Application Directions. For optimal disease control, begin application of **Merivon** at pink bud or prior to the onset of disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Use of Adjuvants and Other Products as Mixes with Merivon.

The use of adjuvants or additives may improve the performance of **Merivon** on stone fruits. However, under certain conditions, mixtures of **Merivon** with adjuvants, additives and/or other products may cause crop injury, particularly to fruit within two weeks of harvest. Caution must be exercised if **Merivon** is tank mixed with products formulated as emulsifiable concentrates (EC) or containing high amounts of solvents since injury may occur. Consult your local BASF representative for more information specific to your area and specific conditions that may cause crop injury.

Up to Two Weeks Before Harvest

Merivon can be used with nonionic, organosilicone or **blended** MSO/OS (methylated seed oil/organosilicone) adjuvants.

DO NOT use **Merivon** with:

- Crop oil concentrate (COC), methylated seed oil (MSO) adjuvants.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Stone fruits crop group 12 *(continued)*

Within Two Weeks of Harvest - Use on Cherries (See below for other Stone Fruit crops)

Merivon can only be used with nonionic adjuvants that **DO NOT** acidify and/or enhance penetration. **DO NOT** exceed 0.0625% (v/v) concentration when using these adjuvants.

DO NOT use **Merivon** with:

- Emulsifiable concentrate (EC) or solvent-based formulation products.
- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS), MSO/OS blend adjuvants.
- Nonionic surfactant (NIS) adjuvant products that acidify or enhance plant penetration.

Within Two Weeks of Harvest - Use on Stone Fruit crops other than Cherries

DO NOT use **Merivon** with:

- Crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS), MSO/OS blend adjuvants.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Merivon** may result from mixing **Merivon** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF advises testing **Merivon** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

For aerial application to stone fruit trees, use a minimum of 10 gallons of spray solution per acre.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

The minimum retreatment interval is 7 days.

DO NOT apply more than 6.7 fl ozs of **Merivon** (0.109 lb fluxapyroxad, 0.109 lb pyraclostrobin) per acre per application.

DO NOT apply more than 20.1 fl ozs of **Merivon** (0.328 lb fluxapyroxad, 0.328 lb pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Strawberries	Leaf spot <i>(Mycosphaerella fragariae, Ramularia tulasnei)</i> Powdery mildew <i>(Sphaerotheca macularis)</i>	4 to 7	3	33	0
	Anthracnose <i>(Colletotrichum spp.)</i>	5.5 to 8			
	Botrytis gray mold <i>(Botrytis cinerea)</i>	8 to 11			

Application Directions. Begin applications of **Merivon** no later than 10% bloom, or prior to disease development and continue on a 7 to 14 day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 11 fl ozs of **Merivon** (0.180 lb fluxapyroxad, 0.180 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs of **Merivon** (0.539 lb/acre fluxapyroxad, 0.539 lb/acre pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p>Tree nuts crop group 14</p> <p>Almond Beechnut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert Hickory nut Macadamia nut Pecan Pistachio Walnut, black Walnut, English</p>	<p>Anthracnose (<i>Colletotrichum</i> spp.)</p> <p>Botrytis blossom and shoot blight (<i>Botrytis cinerea</i>)</p> <p>Brown rot/Blossom blight (<i>Monilinia</i> spp.)</p> <p>Eastern filbert blight* (<i>Anisogramma anomala</i>)</p> <p>Green fruit rot/Jacket rot (<i>Botrytis cinerea</i>, <i>Sclerotinia sclerotiorum</i>, <i>Monilinia laxa</i>)</p> <p>Leaf rust (<i>Tranzschelia discolor</i>)</p> <p>Panicle and shoot blight (<i>Botryosphaeria dothidea</i>)</p> <p>Scab (<i>Cladosporium carpophilum</i>, <i>C. caryigenum</i>)</p> <p>Shothole (<i>Wilsonomyces carpophilus</i>)</p> <p>Suppression only</p> <p>Hull rot (<i>Aspergillus</i> spp.*, <i>Monilinia</i> spp., <i>Rhizopus stolonife</i>)</p>	<p>5 to 6.5</p>	<p>3</p>	<p>19.5</p>	<p>14</p>
	<p>Alternaria late blight (<i>Alternaria</i> spp.)</p>	<p>6.5</p>			

[Optional Text: * State-specific Restrictions - Not registered for use in California.]

Application Directions. For almond, begin applications of **Merivon** prior to the onset of disease development, and continue on a 7 to 14 day interval up to 14 days before harvest.

For filbert, begin applications at budswell to budbreak, or prior to infection and onset of disease development. Continue on a 7 to 14 day interval to cover and protect new growth.

For pecan, begin applications of **Merivon** prior to onset of disease development and continue on a 7 to 21 day interval for the control of scab. Use the shorter interval and/or the higher rate when disease pressure is high.

For pistachio, apply **Merivon** prior to onset of disease development and continue on a 10 to 30 day interval. Use the higher rate and shorter intervals when disease pressure is high.

For all other crops listed, apply **Merivon** prior to disease development and continue on a 7 to 28 day interval. In all cases use the shorter interval when shoot growth is rapid.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Tree nuts crop group 14 *(continued)*

Mechanically pressurized backpack sprayer applications may be used.

No restriction on livestock feeding of almond hulls.

For aerial application to tree nuts, **DO NOT** use less than 10 gallons of spray solution per acre.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

DO NOT apply more than 6.5 fl ozs of **Merivon** (0.106 lb fluxapyroxad, 0.106 lb pyraclostrobin) per acre per application.

DO NOT apply more than 19.5 fl ozs of **Merivon** (0.318 lb fluxapyroxad, 0.318 lb pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* Not registered for use in California.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Tuberous and corm vegetables crop subgroup 1C Arracacha Arrowroot Cassava (bitter and sweet) Chayote (root) Chinese artichoke Chufa Dasheen (taro) Edible canna Ginger Jerusalem artichoke Leren Potato Sweet potato Tanier True yam Turmeric Yam bean	Brown spot and black pit** <i>(Alternaria alternata)</i> Early blight <i>(Alternaria solani)</i> Suppression only Botrytis gray mold** <i>(Botrytis cinerea)</i>	3 to 5.5	3	16.5*	7
	Black dot <i>(Colletotrichum coccodes)</i> Leaf spot <i>(Cercospora spp.)</i> Powdery mildew <i>(Erysiphe spp., Leveillula taurica)</i>	4 to 5.5			
	Rust <i>(Uromyces spp., Puccinia spp.)</i> Suppression only Late blight** <i>(Phytophthora infestans)</i> White mold <i>(Sclerotinia sclerotiorum)</i>	5.5			

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Tuberous and corm vegetables crop subgroup 1C *(continued)*

Application Directions. For optimal disease control, begin applications of **Merivon** prior to disease development and continue on a 7 to 14 day interval. The lower rate and longer interval can be used early season prior to the observance of symptoms and when disease pressure is low.

Use the higher rates and shorter intervals once disease has been confirmed in your area or weather conditions are conducive to disease development.

Use of Adjuvants and Other Products as Mixes with Merivon. The use of additives or adjuvants may improve the performance of **Merivon** on tuberous and corm vegetables.

However, BASF evaluations also indicate that under some conditions, application of **Merivon** in combination with certain rates of organosilicone-based or oil-containing (petroleum, crop or methylated seed oil) additives or adjuvants can cause crop injury. Crop injury also may result from applications of **Merivon** mixed with other products that have solvent-based formulations that increase penetration.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Merivon** may result from mixing **Merivon** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF advises testing **Merivon** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

(continued)

Table 3. Merivon® Xemium® brand fungicide Crop-specific Directions *(continued)*

Tuberous and corm vegetables crop subgroup 1C *(continued)*

No livestock feeding restrictions.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

The minimum retreatment interval is 7 days.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 16.5 fl ozs of **Merivon** (0.269 lb fluxapyroxad, 0.269 lb pyraclostrobin) per acre per year.

DO NOT make more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

* The maximum product rate per year includes the combination of infurrow and foliar uses (for above-listed crops, infurrow use is permitted in potato only).

Instructions for Infurrow Use to Aid in the Control of Soilborne Rhizoctonia in Potatoes

Use 0.4 fl oz of **Merivon** per 1000 row feet (for applications on 32-inch or 34-inch rows, the maximum application rate is 0.325 fl oz/1000 row feet). Refer to the chart below to determine the rate per acre. Apply at planting as an infurrow spray by directing spray pattern to uniformly cover seed pieces and surrounding soil. The spray pattern must be a 4 to 8 inch band that is applied to the seed piece prior to being covered with soil.

When Rhizoctonia disease pressure conditions are expected to be severe or if the field has a history of Rhizoctonia infestations, use **Merivon** at 0.4 fl oz product per 1000 row feet and/or tank mix with a fungicide having a different mode of action.

Use a minimum volume of application of 5 gallons of water per acre.

** Not registered for use in California.

Product Rate per 1000 row feet (fl oz product)	Product Rate per Acre (fl ozs product)			
	34-inch rows or less	36-inch rows	38-inch rows	40-inch rows
0.4	See footnote ¹	5.4	5.1	4.8

¹ For 34-inch rows or less, use a maximum of 0.325 fl oz product per 1000 row feet.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Merivon® Xemium® brand fungicide is specially formulated and sold by BASF for the control of pests according to the directions on this label. The purchase price of **Merivon** includes a royalty whereby the purchaser acquires a prepaid license under U.S. Patent No. 5,438,070 and (pending No. 2008/0153707 A1) under which purchaser agrees to employ the purchased quantity of **Merivon** only for the above-specified uses under BASF's United States patent rights and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of **Merivon** other than those specified on this label are not licensed through the purchase of this product and the use of this product for other purposes may violate this license and patent rights of BASF.

***F500, Merivon and Xemium** are registered trademarks of BASF.*

© 2023 BASF Corporation
All rights reserved.

007969-00310.20230808e.NVA 2023-04-0371-0062

Supersedes: NVA 2020-04-371-0255

Supplementals: NVA 2021-04-371-0186

NVA 2023-04-0371-0120

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry

ACCEPTED

01/18/2024

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7969-310

Fluxapyroxad	Group	7	Fungicide
Pyraclostrobin	Group	11	Fungicide

Supplemental Label

Merivon®

Xemium® Brand Fungicide

For disease control and plant health in avocado

This supplemental label expires on August 31, 2026, and must not be used or distributed after this date.

Powered by **Xemium®** and **F500®** fungicides

Active Ingredients:

fluxapyroxad*: 1 <i>H</i> -Pyrazole-4-carboxamide, 3-(difluoromethyl)-1-methyl-N-(3',4',5'-trifluoro[1,1'-biphenyl]-2-yl)-	21.26%
pyraclostrobin**: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1 <i>H</i> -pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester)	21.26%

Other Ingredients:..... 57.48%

Total: 100.00%

* Equivalent to 2.09 pounds of fluxapyroxad per gallon

** Equivalent to 2.09 pounds of pyraclostrobin per gallon

EPA Reg. No. 7969-310
WARNING/AVISO

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Merivon® Xemium® brand fungicide** container label, EPA Reg. No. 7969-310, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Merivon** before applying.
- Use of **Merivon** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Merivon**.

Application Instructions

Refer to the **Merivon** container label for specific use directions, recommendations and restrictions in registered crops.

Personal Protective Equipment (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils)

Use Restrictions and Limitations

- **Merivon** is not for sale, distribution or use in Nassau and Suffolk counties in New York State.

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709


We create chemistry

Table 1. Merivon® Xemium® brand fungicide Restrictions and Limitations Overview*

Crop	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Number of Sequential Applications	Maximum Product Rate per Year (fl ozs/A)
Avocado	0	5 to 11	3	2	33

* See Table 2. Merivon Crop-specific Directions for additional directions.

Table 2. Merivon Crop-specific Directions

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Avocado	Anthrachnose (<i>Colletotrichum</i> spp.)	5 to 11	3	33	0

Application Directions. Begin applications of **Merivon** prior to onset of disease development and continue on a 14-day interval.

For ground applications to avocado, use a minimum of 100 gallons of spray solution per acre.

Mechanically pressurized backpack sprayer applications may be used.

For aerial applications to avocado, use a minimum of 20 gallons of spray solution per acre.

Apply **Merivon** in sufficient spray volume to provide complete plant coverage without excessive runoff. Thorough plant coverage is required to maximize the effectiveness of **Merivon**.

DO NOT tank mix with other pesticide products, additives, nutrients, or any adjuvants other than NIS. When mixing with NIS, **DO NOT** exceed 1 pint/100 gallons (0.125% v/v).

All varieties and cultivars have not been tested with NIS tank mix combinations. Local conditions can also influence crop tolerance. Therefore, before tank mixing **Merivon** with NIS adjuvants, test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not result.

Restrictions

The minimum retreatment interval is 14 days.

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

DO NOT apply more than 11 fl ozs (0.180 lb fluxapyroxad 0.180 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs (0.539 lb fluxapyroxad, 0.539 lb pyraclostrobin) per acre per year.

DO NOT apply more than 3 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Merivon® Xemium® brand fungicide is specially formulated and sold by BASF for the control of pests according to the directions on this label. The purchase price of **Merivon** includes a royalty whereby the purchaser acquires a prepaid license under U.S. Patent No.5,438,070 and (pending No. 2008/0153707 A1) under which purchaser agrees to employ the purchased quantity of **Merivon** only for the above-specified uses under BASF's United States patent rights and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of **Merivon** other than those specified on this label are not licensed through the purchase of this product and the use of this product for other purposes may violate this license and patent rights of BASF.

F500, Merivon, and Xemium are registered trademarks of BASF.

© 2023 BASF Corporation
All rights reserved.

007969-00310.20211022e.NVA 2021-04-371-0186

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry

ACCEPTED

01/18/2024

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7969-310

Fluxapyroxad	Group	7	Fungicide
Pyraclostrobin	Group	11	Fungicide

Supplemental Label

Merivon[®]

Xemium[®] Brand Fungicide

For disease control and plant health in grapes

This supplemental label expires on August 31, 2026, and must not be used or distributed after this date.

Powered by **Xemium[®]** and **F500[®]** fungicides

Active Ingredients:

fluxapyroxad*: 1 <i>H</i> -Pyrazole-4-carboxamide, 3-(difluoromethyl)-1-methyl-N-(3',4',5'-trifluoro[1,1'-biphenyl]-2-yl)-	21.26%
pyraclostrobin**: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1 <i>H</i> -pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester)	21.26%

Other Ingredients:..... 57.48%

Total: 100.00%

* Equivalent to 2.09 pounds of fluxapyroxad per gallon

** Equivalent to 2.09 pounds of pyraclostrobin per gallon

EPA Reg. No. 7969-310

WARNING/AVISO

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Merivon[®] Xemium[®] brand fungicide** container label, EPA Reg. No. 7969-310, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Merivon** before applying.
- Use of **Merivon** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Merivon**.

Application Instructions

Refer to the **Merivon** container label for specific use directions, recommendations and restrictions in registered crops.

Personal Protective Equipment (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils)

Use Restrictions and Limitations

- **Merivon** is not for sale, distribution or use in Nassau and Suffolk counties in New York State.

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709


We create chemistry

Table 1. Merivon® Xemium® brand fungicide Restrictions and Limitations Overview*

Crop	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Number of Sequential Applications	Maximum Product Rate per Year (fl ozs/A)
Grapes	14	5.5	6	2	33

* See Table 2. Merivon® Xemium® brand fungicide Crop-specific Directions for additional directions.

Table 2. Merivon® Xemium® brand fungicide Crop-specific Directions

Crop	Target Disease	Product Use Rate per Application (fl ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (fl ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Grapes Table Wine and raisin See Crop Safety Precaution below	Angular leaf spot (<i>Mycosphaerella angulata</i>) Anthracnose (<i>Elsinoe ampelina</i>) Black rot (<i>Guignardia bidwellii</i>) Leaf blight (<i>Pseudocercospora vitis</i>) Phomopsis cane and Leaf spot (<i>Phomopsis viticola</i>) Powdery mildew (<i>Erysiphe necator</i>) Ripe rot (<i>Colletotrichum gloeosporioides</i>) <u>Suppression only</u> Botrytis gray mold (<i>Botrytis cinerea</i>) Downy mildew (<i>Plasmopara viticola</i>)	4 to 5.5	6	33	14

Application Directions. DO NOT allow workers to perform cane tying and leaf pulling tasks for 5 days after application. Notify workers of this prohibition.

For powdery mildew control, begin applications of **Merivon** at bud break or prior to onset of disease and continue on 28 day intervals. The minimum retreatment interval is 28 days for powdery mildew control.

For black rot control and downy mildew suppression, begin applications of **Merivon** before pre-bloom and continue on 28 day intervals. The minimum retreatment interval is 28 days for black rot control and downy mildew suppression.

For all other diseases listed, begin applications of **Merivon** prior to onset of disease development and continue on 28 day intervals.

(continued)

Table 2. Merivon® Xemium® brand fungicide Crop-specific Directions (continued)

Grapes (continued)

Use of Adjuvants and Other Products as Mixes with Merivon.

The use of adjuvants or additives may improve the performance of **Merivon** on **wine and raisin** grapes. However, under certain conditions, mixtures of **Merivon** with adjuvants, additives and/or other products may cause crop injury. Caution must be exercised if **Merivon** is tank mixed with products containing high amounts of solvents as crop response to actively growing foliage and/or fruit can occur under certain conditions.

The use of citrus oils or adjuvants containing methylated seed oil (MSO) may increase the likelihood of necrotic spots and/or marginal necrosis on active growing leaves of certain sensitive varieties.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, crop injury, or incompatibility due to additives, adjuvants or other products used in combination with **Merivon** may result from mixing **Merivon** with other products. Refer also to the **Conditions of Sale and Warranty** section of this label.

To minimize the likelihood of crop injury, BASF advises testing **Merivon** in combination with other products for crop safety on a small portion of young foliage before application to the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult your local BASF representative for more information specific to your area and specific conditions that may cause crop response.

Crop Safety Precaution. DO NOT use Merivon on Concord, Noiret and NY73.0136.17 varieties due to possible foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related varieties. Not all varieties have been thoroughly tested.

Resistance Management. To limit the potential for development of resistance, **DO NOT** exceed the maximum limits in the table above.

Restrictions

DO NOT use **handheld/portable fogger/mister**. For mechanically pressurized backpack sprayer applications, refer to PPE requirements under **Precautionary Statements**.

The minimum retreatment intervals are listed above with their corresponding diseases.

DO NOT apply more than 5.5 fl ozs of **Merivon** (0.090 lb fluxapyroxad, 0.090 lb pyraclostrobin) per acre per application.

DO NOT apply more than 33 fl ozs of **Merivon** (0.539 lb/acre fluxapyroxad, 0.539 lb/acre pyraclostrobin) per acre per year.

DO NOT make more than 6 applications per year.

DO NOT make more than two (2) sequential applications of **Merivon** before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide.

Mix **Merivon** with water only for applications to **table grapes**.

DO NOT mix with EC formulation pesticides.

DO NOT mix with petroleum oil-based products such as mineral and paraffinic oil (including phytobland oil) or with blended adjuvants containing petroleum oil-based products, e.g., stilet oil, crop oil concentrate and crop oil (emulsifiable).

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Merivon® Xemium® brand fungicide is specially formulated and sold by BASF for the control of pests according to the directions on this label. The purchase price of **Merivon** includes a royalty whereby the purchaser acquires a prepaid license under U.S. Patent No.5,438,070 and (pending No. 2008/0153707 A1) under which purchaser agrees to employ the purchased quantity of **Merivon** only for the above-specified uses under BASF's United States patent rights and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of **Merivon** other than those specified on this label are not licensed through the purchase of this product and the use of this product for other purposes may violate this license and patent rights of BASF.

F500, Merivon, and Xemium are registered trademarks of BASF.

© 2023 BASF Corporation
All rights reserved.

007969-00310.20230714c.NVA 2023-04-0371-0120

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry