



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7504P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

EPA Reg. Number:

Date of Issuance:

7969-322

APR 2 1 201

Term of Issuance: Conditional

Name of Pesticide Product: Stamina F3 Fungicide Seed Treatment

Name and Address of Registrant (include ZIP Code): Charlotte A. Sanson Product Registration Manager

BASF Corporation Manage

26 Davis Drive

Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

Page 1 of 2

Signature of Approving Official:

Date:

APR 2 1 2011

Shaja B. Joyner PM 20

Fungicide Branch, Registration Division (7504P)

EPA Form 8570-6

Notice of Pesticide Registration Stamina F3 RTU Fungicide Seed Treatment EPA Reg. No. 7969-322 Page 2 of 2

- 2. You must submit the following conditional data before the due date of 03/21/2012:
 - a. Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies. These studies must be conducted in commercial containers and the observations should be made at 0, 6, 9, and 12 month intervals.
- 3. Make the following change to the label:
 - a. Change the product registration number to "EPA Reg. No. 7969-322"
- 4. Submit one copy of the revised final printed label before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted" is enclosed for your records.

Shaja B. Joyner, PM 20 Fungicide Branch Registration Division (7504P)

Enclosure:

Label stamped "Accepted" Product Chemistry Review DP Acute Toxicity Review DP



ACCEPTED with COMMENTS In EPA Letter Dated

Under the Federal Insecticide.
Fungicide, and Rodenticide as amended, for the pestic registered under EPA Reg. No.



For commercial and on-farm use

A liquid flowable for control of various seedborne diseases in barley, oats, rye, triticale and wheat

Active Ingredients:

pyraclostrobin*	1.59%
triticonazole**	1.59%
metalaxyl***	0.93%
Other Ingredients:	95.89%
Total:	00.00%

*Contains 0.14 lb of the active ingredient pyraclostrobin per U.S. gallon **Contains 0.14 lb of the active ingredient triticonazole per U.S. gallon

***Contains 0.08 lb of the active ingredient metalaxyl per U.S. gallon

EPA Reg. No. 7969-xxx

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside booklet for complete First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

BASF Corporation 26 Davis Drive, Research Triangle Park, NC 27709



FIRST AID				
 Immediately call a poison control center or doctor for treatment advice. DO NOT induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. DO NOT give anything by mouth to an unconscious person. 				
 Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after first 5 minutes; then continue rinsing. Call a poison control center or doctor for treatment advice. 				
 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. 				
 Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice. 				

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils. If you want more options, follow the instructions for Category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Shoes plus socks
- · Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber > 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water, or to areas where surface water is present, or intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

If treated seed is spilled outdoors or in areas accessible to birds or endangered species, promptly clean up and dispose of properly to prevent ingestion. Dispose of all excess treated seed by burying seed away from bodies of water.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possesion of the user at the time of fungicide application.

DO NOT APPLY this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions, and **Conditions of Sale and Warranty** are to be followed.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and the restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24 hours**.

EXCEPTION: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- · Long-sleeved shirt and long pants
- Socks and shoes
- Chemical-resistant gloves made of any waterproof material

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original container. Keep container closed and in a secure place. Store in an area that is out of reach of children and animals, away from food, feedstuffs, fertilizers and seed. Protect from frost and freezing.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity > 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(continued)

6.

STORAGE AND DISPOSAL (continued)

Container Handling (continued)

Refillable Container. Refill this container with Stamina® F³ RTU fungicide seed treatment only. DO NOT reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Returnable Containers. After use, return cleaned container to point of purchase or designated locations.

DO NOT reuse container for any other purpose. Refill only with Stamina® F³ RTU fungicide seed treatment. For information on the disposal of unused or unwanted product, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

In Case of Spill

in case of large-scale spillage regarding this product, call:

CHEMTREC

1-800-424-9300

• BASF Corporation

1-800-832-HELP (4357)

Product Information

Stamina F³ RTU is a liquid seed treatment used for control or suppression of certain seedborne and soilborne diseases of barley, oats, rye, triticale and wheat. **Stamina F³ RTU** will generally increase emergence and plant stands by controlling or suppressing these diseases.

Seed treatment applications with **Stamina F³ RTU** have been associated with more rapid and increased emergence of seedlings under certain cold conditions.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm.

Stamina F³ RTU contains an EPA-approved dye that imparts an unnatural color to the treated seed to prevent its accidental use as a food or feed for animals. Refer to 21 CFR, Part 2.25.

See the following table for diseases controlled or suppressed by **Stamina F³ RTU**.

When handling Stamina F³ RTU and when treating seed or while augering or handling treated seed, work in a well-ventilated area.

7:0

Stamina® F³ RTU fungicide seed treatment Crop-specific Instructions

Crop	Disease Controlled	Disease Suppressed¹	Rate (fl ozs/100 lbs seed)
Barley	Common root rot Cochliobolus spp.	Fusarium crown rot Fusarium spp.	4.6
	Covered smut Ustilago hordei	Fusarium root rot Fusarium spp.	
	Dry seed decay Penicillium spp.		
	False loose smut Ustilago nigra		
	Fusarium seed rot Fusarium spp.		
	Fusarium seedling blight Fusarium spp.		
	Pythium damping off Pythium spp.		
	Rhizoctonia root rot Rhizoctonia solani		
	True loose smut Ustilago nuda		
Oats	Common root rot Cochliobolus spp.	Fusarium crown rot Fusarium spp.	4.6
	Covered smut Ustilago kolleri	Fusarium root rot Fusarium spp.	
	Fusarium seed rot Fusarium spp.		
	Fusarium seedling blight Fusarium spp.		
	Loose smut Ustilago avenae		
	Pythium damping off Pythium spp.		

(continued)

Stamina® F3 RTU fungicide seed treatment Crop-specific Instructions (continued)

Crop	Disease Controlled	Disease Suppressed¹	Rate (fl ozs/100 lbs seed)
Rye	Common bunt Tilletia caries	Fusarium crown rot Fusarium spp.	4.6
	Common root rot Cochliobolus spp.	Fusarium root rot Fusarium spp.	
	Fusarium seed rot Fusarium spp.		
	Fusarium seedling blight Fusarium spp.		
	Pythium damping off Pythium spp.		
Triticale and Wheat	Common bunt Tilletia caries	Fusarium crown rot Fusarium spp.	4.6
	Common root rot Cochliobolus spp.	Fusarium root rot Fusarium spp.	
	Dry seed decay Penicillium spp.		
	Flag smut Urocystis agropyri		
	Fusarium seed rot Fusarium spp.		
	Fusarium seedling blight Fusarium spp.	·	
	Loose smut Ustilago nuda var. tritici		
	Pythium damping off Pythium spp.		
	Rhizoctonia root rot Rhizoctonia solani		

Suppression is defined as suboptimal control which is still of commercial benefit.

Application Method

Seed should be well conditioned and free of chaff before treating with **Stamina**° **F**° **RTU** fungicide seed **treatment**. **Stamina F**° **RTU** may be applied using standard commercial seed treatment equipment (such as, but not limited to, slurry or mist-type equipment) or on-farm seed treatment equipment. **Stamina F**° **RTU** may also be used in on-the-go type air seeder treatment systems.

Apply **Stamina F³ RTU** to barley, oats, rye, triticale and wheat at the rate of 4.6 fluid ounces of product per 100 pounds of seed. Thorough seed coverage and proper application equipment calibration is essential for good disease control. To achieve thorough coverage, **Stamina F³ RTU** may be diluted with water, if needed, into a slurry. Direct application into a planter box is prohibited.

Use Restrictions

- DO NOT use treated seed for food, feed or oil processing.
- DO NOT store Stamina F3 RTU near feed or foodstuffs.
- DO NOT contaminate feed or foodstuffs.
- DO NOT contaminate domestic or irrigation waters.
- **DO NOT** plant any crop not listed on this label within 30 days after planting **Stamina F³ RTU** treated seed.
- DO NOT use the same auger for treating seed that will handle commercial or feed grains.
- DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed. Store treated seed in cool, dry conditions.
- Stamina F3 RTU is not for use in California.

Seeds treated with this product that are then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This package or bag contains seed that has been treated with Stamina" F' RTU fungicide seed treatment containing pyraclostrobin, triticonazole, and metalaxyl.

DO NOT use treated seed for food, feed, oil processing or any other purpose except planting. Store away from feed and foodstuffs. DO NOT allow children, pets or livestock to have access to treated seeds.

Wear long pants, long-sleeved shirt, shoes and socks, and protective gloves when opening bag or handling treated seed. After the seeds have been planted in soil or other planting media, **DO NOT** enter or allow worker entry into areas with treated seeds for **12 hours**, except that workers may enter the areas with treated seeds without restriction if there will be no worker contact with the soil/media subsurface.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess seed by burying away from bodies of water.

DO NOT contaminate bodies of water when disposing of planting equipment washwater.

Dispose of seed packaging or containers in accordance with local requirements."

10

Conditions of Sale and Warranty

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as the presence of other or untested or unapproved materials, seed of low quality or low vigor or low germination, use of the product in a manner inconsistent with its labeling, misapplication of the product, or weather conditions at planting or environmental conditions during seed storage, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

ST1108

Stamina and the **S** symbol are registered trademarks of BASF.

© 2010 BASF Corporation All rights reserved.

007969-00xxx.20100820.NVA 2010-04-379-0137

BASF Corporation 26 Davis Drive Research Triangle Park, NC 27709

