



**OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION**

WASHINGTON, D.C. 20460

July 23, 2025

Amy Dugger-Webster  
U.S. Regulatory Manager  
BASF Corporation, Agricultural Products  
26 Davis Drive, P.O. Box 13528  
Research Triangle Park, NC 27709-3528

Subject: PRIA Label Amendment – Afidopyropen: Addition of New IR-4 Greenhouse Use  
on Leaf Lettuce  
Product Name: Ventigra® Insecticide  
EPA Registration Number: 7969-393  
Application Date: December 15, 2023  
Petition Number: 4E9104  
Action Case Numbers: 496587 & 496589

Dear Amy Dugger-Webster:

The application referred to above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable under FIFRA section 3(c)(5).

You must submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website

is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Carmen Rodia via e-mail at [Rodia.Carmen@epa.gov](mailto:Rodia.Carmen@epa.gov).

Sincerely,



Digitally signed by Ideliz Negrón-Encarnación  
Date: 2025.07.23 11:20:18 -04'00'

Ideliz Negrón-Encarnación, Ph.D.  
Chief, Invertebrate-Vertebrate Branch 2  
Registration Division (7505T)

Enclosure:      *Stamped "Accepted" Master & Supplemental Labels, July 23, 2025*

{NOTE to PM: This following text is to appear at the top of page 1 of the container booklet and to the first page of the labeling accessed via the QR code.} The label accessed via the QR code is legally valid only for use in registered states and only with a container of **Ventigra<sup>®</sup> insecticide**, EPA Reg. No. 7969-393, that bears a "released for shipment date" on or before {date of website access}. See Page 1 of this document for additional legal requirements on the use of this labeling. For pesticide applications following the label accessed via the QR code, the user must possess a copy of the labeling accessed via the QR Code at the time of application. If the labeling accessed via the QR code conflicts with the labeling found on the pesticide product, use only one set of the labeling instructions in its entirety. **DO NOT** mix and match labeling directions. The labeling accessed via the QR code is the most current version of the **Ventigra**, EPA Reg. No. 7969-393, label approved by US EPA and state lead agencies. Currently the user is responsible for complying with the requirements of product labeling obtained when the user took possession of the product's container. The user must comply with all instructions and requirements in the labeling and must keep the labeling as long as the user has the container. Similarly, the user who relies upon the labeling accessed via the QR code may meet this requirement by maintaining a printed or electronic copy of the labeling to meet federal and state record keeping requirements. In the situations described above where the labeling accessed via the QR code conflicts with labeling on the pesticide container, the user must be able to identify which labeling (i.e., container label or labeling accessed via the QR code) was used and followed at the time of the pesticide's application.

{NOTE to PM: The following text is to appear on each page of the labeling accessed via the QR code.} This web-distributed labeling was rendered by accessing the associated container's QR code on (date). (**Ventigra<sup>®</sup> insecticide**, EPA Reg. No. 7969-393) Web-Distributed Labeling, Page (page #) of (total # of pages).



We create chemistry

**AFIDOPYROPEN    GROUP    9D    INSECTICIDE**

{NOTE to PM: [Text] in brackets denotes optional language.  
{NOTE to PM: {Text} in braces denotes where in the final label text will appear.}


{NOTE to PM: (1) This is a placeholder. The final printed label/container label will bear a unique QR code that when scanned takes the user to the PDF version of the most current label approved by US EPA and state lead agencies where the product is sold. The container label and QR code PDF label both bear EPA required web distributed labeling text per PR-Notice 2014-1. (2) The user must choose a single, valid label version to follow in its entirety. (3) The QR code locations include (based on package type) carton, front panel of booklet, container decal/stay-behind.}



**Ventigra<sup>®</sup>**

**Insecticide**

[Optional text: **ABN: Ventigra<sup>®</sup> Inscalis<sup>®</sup> Insecticide**]



**For control of piercing and sucking insects in ornamental plants, vegetable transplants for the home consumer market, and greenhouse crops grown to production**

**Active Ingredient:**

Afidopyropen, [(3S,4R,4aR,6S,6aS,12R,12aS,12bS)-3-[(cyclopropylcarbonyl)oxy]-1,3,4,4a,5,6,6a,12,12a,12b-decahydro-6,12-dihydroxy-4,6a,12b-trimethyl-11-oxo-9-(3-pyridinyl)-2H,11H-naphtho[2,1-b]pyrano[3,4-e]pyran-4-yl)methyl cyclopropanecarboxylate\* . . . . . 9.78%

**Other Ingredients:** . . . . . 90.22%

**Total:** . . . . . 100.00%

\*Contains 0.83 pound of afidopyropen per gallon, formulated as a dispersible concentrate.

**EPA Reg. No. 7969-393** **EPA Est. No.**

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See full label for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

**In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

**Net Contents:**

BASF Agricultural Solutions US LLC  
2 TW Alexander Drive  
Research Triangle Park, NC 27713

FIRST AID	
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may contact BASF Agricultural Solutions US LLC (hereafter "BASF") at 1-800-832-HELP (4357), twenty-four (24) hours per day seven (7) days per week.</p>	

## Precautionary Statements

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed or inhaled. Avoid contact with skin or clothing. Avoid breathing spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves made of barrier laminate or butyl rubber ≥ 14 mils or nitrile rubber ≥ 14 mils or neoprene rubber ≥ 14 mils or polyvinyl chloride (PVC) ≥ 14 mils or viton ≥ 14 mils, and
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

#### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## Environmental Hazards

### Terrestrial Use

**DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. **DO NOT** pour equipment washwaters or rinsate down into a natural drain or water body.

**DO NOT** exceed the maximum seasonal use rate or the total number of applications of **Ventigra® insecticide** per season. To reduce the potential for developing insecticide-resistance, rotate to an insecticide with a different mode of action.

### Non-target Organisms

**Ventigra** is toxic to aquatic invertebrates. Drift and run-off from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** apply when weather conditions favor drift from treated areas.

### Pollinator Advisory

Although **Ventigra** is not acutely toxic to bees, use at the maximum single application rate may have some short-term behavioral effects on adult bees, but is not expected to have long-term impacts on bees and overall colony health.

### Ground Water Advisory

Afidopyropen and a degradate of concern may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

### Surface Water Advisory

**Ventigra** may impact surface water quality due to run-off of rain water. This is especially true for poorly draining soils and soils with shallow ground water. **Ventigra** is classified as having a medium-high potential for reaching both surface water and aquatic sediment via run-off for several weeks to months after application. A well-maintained

vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs, as required under the **Directions For Use**, will reduce the potential for loading of afidopyropen from run-off and sediment.

Run-off of afidopyropen will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

## Physical or Chemical Hazards

**DO NOT** mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

---

## Directions For Use

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

---

## Use Restrictions

---

- Apply this product only as specified on the EPA approved label and any supplemental labeling.
- **Ventigra® insecticide** is for use by professional applicators.
- **Ventigra** is **NOT** for homeowner use.
- **Ventigra** is **NOT** for sale, distribution, or use in Nassau or Suffolk counties in New York state except by New York-specific supplemental labeling.
- **DO NOT** make more than 2 sequential applications before rotating to a product of a different mode of action group.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- **DO NOT** make **Ventigra** applications at intervals shorter than 7 days.
- **DO NOT** apply to plants exhibiting symptoms of stress or injury, such as stunting, wilting, leaf burn or drop, or abnormal growth.
- A 30-day plant-back interval is appropriate for all food crops not listed on this label.
- For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## Buffer Zones

**Vegetative Buffer Strips.** Construct and maintain a minimum 10-foot vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (e.g., lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuarine/marine habitats). Only apply products containing afidopyropen onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat. For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses*. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21pp. <https://permanent.access.gpo.gov/lps9018/www.wcc.nrcs.usda.gov/water/quality/common/pestmgt/files/newconbuf.pdf>

### Buffer Zone for Ground Application (e.g., ground boom, overhead chemigation, airblast).

- **DO NOT** apply within 10 feet of aquatic habitats (e.g., lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuarine/marine habitats).
- In **California, Florida, and, New York:**
  - **DO NOT** apply within 25 feet of freshwater bodies (e.g., reservoirs, rivers, permanent streams, marshes, natural ponds).
  - **DO NOT** apply within 100 feet of estuarine/marine habitats.

### Buffer Zone for Aerial Application.

- **DO NOT** apply within 150 feet of aquatic habitats (e.g., lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuarine/marine habitats).
- In **California, Florida, and, New York:**
  - **DO NOT** apply within 150 feet of freshwater bodies (e.g., reservoirs, rivers, permanent streams, marshes, natural ponds).
  - **DO NOT** apply within 1,000 feet of estuarine/marine habitats.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of barrier laminate or butyl rubber  $\geq 14$  mils or nitrile rubber  $\geq 14$  mils or neoprene rubber  $\geq 14$  mils or polyvinyl chloride (PVC)  $\geq 14$  mils or viton  $\geq 14$  mils, and
- Shoes plus socks

## NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard of agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses or shadehouses.

**REENTRY: Keep children, pets, and unprotected persons out of treated area until sprays have dried.**

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

### Pesticide Storage

Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals. **DO NOT** store near food or feed.

### Pesticide Disposal

Wastes resulting from using this product must be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

(continued)

## STORAGE AND DISPOSAL (continued)

### Container Handling

**Nonrefillable Container. DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake (capacity  $\leq 5$  gallons) as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Triple rinse containers too large to shake (capacity  $> 5$  gallons) as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

**Triple rinse as follows:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

(continued)



## STORAGE AND DISPOSAL *(continued)*

### Container Handling *(continued)*

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or by other procedures approved by state and local authorities.

### In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF 1-800-832-HELP (4357)

### Steps to take if this material is released or spilled:

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

## Product Information

**Ventigra® insecticide** contains the active ingredient **Inscalis® insecticide**, commonly known as afidopyropen. When used as directed, **Ventigra** controls aphids, mealybugs, and whiteflies, and suppresses certain species of mealybugs and scale insects within the crops and use sites included on this label (refer to the **Application Information** and **Use-specific Application Instructions** sections).

**Ventigra** can be used on ornamental plants, vegetable transplants for the home consumer market, and greenhouse crops grown to production.

**Ventigra** is primarily active through ingestion by the insect and demonstrates translaminar activity in the plant; it is not fully systemic. Thorough coverage of plant surfaces will result in the most effective control. **Ventigra** acts quickly to inhibit insect feeding.

**Ventigra** can be effectively used in Integrated Pest Management (IPM) and resistance management programs for the labeled pests in ornamentals, vegetable transplants for the home consumer market, and greenhouse crops grown to production.

### Mode of Action

The active ingredient in **Ventigra** is classified by the IRAC (Insecticide Resistance Action Committee) as a target-site-of-action **Group 9** insecticide, a chordotonal organ TRPV (Transient Receptor Potential Vanilloid) channel modulator. **Ventigra** disrupts feeding and other behaviors in target insects. Repeated use of insecticides with similar modes of action can lead to the buildup of resistant pest populations.

### Resistance Management

For resistance-management, **Ventigra** contains a **Group 9** insecticide. Any insect population may contain individuals naturally resistant to **Ventigra** and other **Group 9** insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed. To reduce the potential for developing insect resistance, rotate to an insecticide with a different mode of action. Monitor treated pest populations for resistance development. Read product label before applying any insecticide and follow label directions.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Ventigra** or other **Group 9** insecticides within a growing season, or among growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of **Ventigra** per season.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.

- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.

- Adopt an Integrated Pest Management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact BASF representatives at 1-800-832-HELP (4357).

---

## Application Information

---

**Ventigra® insecticide** can be used in ornamental plants, vegetable transplants for the home consumer market, and greenhouse crops grown to production. Refer to the **Use-specific Application Instructions** in **Tables 2, 4, and 5** for specific application information.

### Use Sites

- Commercial and retail nurseries, field and container
- Commercial, public, and recreational gardens
- Commercial and residential landscapes, including golf courses, parks, and recreation areas
- Forest tree and conifer nurseries and plantations
- Greenhouses, lathhouses, and shadehouses
- Interiorscapes

**Ventigra is rainfast one (1) hour after an application has dried.**

**Ventigra** is not a rescue treatment. For **Ventigra** to be most effective, apply at the first sign of insect pests and before the build-up of heavy pressure. Application must be timed to coincide with locally recommended treatment thresholds for label-specific pest listed.

### Water Volume

**Ventigra** is not systemic but has translaminar activity. Thorough and uniform coverage of the leaf surface and/or direct contact of the spray mixture with the target pest is required for control. Make foliar application using properly calibrated sprayers. To provide optimum coverage of ornamental plants with hard-to-wet foliage an adjuvant may be used. Refer to the **Additive and Tank Mixing Information** section of the label.

- BASF recommends 100 gallons/acre spray solution to ensure complete coverage of all leaf surfaces. Minimum spray carrier volume per acre: 20 gallons for ground; 5 gallons for air.

### Aerial Application

For all crops, thorough coverage is required to obtain optimum insect control when aerial applications are employed. Avoid applications under conditions when uniform coverage cannot be obtained. Use no less than the minimum allowed gallons of spray solution per acre as indicated in the **Use-specific Application Instructions** of this label. **DO NOT** apply when conditions favor drift from target area. **DO NOT** apply directly to humans or animals.

### Spray Drift Reduction Management

**DO NOT** apply when wind speed favors drift beyond the area intended for treatment. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

**Importance of Droplet Size.** An important factor influencing drift is droplet size. Small droplets (<150 to 200 microns) drift more than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Use only medium or coarser spray nozzles (for ground and aerial applications) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

**Ground Applications.** Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application. For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy. For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two (2) rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

**Aerial Applications.** The spray boom should be mounted on the aircraft to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter. Flight speed and nozzle orientation must be considered in determining droplet size. Spray must be released at the lowest height consistent with pest control and flight safety. **DO NOT** release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

**Wind Speed Restrictions.** Drift potential increases at wind velocities of less than 3 mph (due to inversion)



potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Only apply this product if the wind direction favors on-target deposition. **DO NOT** apply when wind velocity exceeds 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

#### Restrictions During Temperature Inversions.

**DO NOT** make ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by stable air and increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by mist or ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally near the ground surface in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

**Terrain Precaution.** Local terrain can influence wind patterns. The applicator must be familiar with these patterns and understand their impact on drift.

**Sensitive Areas.** To ensure the protection of threatened or endangered species, it is important to maintain spray drift loadings below levels of concern for any area adjacent to the application site that is not excluded as possible habitat for these organisms.

#### Cleaning Spray Equipment

Before application, start with clean, well-maintained application equipment. Following spray application, thoroughly clean all application equipment. Drain application equipment of any excess product. Thoroughly rinse application equipment and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean equipment near wells, water sources, or desirable vegetation.

---

### Additives and Tank Mixing Information

---

Shake container well before use.

**Ventigra® insecticide** can be tank mixed with most recommended fungicides, insecticides, liquid fertilizers, adjuvants, and additives.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Under certain conditions or when recommended in the **Use-specific Application Instructions**, adjuvants can improve the performance of **Ventigra**. However, all varieties and cultivars have not been tested with all possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced pest control, or crop injury can result from mixing **Ventigra** with other products.

Before using any tank mix (fungicides, insecticides, liquid fertilizers, adjuvants, and additives), test the combination on a small portion of the crop to be treated (including plant cultivars) to ensure that a phytotoxic response will not occur as a result of application. Always follow the most restrictive label use directions.

When an adjuvant is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

---

### Mixing Order

---

Make sure each component is thoroughly mixed and suspended before adding tank mix partners. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Maintain constant agitation during application.

Shake the **Ventigra** container well before use.

1. **Water** - Begin by filling a thoroughly clean sprayer tank 1/2 to 3/4 full with clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-soluble additives** (including dry and liquid fertilizers such as ammonium sulfate or urea ammonium nitrate)
6. **Water-dispersible products** (such as **Ventigra**, dispersible concentrates, dry flowables, wettable powders, other suspension concentrates, or suspo-emulsions).
7. **Water-soluble products**
8. **Emulsifiable concentrates** (such as oil concentrates when applicable)
9. **Water-soluble additives** (such as AMS or UAN when applicable)
10. **Remaining quantity of water**

## Dilution Rate Conversion

Use the following for information on use rates and respective spray volumes of **Ventigra®** insecticide.

**Table 1. Dilution Rate Conversion**

Use rate (fl ozs product/ 100 gallons)	Spray Volume mL of product/ 2 gallons	Spray Volume mL of product/ 5 gallons	Spray Volume mL of product/ 10 gallons	Spray Volume mL of product/ 25 gallons	Spray Volume mL of product/ 50 gallons
1.4	0.8	2	4	10	20
4.8	2.8	7	14	35	70
6	3.6	9	18	45	90
7	4.2	10.5	21	52.5	105

## Use-specific Application Instructions

### Ornamental Plants

**Ventigra** insecticide can be used on ornamental plants such as:

- Annual bedding plants
- Flowering and foliage plants
- Juvenile fruit and nut trees, vines, brambles, and bushberries (Note: Immature and/or inedible fruits, nuts, or berries are not intended for harvest or consumption.)
- Perennials: herbaceous, and woody
- Woody plants and trees

**Table 2. Ventigra® insecticide Use-specific Application Instructions: Ornamental Plants**

Pest	Product Use Rate per Application (fl ozs/100 gallons)
<b>Aphids</b> such as: Cotton/Melon aphid ( <i>Aphis</i> spp.) Green peach aphid ( <i>Myzus</i> spp.) Rose aphid ( <i>Macrosiphum</i> spp.)	1.4
<b>Whiteflies</b> such as: Greenhouse Whitefly ( <i>Trialeurodes</i> spp.) Silverleaf Whitefly ( <i>Bemisia</i> spp.)	4.8 to 7.0
<b>Mealybugs</b> such as: Citrus mealybug ( <i>Planococcus</i> spp.) Long-tailed mealybug ( <i>Pseudococcus</i> spp.) <b>Suppression of:</b> Mexican mealybug ( <i>Phenacoccus</i> gossypii)	4.8 to 7.0
<b>Scale (nymphs and crawlers):</b> <b>Suppression of:</b> Euonymus scale ( <i>Unaspis</i> spp.) Tea scale ( <i>Fiorinia</i> spp.)	4.8 to 7.0

(continued)

**Table 2. Ventigra® insecticide Use-specific Application Instructions: Ornamental Plants** *(continued)*

For <b>Ventigra</b> to be most effective, apply at the first sign of insect pests and before the build-up of heavy pressure. Application must be timed to coincide with locally recommended treatment thresholds for label-specific pest listed.
The addition of a tank-mixed adjuvant enhances control for most species except for aphids. <b>No adjuvant is needed for aphid-only applications.</b> Refer to the <b>Additives and Tank Mixing Information</b> section of this label.
Apply <b>Ventigra</b> at spray volumes sufficient to ensure thorough and uniform coverage for optimal performance.
• <b>Minimum spray carrier volume per acre:</b> 20 gallons for ground; 5 gallons for air.
<b>Use Restrictions for Ornamental Plants</b>
<b>DO NOT</b> make <b>Ventigra</b> applications at intervals shorter than 7 days.
<b>Resistance Management. DO NOT</b> make more than 2 sequential applications of <b>Ventigra</b> before using an effective insecticide with a different mode of action.
<b>DO NOT</b> apply more than 14 fl ozs/A (0.09 lb afidopyropen/A) of <b>Ventigra</b> per crop season. For indoor or greenhouse-grown ornamentals, <b>DO NOT</b> apply more than 17 fl ozs/A (0.11 lb afidopyropen/A) of <b>Ventigra</b> per crop season.
<b>DO NOT</b> apply more than 42 fl ozs/A (0.27 lb afidopyropen/A) of <b>Ventigra</b> per year per crop.

**Plant Phytotoxicity**

**Ventigra** has been applied to a wide variety of common ornamental plants without observed plant injury. Not all species, varieties, and cultivars have been tested for tolerance to **Ventigra**, possible tank mix combinations with **Ventigra**, pesticide treatments preceding or following those with **Ventigra**, and combinations of **Ventigra** with surfactants or adjuvants. Local conditions can also influence plant tolerance and may not match those under which BASF has conducted testing. Because many cultivars within a plant species vary in tolerance to chemical applications and growing conditions, the grower must recognize these differences and test the product accordingly.

At a minimum, always test a small group of representative plants for tolerance to **Ventigra** under local growing conditions and before large-scale use. Grower assumes responsibility for testing species suitability under local growing conditions by treating a small number of plants at the specified rate. At a minimum, this should include evaluating treated plants for several weeks following treatment for possible injury or other effects. To the extent consistent with applicable law, by applying **Ventigra**, the user assumes responsibility for any crop damage or other liability associated with factors beyond the manufacturer’s control, such as weather, presence of other materials, and manner or use of application. Plants with known potential for injury, such as Poinsettia at bract stage, are listed in **Table 3. Sensitive Plant Species**.

**Use caution in applying Ventigra:** Always treat a small area to ensure that plant injury will not occur during large-scale use.

**Table 3. Sensitive Plant Species**

Plant species NOT Tolerant to <b>Ventigra</b>	Coleus	<i>Solenostemum</i> spp. ‘Rustic Orange’
	Ficus	<i>Ficus benjamina</i>
	Poinsettia – at bract formation	<i>Euphorbia pulcherrima</i>
Applications to flowers have occasionally shown discoloration to flowers	Impatiens	<i>Impatiens walleriana</i>
	Petunia	<i>Petunia x hybrida</i> , <i>Petunia</i> spp.

## Vegetable Transplants for the Home Consumer Market

**Ventigra® insecticide** can be used on the following vegetable transplants for the home consumer market only.

**Table 4. Ventigra® insecticide Use-specific Application Instructions: Vegetable Transplants for the Home Consumer Market**

Crop*	Pest	Product Use Rate per Application (fl ozs/100 gallons)
<p>Unless noted with an *, application is not permitted in California.</p> <p><b>Berry, Low Growing (Subgroup 13-07G):</b> Strawberry</p> <p><b>Vegetable, Brassica, Head and Stem (Group 5-16):</b> *Broccoli; Brussels sprouts; *cabbage; cabbage, Chinese (napa); cauliflower; and cultivars, varieties, and/or hybrids of these</p> <p><b>Vegetable, Cucurbit (Group 9):</b> Chayote; Chinese waxgourd (Chinese preserving melon); citron melon; *cucumber; gherkin; gourd, edible (including Chinese okra, cucuzza, hechima, hyotan); <i>Momordica</i> spp. (including balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelon (including true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon); pumpkin; summer squash (including crookneck squash, scallop squash, straightneck squash, vegetable marrow, *zucchini); winter squash (including butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); *watermelon</p> <p><b>Vegetable, Fruiting (Group 8-10):</b> African eggplant; bush tomato; cocona; currant tomato; *eggplant; garden huckleberry; goji berry; ground cherry; martynia; naranjilla; okra; pea eggplant; pepino; *pepper, bell; pepper, non-bell; roselle; scarlet eggplant; sunberry; tomatillo; *tomato; tree tomato; cultivars, varieties and hybrids of these commodities</p> <p><b>Vegetable, Leafy Greens (Subgroup 4-16A):</b> Amaranth, Chinese; amaranth, leafy; arugula; aster, Indian; blackjack; broccoli raab; broccoli, Chinese; cabbage, Abyssinian; cabbage, seakale; Cat's Whiskers; cham-chwi, cham-na-mul; chervil, fresh leaves; Chinese cabbage, bok choy; chipilin; chrysanthemum, garland; cilantro, fresh leaves; collards; corn salad; including lamb's lettuce, and Italian corn salad; cosmos; cress, garden; cress, upland; dandelion; dang-gwi; dillweed, fresh leaves; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; Good King Henry; Hanover salad; huauzontle; jute leaves; kale; *lettuce, bitter, *lettuce, head; *lettuce, leaf (Romaine); maca; mizuna; mustard greens; orach; parsley, fresh leaves; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; radish, leaves; rape greens; rocket, wild; shepherd's purse; *spinach; spinach, Malabar; spinach, New Zealand; spinach, tree; Swiss chard; tanager spinach; *turnip greens; violet, Chinese; watercress; cultivars and/or hybrids of these</p> <p><b>Vegetable, Leaf Petiole (Subgroup 22B):</b> cardoon; celery; celery, Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities</p>	<p><b>Aphids</b> such as: Cotton/Melon aphid (<i>Aphis</i> spp.) Green peach aphid (<i>Myzus</i> spp.) Rose aphid (<i>Macrosiphum</i> spp.) Strawberry aphid (<i>Chaetosiphon</i> spp.)</p>	1.4
	<p><b>Whiteflies</b> such as: Greenhouse Whitefly (<i>Trialeurodes</i> spp.) Silverleaf Whitefly (<i>Bemisia</i> spp.)</p>	4.8 to 7.0
	<p><b>Mealybugs</b> such as: Citrus mealybug (<i>Planococcus</i> spp.) Long-tailed mealybug (<i>Pseudococcus</i> spp.) <b>Suppression of:</b> Mexican mealybug (<i>Phenacoccus gossypii</i>)</p>	4.8 to 7.0
	<p><b>Scale (nymphs and crawlers):</b> <b>Suppression of:</b> Euonymus scale (<i>Unaspis</i> spp.) Tea scale (<i>Fiorinia</i> spp.)</p>	4.8 to 7.0

(continued)

**Table 4. Ventigra® insecticide Use-specific Application Instructions: Vegetable Transplants for the Home Consumer Market** (continued)

For **Ventigra** to be most effective, apply at the first sign of insect pests and before the build-up of heavy pressure. Application must be timed to coincide with locally recommended treatment thresholds for label-specific pest listed.

The addition of a tank-mixed adjuvant enhances control for most species except for aphids. **No adjuvant is needed for aphid-only applications.** Refer to the **Additives and Tank Mixing Information** section of this label.

Apply **Ventigra** at spray volumes sufficient to ensure thorough and uniform coverage for optimal performance.

- **Minimum spray carrier volume per acre:** 20 gallons for ground; 5 gallons for air.

**Use Restrictions for Vegetable Transplants for the Home Consumer Market**

**DO NOT** make **Ventigra** applications at intervals shorter than 7 days.

**Resistance Management. DO NOT** make more than 2 sequential applications of **Ventigra** before using an effective insecticide with a different mode of action.

**DO NOT** apply more than 14 fl ozs/A (0.09 lb afidopyropen/A) of **Ventigra** per crop season.

**DO NOT** apply more than 42 fl ozs/A (0.27 lb afidopyropen/A) of **Ventigra** per year per crop.

### Greenhouse Crops Grown to Production

**Ventigra** can be used on the following greenhouse crops grown to production (edible fruit). Refer to **Table 1. Dilution Rate Conversion** for information on use rates and respective spray volumes.

**Table 5. Ventigra® insecticide Use-specific Application Instructions: Greenhouse Crops Grown to Production**

Crop	Pest	Product Use Rate per Application (fl ozs/100 gallons)	Pre-Harvest Interval (PHI) (days)
<b>Berry, Low Growing (Subgroup 13-07G)</b> Strawberry	<b>Aphids</b> such as: Cotton/Melon aphid ( <i>Aphis</i> spp.) Green peach aphid ( <i>Myzus</i> spp.) Rose aphid ( <i>Macrosiphum</i> spp.) Strawberry aphid ( <i>Chaetosiphon</i> spp.)	1.4	0
<b>Leaf Lettuce (Greenhouse and Shadehouses)</b>			
<b>Vegetable, Cucurbit (Group 9)</b> Cucumber Cultivars, varieties, and/or hybrids of these			
<b>Vegetable, Fruiting (Group 8-10)</b> African eggplant Bush tomato Cocona Currant tomato Eggplant Garden huckleberry Goji berry Groundcherry Martynia Naranjilla Okra Pea eggplant Pepino Pepper, bell Pepper, non-bell Roselle Scarlet eggplant Sunberry Tomatillo Tomato Tree tomato Cultivars, varieties, and/or hybrids of these commodities	<b>Psyllids</b> such as: Potato/tomato psyllid ( <i>Bactericera cockerelli</i> )	7.0	
	<b>Whiteflies</b> such as: Greenhouse Whitefly ( <i>Trialeurodes</i> spp.) Silverleaf Whitefly ( <i>Bemisia</i> spp.) Sweetpotato whitefly ( <i>Bemisia</i> spp.) B and Q biotype		



**Table 5. Ventigra® insecticide Use-specific Application Instructions: Greenhouse Crops Grown to Production** *(continued)*

<p>For <b>Ventigra</b> to be most effective, apply at the first sign of insect pests and before the build-up of heavy pressure. Application must be timed to coincide with locally recommended treatment thresholds for label-specific pest listed.</p> <p>The addition of a tank-mixed adjuvant enhances control for most species except for aphids. <b>No adjuvant is needed for aphid-only applications.</b> Refer to the <b>Additives and Tank Mixing Information</b> section of this label.</p> <p>Apply <b>Ventigra</b> at spray volumes sufficient to ensure thorough and uniform coverage for optimal performance.</p> <ul style="list-style-type: none"><li>• <b>Minimum spray carrier volume per acre:</b> 20 gallons for ground; 5 gallons for air.</li></ul> <p><b>Use Restrictions for Greenhouse Crops Grown to Production</b></p> <p><b>DO NOT</b> make <b>Ventigra</b> applications at intervals shorter than 7 days.</p> <p><b>Resistance Management. DO NOT</b> make more than 2 sequential applications of <b>Ventigra</b> before using an effective insecticide with a different mode of action.</p> <p><b>DO NOT</b> apply more than 14 fl ozs/A (0.09 lb afidopyropen/A) of <b>Ventigra</b> per crop season.</p> <p><b>DO NOT</b> apply more than 42 fl ozs/A (0.27 lb afidopyropen/A) of <b>Ventigra</b> per year per crop.</p>
--

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Agricultural Solutions US LLC ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Inscalix and Ventigra are registered trademarks of BASF.*

© 2025 BASF Agricultural Solutions US LLC  
All rights reserved.

007969-00393.20231212d.**NVA 2023-04-0533-0190**

Supersedes: NVA 2020-04-533-0074

Supplemental: NVA 2023-04-0533-0191

BASF Agricultural Solutions US LLC  
2 TW Alexander Drive  
Research Triangle Park, NC 27713



We create chemistry

13 of 16

## Supplemental Label

# Ventigra<sup>®</sup>

## Insecticide

### For control of piercing and sucking insects in greenhouse leaf lettuce (including in shadehouses) grown to production

This supplemental label expires July 31, 2028, and must not be used or distributed after this date.

#### Active Ingredient:

Afidopyropen, [(3*S*,4*R*,4*aR*,6*S*,6*aS*,12*R*,12*aS*,12*bS*)-3-[(cyclopropylcarbonyl)oxy]-1,3,4,4*a*,5,6,6*a*,12,12*a*,12*b*-decahydro-6,12-dihydroxy-4,6*a*,12*b*-trimethyl-11-oxo-9-(3-pyridinyl)-2*H*,11*H*-naphtho[2,1-*b*]pyrano[3,4-*e*]pyran-4-yl)methyl

cyclopropanecarboxylate ..... 9.78%

**Other Ingredients:** ..... 90.22%

**Total:** ..... 100.00%

Contains 0.83 pound of afidopyropen per gallon, formulated as a dispersible concentrate.

EPA Reg. No. 7969-393

### CAUTION/PRECAUCIÓN

#### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Ventigra<sup>®</sup> insecticide** container label, EPA Reg. No. 7969-393, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Ventigra** before applying.
- Use of **Ventigra** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Ventigra**.

#### Application Information

Refer to the **Ventigra** container label for additional application instructions and restrictions.

#### Use-specific Application Instructions

**Ventigra** can be used on the following greenhouse crops grown to production (edible fruit):

- Leaf lettuce (including in shadehouses)

See the container label for **Table 1. Dilution Rate Conversion** for information on use rates and respective spray volumes of **Ventigra**.

**Table 1. Ventigra® insecticide Use-specific Application Instructions: Greenhouse Crops Grown to Production**

Crop	Pest	Product Use Rate per Application (fl ozs/100 gallons)	Pre-Harvest Interval (PHI) (days)
Leaf lettuce (including in shadehouses)	<b>Aphids</b> such as: Cotton/Melon aphid ( <i>Aphis</i> spp.) Green peach aphid ( <i>Myzus</i> spp.) Rose aphid ( <i>Macrosiphum</i> spp.) Strawberry aphid ( <i>Chaetosiphon</i> spp.)	1.4	0
	<b>Psyllids</b> such as: Potato/tomato psyllid ( <i>Bactericera cockerelli</i> )	7.0	
	<b>Whiteflies</b> such as: Greenhouse Whitefly ( <i>Trialeurodes</i> spp.) Silverleaf Whitefly ( <i>Bemisia</i> spp.) Sweetpotato whitefly ( <i>Bemisia</i> spp.) B and Q biotype		
<p>For <b>Ventigra</b> to be most effective, apply at the first sign of insect pests and before the build-up of heavy pressure. Application must be timed to coincide with locally recommended treatment thresholds for label-specific pest listed.</p> <p>The addition of a tank-mixed adjuvant enhances control for most species except for aphids. <b>No adjuvant is needed for aphid-only applications.</b> Refer to the <b>Additives and Tank Mixing Information</b> section of this label.</p> <p>Apply <b>Ventigra</b> at spray volumes sufficient to ensure thorough and uniform coverage for optimal performance.</p> <ul style="list-style-type: none"><li>• <b>Minimum spray carrier volume per acre:</b> 20 gallons for ground; 5 gallons for air.</li></ul> <p><b>Use Restrictions for Greenhouse Crops Grown to Production</b> <b>DO NOT</b> make <b>Ventigra</b> applications at intervals shorter than 7 days.</p> <p><b>Resistance Management. DO NOT</b> make more than 2 sequential applications of <b>Ventigra</b> before using an effective insecticide with a different mode of action.</p> <p><b>DO NOT</b> apply more than 14 fl ozs/A (0.09 lb afidopyropen/A) of <b>Ventigra</b> per crop season.</p> <p><b>DO NOT</b> apply more than 42 fl ozs/A (0.27 lb afidopyropen/A) of <b>Ventigra</b> per year per crop.</p>			

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Agricultural Solutions US LLC (hereafter "BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Ventiga is a registered trademark of BASF.*

© 2025 BASF Agricultural Solutions US LLC  
All rights reserved.

007969-00393.20231212d.NVA 2023-04-0533-0191

BASF Agricultural Solutions US LLC  
2 TW Alexander Drive  
Research Triangle Park, NC 27713



We create chemistry