



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION
WASHINGTON, D.C. 20460

January 19, 2026

Casey Pehrson
Sr. Regulatory Affairs Manager (US)
BASF Agricultural Solutions US LLC
2 TW Alexander Drive
Research Triangle Park, NC 27713

Subject: Approval of Label Amendment; Only Indicated Changes Reviewed: addition of diseases (pests) for soybean and corn; minor administrative updates
Product Name: Fluopyram ST
EPA Registration Number: 7969-441
Application Date: October 21, 2025
Case Number: 672601

Dear Casey Pehrson:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. However, EPA reviewed only the label changes highlighted, marked, or otherwise indicated on the submitted label. Any other changes to the previously approved label that were not clearly highlighted, marked, or otherwise indicated in your submission were not reviewed and may form the basis of regulatory and/or enforcement action if later discovered by the Agency. Further, submission of a label amendment application with unidentified changes may be considered a knowing submission of false information to the Agency. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The label submitted with the application has been stamped "Accepted Only Indicated Revisions Reviewed" and is enclosed for your records.

This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 C.F.R. § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 C.F.R. § 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website contains any false or misleading statement, design, or graphic, the product may be misbranded and unlawful to sell or distribute under FIFRA Sections 2(q)(1)(A) and 12(a)(1)(E). 40 C.F.R. § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on the product label, claims made as part of the product's sale or distribution may not substantially differ from those claims approved through the registration process under FIFRA Section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the product will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Julie Kelleher at Kelleher.Julie@epa.gov.

Sincerely,

Kable Bo Davis

Kable Bo Davis, Senior Advisor
Office of Pesticide Programs
Registration Division; Immediate Office

Enclosure

Fluopyram ST

Alternate Brand Names: ILEVO®; Copeo®; Copeo® PRIME; ILEVO® seed treatment; Copeo® seed treatment; Copeo® Prime seed treatment; Aramax™ XC seed treatment; VERCORAS™ XC seed treatment

A systemic seed treatment for use on corn, cotton (subgroup 20C), peanut, sorghum, soybean, and wheat for the protection against damage caused by early season plant pathogenic nematodes, and rapeseed (subgroup 20A) for the protection from soilborne Blackleg (*Leptosphaeria maculans*). As a soybean seed treatment provides protection from seedling infections by *Fusarium virguliforme*, the causal agent of Sudden Death Syndrome, and suppression of Red Crown Rot (*Calonectria illicicola*). As a peanut seed treatment provides protection from seedborne fungi. As a corn seed treatment for the suppression of *Bipolaris maydis*, the causal agent of Southern corn leaf blight.

Active Ingredient:

fluopyram: *N*-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-(trifluoromethyl)benzamide* 49.02%

Other Ingredients: 50.98%

Total: 100.00%

Contains 5 lbs fluopyram per gallon (600 g fluopyram per liter)

*(CAS Number 658066-35-4)

EPA Reg. No. 7969-441

SUSPENSION CONCENTRATE

EPA Est. No.

ACCEPTED
ONLY INDICATED
REVISIONS REVIEWED

01/19/2026

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

7969-441

No label revisions other than those indicated were reported to the Agency.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional **Precautionary Statements** and **Directions For Use** on label.

In case of an emergency call 24 hours a day 1-800-832-HELP (4357).

Net Contents:

BASF Agricultural Solutions US LLC
2 TW Alexander Drive
Research Triangle Park, NC 27713

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • DO NOT give anything by mouth to an unconscious person.
<p>For MEDICAL Emergencies call 24 hours a day 1-800-832-HELP (4357). In case of emergency call toll free BASF Agricultural Solutions US LLC (hereafter "BASF") 1-800-832-HELP (4357). Have a product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p>NOTE TO PHYSICIAN: Treat symptomatically.</p>	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE)

Workers involved with treating the seed

(e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) and others exposed to the concentrate, and cleaners/repairers of seed treatment equipment must wear:

- Long-sleeve shirt and long pants
- Shoes plus socks
- Waterproof gloves (such as natural rubber, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyethylene, polyvinyl chloride, or viton)

Baggers and bag sewers must wear:

- Long-sleeve shirt and long pants
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
Users should:
<ul style="list-style-type: none"> • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluopyram. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading.

Directions For Use

STOP **READ THE ENTIRE LABEL BEFORE USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read the entire label and observe all label directions and precautions before use.

STORAGE AND DISPOSAL	STORAGE AND DISPOSAL (continued)
<p>DO NOT contaminate water, food or feed by storage, disposal or cleaning of equipment.</p>	
<p>Pesticide Storage</p>	<p>Container Handling (continued)</p>
<p>Store in original container away from feed and food. Store in cool, dry area. DO NOT store in direct sunlight.</p>	<p>Dilutable Seed Treatment Products in Nonrefillable Plastic Containers (continued)</p>
<p>DO NOT allow prolonged storage in temperatures that exceed 105° F (40° C) or in temperatures that fall below 14° F (-10° C).</p>	<p>Rigid nonrefillable containers</p>
<p>Pesticide Disposal</p>	<p>that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)</p>
<p>To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).</p>	<p>Nonrefillable Container. DO NOT reuse or refill this container. After emptying product from container, either return container to BASF, or rinse and either recycle or dispose of the container as follows:</p>
<p>Container Handling</p>	<p>Bottom discharge IBC (e.g. Schuetz Caged IBC or Snyder Square Stackable)</p>
<p>Dilutable Seed Treatment Products in Nonrefillable Plastic Containers</p> <p>Rigid nonrefillable containers (equal to or less than 5 gallons)</p>	<p>Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.</p>
<p>Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.</p>	<p>Top discharge IBC, drums, kegs (e.g. Snyder 120 Next Gen, Bonar B120, drums, and kegs)</p>
<p>Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.</p>	<p>Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.</p>
<p>Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.</p>	<p>Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.</p>

(continued)

(continued)

STORAGE AND DISPOSAL (continued)

Container Handling (continued)

Dilutable Seed Treatment Products in Nonrefillable Metal Containers

Nonrefillable Container. DO NOT reuse or refill this container. After emptying product from container, either return container to BASF, or rinse and either recycle or dispose of the container as follows:

Liquid dilutables in containers small enough to shake (5 gallons or less)

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Solid dilutables in containers small enough to shake (5 gallons or 50 pounds or less)

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Any dilutable pesticides in containers too large to shake (larger than 5 gallons or 50 pounds)

Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

(continued)

STORAGE AND DISPOSAL (continued)

Container Handling (continued)

Nondilutable Seed Treatment Products in Nonrefillable Containers

Seed treatment products in nonrefillable fiber drums with liners

Nonrefillable Container. DO NOT reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Seed treatment products in nonrigid, nonrefillable containers

Nonrefillable Container. DO NOT reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Seed Treatment Products in Refillable Containers

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. After emptying product from container, either return container to BASF, or rinse and either recycle or dispose of the container as follows:

Bottom discharge IBC (e.g. Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

(continued)

STORAGE AND DISPOSAL (continued)

Container Handling (continued)

Seed Treatment Products in

Refillable Containers (continued)

Top discharge IBC, drums, kegs (e.g. Snyder 120

Next Gen, Bonar B120, drums, and kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container.

Product Information

- **Fluopyram ST** is a seed treatment fungicide labeled for the control or suppression of certain plant pathogenic fungi and nematodes.
- **Fluopyram ST** is a seed treatment fungicide that when applied to seed, aids in protecting the seed and seedling against certain early season plant pathogenic fungi and nematodes that attack the root system.
- The use of **Fluopyram ST** can protect against early season infection by *Fusarium virguliforme*, the causal agent of Sudden Death Syndrome.
- In areas of high nematode infestation additional control measures may be warranted.

Use Restrictions

- For use only in commercial seed treatment equipment.
- Not for use on agricultural establishments in hopper box, planter box, slurry box, or other on-farm seed treatment applications at or immediately before planting.
- **DO NOT** use this product except as directed on this label or on other BASF supplemental labeling for this product.
- **DO NOT** apply more than 0.45 lb ai/A (11.52 fl ozs product) of fluopyram per year (including seed treatment, foliar, in-furrow or banded applications).

- Refer to the specific use directions and restrictions in each crop, crop group or crop subgroup table.
- This product does not contain colorant. The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

Labeled Uses

corn, cotton (subgroup 20c), peanut, rapeseed (subgroup 20A), sorghum, soybean, wheat

Fungicide Resistance Management (FRAC)

For resistance management, **Fluopyram ST** contains a **Group 7** fungicide. Any fungal population may contain individuals naturally resistant to **Fluopyram ST** and other **Group 7** fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Fluopyram ST** or other **Group 7** fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your BASF representative, pesticide distributor, or university extension specialist.

Application Instructions

Use in commercial seed treatment facilities with commercial or professional seed treatment application equipment only.

- Use good quality, well-conditioned seed free of chaff.
- Thorough and uniform seed coverage is required for optimum performance.
- Apply at specified rates with either seed treatment equipment that has “direct inject” or dedicated pump capabilities or in a water-based mixture using standard slurry seed treatment application equipment. Consult a seed treatment specialist about crop-specific slurry rates and seed treatment calibration and operation procedures. If applying in a water-based mixture, mix only as much product as is needed for immediate application to seed.

Compatibility Testing and Tank Mix Partners

- **Fluopyram ST** can be used in combination with other registered BASF seed treatment fungicides and insecticides. Pretest for compatibility with other seed treatment products.
- It is the pesticide user’s responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility

It is essential that before using **Fluopyram ST** seed treatment in any tank mixture, the compatibility of the mixture be established. Add **Fluopyram ST** at the labeled rate to a clean quart jar containing approximately one-half the amount of water intended for a final slurry application rate. Next, follow with all other tank mix components that will be used in the total slurry application. Add last the remaining balance of water. The total amount of volume is determined by the seed size and how much is necessary to ensure complete and uniform coverage and distribution on the seed, as well as the type of commercial seed treating application equipment that will be used.

DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GEL. USE TANK MIXTURES IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

Mixing Instructions

When applied with seed treatment application equipment designed to treat seed utilizing a treatment slurry mixture:

- Add approximately 1/2 to 2/3 of the required water to the treatment slurry tank followed by the accompanying seed treatment products and **Fluopyram ST**.
- The order of product addition is dependent on the accompanying seed treatment products and application equipment used.
- Allow each slurry component to disperse completely prior to the next addition until a uniform suspension is obtained.
- After all slurry components have been added and dispersed completely, add the remaining 1/2 to 1/3 of required water for proper slurry volume and mix until uniform.
- **DO NOT** store mixed slurries for greater than 72 hours.
- Maintain adequate agitation until treatment slurry is applied.
- For additional instructions for use, slurry component mixing order, and other application suggestions contact your BASF representative.

Specific Crop Use Directions

Corn

Pests Controlled - Soilborne nematodes: *Belonolaimus* spp., *Criconemella* spp., *Meloidogyne incognita*, *Paratrichodorus* spp., *Pratylenchus* spp., suppression of Southern Corn Leaf Blight (*Bipolaris maydis*).

Rate of Fluopyram ST				
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/80,000 seeds (fl ozs/80,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
3,706 (1,681)	15.05 to 154.82 (0.23 to 2.37)	3.25 to 33.42 (0.11 to 1.13)	0.025 to 0.25	1.01 to 10.06 (0.0022 to 0.022)

¹ Rate is calculated based on number of seeds/kg averages of corn (3,706).

² Based on seeding rate of 40,250 seeds/acre [(40,250 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

Fluopyram ST provides early-season suppression of Southern Corn Leaf Blight (*Bipolaris maydis*) at rates between 0.125 to 0.25 mg ai/seed. When used as part of an IPM system that includes tolerant hybrids and foliar fungicide application, **Fluopyram ST** provides additional late season control of Southern Corn Leaf Blight caused by *Bipolaris maydis*.

Crop-specific Restrictions

DO NOT exceed 2.37 fl ozs/100 lbs seed (0.25 mg ai/seed).

DO NOT exceed 0.45 lb ai/A (11.52 fl ozs product) per year, including seed treatment, foliar, in-furrow or banded applications.

Seeds must not be broadcast planted.

Cotton (subgroup 20C)

Pests Controlled - Soilborne nematodes: *Hoplolaimus* spp., *Meloidogyne incognita*, *Rotylenchulus reniformis*.

Rate of Fluopyram ST				
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/100,000 seeds (fl ozs/100,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
9,921 (4,500)	289.4 to 578.7 (4.45 to 8.87)	29.17 to 58.33 (0.99 to 1.97)	0.175 to 0.35	14.88 to 29.75 (0.033 to 0.066)

¹ Rate is calculated based on number of seeds/kg averages of cotton (9,921).

² Based on seeding rate of 85,000 seeds/acre [(85,000 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

Crop-specific Restrictions

DO NOT exceed 8.87 fl ozs/100 lbs seed (0.35 mg ai/seed).

DO NOT exceed 0.45 lb ai/A (11.52 fl ozs product) per year, including seed treatment, foliar, in-furrow or banded applications.

Seeds must not be broadcast planted.

Peanut

Diseases Controlled - Seedborne Fungi: *Rhizopus* spp., *Fusarium* spp., *Aspergillus* spp., *Penicillium* spp.

Pests Controlled - Soilborne nematodes: *Meloidogyne* spp., *Pratylenchus* spp., *Belonolaimus longicaudatus*, *Criconemoides ornatus*

Rate of Fluopyram ST				
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/100,000 seeds (fl ozs/100,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
1,521 (690)	83.6 to 207 (1.28 to 3.19)	55 to 136 (1.86 to 4.62)	0.33 to 0.82	23.9 to 59.5 (0.05 to 0.13)

¹ Rate is calculated based on number of seeds/kg averages of peanut (1,521).

² Based on seeding rate of 72,600 seeds/acre [(72,600 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

Crop-specific Restrictions

DO NOT exceed 3.19 fl ozs/100 lbs seed (0.82 mg ai/seed).

DO NOT exceed 0.45 lb ai/A (11.52 fl ozs product) per year, including seed treatment, foliar, in-furrow or banded applications.

Seeds must not be broadcast planted.

Rapeseed (subgroup 20A)

Pest Controlled - Residue or soilborne Blackleg (*Leptosphaeria maculans*).

Rate of Fluopyram ST			
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	mg ai/seed ¹	g/A ² (lb/A)
200,000 (90,718)	125 to 250 (1.92 to 3.83)	0.00375 to 0.0075	1.59 to 3.19 (0.0035 to 0.007)

¹ Rate is calculated based on number of seeds/kg averages of canola (200,000 seeds per kg).

² Based on seeding rate of 425,000 seeds/acre [(425,000 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

Crop-specific Directions

Fluopyram ST will reduce blackleg infection on the cotyledons and early leaves and will reduce lodging and the degree of stem infection later in the season.

Use the higher rate in situations conducive to greater risk of blackleg disease.

Crop-specific Restrictions

DO NOT exceed 3.83 fl ozs/100 lbs seed (0.0075 mg ai/seed).

DO NOT exceed 0.45 lb ai/A (11.52 fl ozs product) per year, including seed treatment, foliar, in-furrow or banded applications.

Seeds must not be broadcast planted.

Sorghum

Pests Controlled - Soilborne nematodes: *Belonolaimus* spp., *Criconemella* spp., *Meloidogyne incognita*, *Paratrichodorus* spp., *Pratylenchus* spp.

Rate of Fluopyram ST				
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/100,000 seeds (fl ozs/100,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
39,683 (18,000)	165.48 to 330.05 (2.52 to 5.04)	4.17 to 8.34 (0.14 to 0.28)	0.025 to 0.05	2.5 to 5 (0.005 to 0.01)

¹ Rate is calculated based on number of seeds/kg averages of sorghum (39,683).

² Based on seeding rate of 100,000 seeds/acre [(100,000 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

Crop-specific Restrictions

DO NOT exceed 5.04 fl ozs/100 lbs seed (0.05 mg ai/seed).

DO NOT exceed 0.45 lb ai/A (11.52 fl ozs product) per year, including seed treatment, foliar, in-furrow or banded applications.

Soybean

Pests Controlled - Sudden Death Syndrome (SDS) caused by *Fusarium virguliforme*. Early season Septoria brown spot (*Septoria glycines*). Soilborne nematodes: *Heterodera glycines*, *Hoplolaimus* spp., *Meloidogyne incognita*, *Pratylenchus* spp., *Rotylenchulus reniformis*. Suppression of Red Crown Rot (*Calonectria ilicicola*).

Rate of Fluopyram ST				
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/140,000 seeds (fl ozs/140,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
5,662 (2,550)	141.55 to 235.92 (2.17 to 3.62)	35 to 58 (1.18 to 1.97)	0.15 to 0.25	37.5 to 62.5 (0.08 to 0.14)

Pests Controlled - Soilborne nematodes: *Heterodera glycines*, *Hoplolaimus* spp., *Meloidogyne incognita*, *Pratylenchus* spp., *Rotylenchulus reniformis*.

Rate of Fluopyram ST				
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/140,000 seeds (fl ozs/140,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
5,662 (2,550)	70.78 to 235.92 (1.08 to 3.62)	18 to 58 (0.6 to 1.97)	0.075 to 0.25	18.75 to 62.5 (0.04 to 0.14)

¹ Rate is calculated based on number of seeds/kg averages of soybean (5,662).

² Based on seeding rate of 250,000 seeds/acre [(250,000 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

Crop-specific Directions

Use the higher rate for suppression of red crown rot.

Crop-specific Restrictions

DO NOT exceed 3.62 fl ozs/100 lbs seed (0.25 mg ai/seed).

DO NOT exceed 0.45 lb ai/A (11.52 fl ozs product) per year, including seed treatment, foliar, in-furrow or banded applications.

DO NOT graze or feed soybean forage and hay to livestock after treating with **Fluopyram ST**.

Seeds must not be broadcast planted.

Wheat

Pests Controlled - Soilborne nematodes: *Belonolaimus* spp., *Criconemella* spp., *Meloidogyne incognita*, *Paratrichodorus* spp., *Pratylenchus* spp.

Rate of Fluopyram ST				
# seeds/kg (# seeds/lb)	ml/100 kg seed (fl ozs/100 lbs seed)	ml/100,000 seeds (fl ozs/100,000 seeds)	mg ai/seed ¹	g/A ² (lb/A)
31,967 (14,500)	133.3 to 266.6 (2.03 to 4.06)	4.17 to 8.34 (0.14 to 0.28)	0.025 to 0.05	58.5 to 117 (0.13 to 0.26)

¹ Rate is calculated based on number of seeds/kg averages of wheat (31,967).

² Based on seeding rate of 2,340,000 seeds/acre [(2,340,000 seeds/A) x (mg ai/seed) x (g/1,000 mg)]

Crop-specific Restrictions

DO NOT exceed 4.06 fl ozs/100 lbs seed (0.05 mg ai/seed).

DO NOT exceed 0.45 lb ai/A (11.52 fl ozs product) per year, including seed treatment, foliar, in-furrow or banded applications.

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with **Fluopyram ST** must be labeled with the following statements:

- This seed has been treated with:
 - **Fluopyram ST**, which contains **fluopyram**.
- **DO NOT** use treated seed for food, feed, or oil purposes.

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

- Store treated seeds away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment washwaters.
- Treated seed must be adequately covered with soil at planting.
- Dispose of seed packaging or containers in accordance with local requirements. **DO NOT** use empty seed bags for any other purpose.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's specifications. Surplus seed or empty seed containers should be stored or disposed according to local federal regulations.
- Treat at label specified rate.

SEED BAG TAG (continued)

• **Restrictions:**

- **DO NOT** graze or feed soybean forage and hay to livestock after treating with **Fluopyram ST**.

• **Rotational Crops Plant-back Restrictions:**

- The following crops can be replanted immediately following the last application of fluopyram:
artichoke (globe); Brassica (cole) leafy vegetables (group 5); bulb vegetables (subgroup 3-07A and 3-07B); carrot; cereals (group 15) [which include: barley, buckwheat, field corn, popcorn, sweet corn, millet (pearl and proso), oats, rice; rye, sorghum, teosinte, triticale, and wheat]; citrus (group 10-10); cotton (subgroup 20C); cucurbits (group 9); dill seed; fruiting vegetables (group 8-10); ginseng; grapes and small vines (except fuzzy kiwifruit) (subgroup 13-07F); herb (subgroup 19A); hop; leafy vegetables (except watercress) (group 4); legume vegetables (except soybean, cowpea and dried peas); peanut; pome fruit (group 11-10); potato and other root, tuberous and corm vegetables (except sugar beet) (subgroup 1B and 1C); rapeseed (subgroup 20A); small berries (caneberries and bushberries) (subgroup 13-07A and 13-07B); soybean; stone fruits (group 12-12); strawberry and other low-growing berries, except cranberry (subgroup 13-07G); sugar beet; sunflower (subgroup 20B); tobacco; tree nut (group 14-12).
- Alfalfa can be planted after 14 days and sugarcane can be planted after 14 days in Region 3.
- **DO NOT** rotate to crops other than those listed above.

(continued)

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as the presence of other or untested or unapproved materials, seed of low quality or low vigor or low germination, use of the product in a manner inconsistent with its labeling, misapplication of the product, or weather conditions at planting or environmental conditions during seed storage, all of which are beyond the control of BASF Agricultural Solutions US LLC ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

ST1108

Fluopyram ST is specially formulated and sold by BASF for the control of various pathogens according to the directions on this label. The purchase price of **Fluopyram ST** includes a prepaid license under which purchaser agrees to employ the purchased quantity of **Fluopyram ST** only for the above-specified uses and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of **Fluopyram ST** other than those specified on this label are not licensed through the purchase of this product.

Aramax and **VERCORAS** are trademarks of BASF.

Copeo and **ILEVO** are registered trademarks of BASF.

© 2025 BASF Agricultural Solutions US LLC
All rights reserved.

007969-00441.20250926.**NVA 2025-04-0597-0243**
Supersedes: NVA 2020-04-597-0164

BASF Agricultural Solutions US LLC
2 TW Alexander Drive
Research Triangle Park, NC 27713

BASF
We create chemistry