



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

7969-506

Date of Issuance:

12/6/2024

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Poncho 750 Seed Treatment

Name and Address of Registrant (include ZIP Code):

Matthew Feinberg
Product Registration Manager
BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Venus Eagle, Product Manager 01
Invertebrate and Vertebrate Branch 3, Registration Division

Date:

12/6/24

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 7969-506."
3. Submit one copy of the final printed label for the record before you release the product for shipment.
4. If, after formal consultation with the appropriate Service(s), additional modifications are identified in the Service's Biological Opinion(s) for clothianidin, EPA will notify BASF in writing consistent with the terms in the Biological Opinion of any necessary required changes. BASF must submit an application for amendment incorporating any required changes, including amended labels, consistent with the timeline specified in EPA's notification. If BASF fails to comply with this term, BASF has agreed in prior written acceptance on **November 26, 2024** of these terms that EPA may cancel the registration under an expedited process under FIFRA 6(e).

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 1/30/2024

The following alternate brand names have been added to the product record:

- Modalex 750 Seed Treatment

If you have any questions, please contact Jasmin Jackson at jackson.jasmin@epa.gov.

Enclosure: Final Approved Label



We create chemistry

Poncho[®] 750

Seed Treatment

Alternate Brand Name: Modalex[™] 750 seed treatment

ACCEPTED

12/6/2024

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 7969-506

Seed treatment use on broccoli, canola, carrot, cereal grains (except corn, rice, sorghum and wild rice), corn, cotton, leafy greens, leek, onion (bulb and bunching), potato (seed piece treatment), rapeseed, rice, sorghum, soybean, and sugar beet for the control of listed insect pests.

All seed treated with this product must be conspicuously colored at the time of treatment.

Active Ingredient:

clothianidin 57.25%

Other Ingredients: 42.75%

Total: 100.00%

Contains 6.26 lbs clothianidin per U.S. gallon

EPA Reg. No. 7969-XXX

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See [inside booklet] [back panel] for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions. [Note to reviewer: Location of additional precautionary statements and directions for use will vary between those listed, depending on container type/size.]

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • DO NOT induce vomiting unless told to by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • DO NOT give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).	
NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeve shirt
- Long pants
- Chemical-resistant gloves made out of: barrier laminate, nitrile rubber \geq 14 mils, butyl rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to birds, mammals and aquatic invertebrates.

Treated seeds exposed on soil surface may be hazardous to birds and mammals. Promptly cover or collect treated seed spilled during loading or planting.

This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust, which can drift to blooming crops or weeds.

Surface Water Label Advisory

DO NOT apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwater or rinseate. **DO NOT** apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on the label.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spill or equipment leak and dispose of wastes according to the disposal instructions on this label.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using this product.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.

Excess treated seed may be used for ethanol production only if (1) By-products are not used for livestock feed and (2) No measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeve shirt
- Long pants
- Chemical-resistant gloves made out of: barrier laminate, nitrile rubber \geq 14 mils, butyl rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils or viton \geq 14 mils
- Shoes plus socks

In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to take if material is released or spilled:

- Wear **Personal Protective Equipment (PPE)** and avoid exposure when managing a spill. (See **Precautionary Statements** section of this label for required PPE.)
- Dike and contain the spill with inert, absorbent material (e.g., sand, earth) and transfer liquid and solid diking material to separate containers for disposal. Small-scale spills of this product (that can be cleaned up with a typical spill kit) may be applied to labeled sites.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep spill out of all sewers and open bodies of water.

Product Information

Poncho® 750 seed treatment:

- Systemic insecticide seed treatment for use on broccoli, canola, rapeseed, carrot, cereal grains (except corn, rice, sorghum and wild rice), corn, cotton, leafy greens,

leek, onion (bulb and bunching), potato (seed piece treatment), rice, sorghum, soybean, and sugar beet for the control of listed insects.

- Provides early season protection against the listed insect pests.
- Provides protection under low-to-moderate corn root-worm pressure.

Application Instructions

Poncho 750:

- To be used in liquid or slurry treaters. Mix thoroughly before use or use entire container at one time. Pretest all tank mixes to determine physical compatibility between formulations. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture.
- Treatment of highly mechanically damaged seed, or seed of known low vigor or poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of BASF, no claims are made to guarantee germination of carryover seed.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's specifications. Surplus seed or empty seed containers must be stored or disposed according to local federal regulations.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inches or greater to minimize exposure.

Use Restrictions

- All seed treated with this product must be conspicuously colored at the time of treatment. This product contains no colorant. An EPA-approved dye or colorant must be added when this product is applied to seed to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals. Regulations pertaining to coloration of treated seed enforced by 40 CFR 153.155 must be strictly adhered to when using this product.
- For use only in commercial seed treatment facilities. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications except for cereal grains (except corn, rice, sorghum, and wild rice) and potato seed pieces.
- Regardless of method of application (seed treatment, soil, or foliar), **DO NOT** apply more than 0.2 lb of active ingredient clothianidin per acre per year.
- Follow-up foliar sprays of a registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.
- **DO NOT** broadcast plant seeds.

Refer to the specific use directions and restrictions in each crop table.

Detailed Use Directions

Commercial Seed Treatment Facilities

Broccoli

Insect Pests	Product Rate fl oz/1,000 seeds (mg ai/seed)
Aphids Bagrada bug Cabbage maggot Flea beetle	0.054 (1.1925)

Crop-specific Restrictions

- **Poncho® 750 seed treatment** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Canola; Rapeseed

(cultivars, varieties and/or hybrids, including *Brassica napus*, *Brassica rapa*, and *Brassica juncea*)

Insect Pests	Product Rate fl ozs/100 lbs seed (mg ai/seed) ¹
Flea beetle Wireworms	3.07* (0.00664) 4.10** (0.008863) 8.18*** (0.0177)

* Use this rate under **low-to-moderate** flea beetle pressure.

** Use this rate under **moderate** flea beetle pressure.

*** Use this rate under **high-to-extreme** flea beetle pressure where extended control is required.

¹ Rate is calculated based on number of seeds/lb averages of canola (102,500).

Crop-specific Restrictions

- Rapeseed greens and seed grown or harvested from **Poncho 750**-treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from **Poncho 750**-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.

Carrot

Insect Pest	Product Rate fl oz/1,000 seeds (mg ai/seed)
Carrot rust fly	0.0015 to 0.003 (0.0338 to 0.0675)

Crop-specific Restrictions

- **Poncho® 750 seed treatment** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Cereal Grains (except corn, rice, sorghum, wild rice) barley, buckwheat, millet (pearl, proso), oats, rye, teosinte, triticale, wheat

Insect Pests	Product Rate fl ozs/100 lbs seed (mg ai/seed) ¹
Wireworms	0.2 to 1.43 (0.00324 to 0.02318)
Aphids (including bird cherry-oat, English grain, greenbug, Russian wheat aphid)	0.6 to 1.43 (0.00971 to 0.02318)
Grasshoppers ² Hessian fly	1.43 (0.02318)

¹ Rate is calculated based on number of seeds/lb averages of wheat (13,700).

² Not registered for use in California.

Crop-specific Directions

- **Wireworm seed protection** - Use 0.2 fl oz/100 lbs seed rate or higher, depending on field history and site populations.
- **Hessian fly protection to wheat seedlings in the Fall season** - Use 1.43 fl ozs/100 lbs seed.
- For maximum effectiveness, treat seed uniformly. Use the higher rate to provide increased length of protection and from heavy insect pressure and to reduce potential spread of barley yellow dwarf virus due to aphid vectors.
- Grasshopper feeding damage to seedling plants under low-to-moderate insect pressure will be reduced when **Poncho 750** is used. Treated seed may be planted as a 50 to 60 foot border around the edges of the crop field as a barrier to grasshoppers advancing into the interior of the field.

Crop-specific Restrictions

- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Corn (field, popcorn)

Insect Pests	Product Rate fl ozs/80,000 seeds (mg ai/seed)
Corn flea beetle (through first true seed leaf) Seedcorn maggot ² Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) White grub ² (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae)	0.45 (0.125)
Brown stinkbug Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) Sugarcane beetle Thrips White grub (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae) Wireworms	0.9 to 1.8 (0.25 to 0.5)
Corn leafhopper Corn rootworm (including Mexican, northern, southern, western) Southern corn billbug	4.51 (1.25)

¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed.

² For seed protection only.

Crop-specific Restrictions

- **Poncho® 750 seed treatment** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Corn (sweet)

Insect Pests	Product Rate fl ozs/80,000 seeds (mg ai/seed)
Corn flea beetle (through first true seed leaf) Seedcorn maggot ² Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) White grub ² (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae)	0.45 (0.125)
Brown stinkbug Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle) Sugarcane beetle Thrips White grub (including European, Chafer larvae, Japanese beetle larvae, May/June beetle larvae) Wireworms	0.9 to 1.8 (0.25 to 0.5)

¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed.

² For seed protection only.

Crop-specific Restrictions

- **Poncho® 750 seed treatment** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Cotton (delinted only)

Insect Pests	Product Rate fl ozs/100,000 seeds (mg ai/seed)
Aphids Cutworm, black Early season thrips Storage pests ¹ (including Indian meal moth, lesser grain borer, red flour beetle, rice weevil, rusty grain beetle)	1.59 (0.353)

¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed.

Crop-specific Restrictions

- Not registered for use in California.
- **Poncho® 750 seed treatment** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Leafy Greens (subgroup 4A)

Insect Pests	Short Season Crops	Short Season Product Rate fl ozs/100 lbs seed ¹ (0.025 to 0.05 mg ai/seed)	Seed Size seeds/lb ¹
Aphids Leafminers (Suppression)	amaranth chrysanthemum (edible-leaved, garland)	45.1 to 90.2	400,000
	arugula	29.3 to 58.6	260,000
	chervil	18.3 to 36.6	162,532
	corn salad	19.4 to 38.9	172,400
	cress (garden, upland)	21.7 to 43.4	192,320
	dandelion	3.9 to 7.9	35,000
	dock	36.6 to 73.3	325,000
	endive	48.1 to 96.1	426,400
	lettuce (lettuce, spring mix)	52.9 to 105.9	469,933
	orach	15.0 to 30.1	133,329
	parsley	33.1 to 66.3	293,920
	purslane (garden, winter)	122.7 to 245.4	1,088,400
	radicchio	27.1 to 54.1	240,000
	spinach (baby)	5.11 to 10.23	45,360
	Full Season Crops	Full Season Product Rate fl ozs/100 lbs seed ¹	Seed Size seeds/lb ¹
	lettuce ² (head, leaf, Romaine)	847.5 to 1,589 (0.4 to 0.75 mg ai/seed)	469,933
	spinach (spinach, New Zealand, vine)	30.68 (0.15 mg ai/seed)	45,360

(continued)

Leafy Greens (subgroup 4A)

(continued)

¹ Fl ozs/100 lbs seed is calculated based on seed size (seeds/lb).

² For high application rate of 0.4 to 0.75 mg ai/seed for lettuce, incorporate **Poncho® 750 seed treatment** into pelleting process. Minimum pelleting buildup weight of 40X is required for the highest rate of 0.75 mg ai/seed. The minimum pelleting buildup weight is allowed to be reduced proportionally [buildup = 40X (target rate/0.75)] for the lower rate. **DO NOT** apply the product on raw seeds.

Crop-specific Restrictions

- Each fluid ounce of **Poncho 750** contains 22.18 grams of clothianidin.
- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Rate Guide Chart

Maximum seeding rates are based on the application rate and range from 120,000 to 3.6 million seeds per acre.
(see **Rate Guide Chart** below)

Application Rate (mg ai/seed)	Maximum Seeding Rate (seeds/A)
0.025	3,600,000
0.05	1,800,000
0.15	600,000
0.4	225,000
0.75	120,000

Leek

Insect Pests	Product Rate fl oz/1,000 seeds (mg ai/seed)
Onion maggot Seedcorn maggot Thrips	0.009 (0.2)

Crop-specific Restrictions

- **Poncho® 750 seed treatment** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Onion (bulb)

Insect Pests	Product Rate fl oz/1,000 seeds (mg ai/seed)
Onion maggot Seedcorn maggot Thrips	0.0053 to 0.0081 (0.118 to 0.18)

Crop-specific Restrictions

- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Onion (bunching)

Insect Pests	Product Rate fl oz/1,000 seeds (mg ai/seed)
Onion maggot Seedcorn maggot Thrips	0.004 to 0.0081 (0.09 to 0.18)

Crop-specific Restrictions

- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert..

Rice

Insect Pests	Product Rate fl ozs/100 lbs seed (mg ai/seed) ¹
Aphids Black bug Chinch bug Grape colaspis Rice water weevil Thrips	1.54 (0.01832)

Provides early season protection against listed pests.

¹ Rate is calculated based on number of seeds/lb averages of rice (18,650).

Crop-specific Restrictions

- **DO NOT** apply to wild rice.
- Not registered for use in California.
- **Poncho® 750 seed treatment** is to be used only as a dry-seeded rice seed treatment.
- **DO NOT** use **Poncho 750** in areas where rice/crawfish aquaculture practices are in place or in or near fish farms; shrimp, prawn, or crab ponds; or nursery operations.
- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Sorghum

Insect Pests	Product Rate fl ozs/100 lbs seed (mg ai/seed) ¹
Aphids (including greenbug, corn leaf aphid, English grain aphid, sugarcane aphid, yellow sugarcane aphid) Black granulate cutworm Chinch bugs White grubs Wireworms	4.06 to 4.87 (0.05 to 0.06)

¹ Rate is calculated based on number of seeds/lb averages of sorghum (18,000).

Crop-specific Restrictions

- Not registered for use in California.
- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

Soybean

Insect Pests	Product Rate fl oz/140,000 seeds (mg ai/seed)
Aphids Bean leaf beetle Grape colaspis Leafhoppers Seedcorn maggot Three-cornered alfalfa hopper Thrips White grubs Wireworms	0.69 (0.11)

Crop-specific Restrictions

- Not registered for use in California.
- Foliar sprays of clothianidin can be applied 45 days after planting **Poncho® 750 seed treatment**-treated seeds for mid-to-late season foliar insects.

Sugar Beet

Insect Pests	Product Rate fl ozs/100,000 seeds (mg ai/seed)
Black bean aphid Cutworms Flea beetle Garden springtail Sugar beet leafhopper Sugar beet leafminer Sugar beet root maggot Wireworms	2.72 (0.603)

Crop-specific Directions

- Protection against beet leafhopper injury aids in reducing potential spread of beet curly top virus, vectored by beet leafhopper.
- Apply **Poncho 750** to seed with a coating system giving a minimum weight gain buildup of approximately 30%.

Crop-specific Restrictions

- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

On-farm Seed Treatment

Cereal Grains (except corn, rice, sorghum, wild rice)

barley, buckwheat, millet (pearl, proso), oats, rye, teosinte, triticale, wheat

Insect Pests	Product Rate fl ozs/100 lbs seed (mg ai/seed) ¹
Wireworms	0.20 to 1.43 (0.00324 to 0.02318)
Aphids (including bird cherry-oat, English grain, greenbug, Russian wheat aphid)	0.60 to 1.43 (0.00971 to 0.02318)
Grasshoppers Hessian fly	1.43 (0.02318)

¹ Rate is calculated based on number of seeds/lb averages of wheat (13,700).

Crop-specific Directions

- **Wireworm seed protection** - Use 0.2 fl oz/100 lbs seed rate or higher, depending on field history and site populations.
- **Hessian fly protection to wheat seedlings in the Fall season** - Use 1.43 fl ozs/100 lbs seed.
- For maximum effectiveness, treat seed uniformly. Use the higher rate to provide increased length of protection and from heavy insect pressure and to reduce potential spread of barley yellow dwarf virus due to aphid vectors.
- Grasshopper feeding damage to seedling plants under low-to-moderate insect pressure will be reduced when **Poncho® 750 seed treatment** is used. Treated seed may be planted as a 50 to 60 foot border around the edges of the crop field as a barrier to grasshoppers advancing into the interior of the field.

Crop-specific Restrictions

- Regardless of application method (seed treatment, soil, or foliar), **DO NOT** apply more than 0.2 lb active ingredient clothianidin per acre per year.
- **Poncho 750** is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.

On-farm Seed Piece Treatment

Potato

Insect Pests	Product Rate fl oz/100 lbs seed (mg ai/seed) ¹
Aphids (including potato, green peach, foxglove, buckthorn aphids) Colorado potato beetle Potato flea beetle (overwintered adults and suppression of second generation) Potato leafhopper	0.128 to 0.256 (3.55 to 7.1)
Wireworms (seed piece protection only)	0.256 (7.1)

¹ Rate is calculated based on number of seeds/lb averages of potato seed pieces (8).

Crop-specific Directions

- Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust.
- Apply specified dosage as a diluted spray onto seed pieces using a shielded spray system that is well contained and will prevent the loss of any liquid.
- Agitate or stir spray solution as needed. For optimal insect control, good coverage of the seed pieces is required.
- As part of the seed cutting and treating process, treatment and/or an inert absorbent ingredient is recommended to improve suberization.
- Plant seed pieces as soon as possible after cutting and treating.
- Consult local specialists, BASF representative or dealer for information relevant to your area.
- For extended residual control of pests other than wireworms, apply the higher rate.

Crop-specific Restrictions

- **DO NOT** enter or allow worker re-entry into treated areas during the restricted-entry interval of 12 hours.
- Regardless of formulation or type of application (seed treatment, soil, or foliar), **DO NOT** apply more than a total of 0.2 lb active ingredient clothianidin per acre per year.
- **DO NOT** apply any subsequent application of a **Group 4** insecticide (i.e., infurrow or foliar application) following treatment with **Poncho® 750 seed treatment** insecticide.
- **DO NOT** dilute with any more than 6 parts water to 1 part **Poncho 750** insecticide.

Rotational Crop Plant-back Intervals					
Crop Planted	Immediate Plant-back	30-day Plant-back	4-month Plant-back	8-month Plant-back	12-month Plant-back
broccoli, carrot, cereal grains, field corn, popcorn, cotton, leek, onions, potato, soybean	canola, cereal grains, cotton, soybean, Brassica (cole) vegetables, bulb vegetables, cucurbit vegetables, fruiting vegetables, leafy green vegetables, root and tuber vegetables	dry beans, grasses, nongrass animal feeds		Sugarcane	Any crop without an earlier plant-back interval
canola, rapeseed, sweet corn, sorghum			Any crop without an earlier plant-back interval		

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry secured storage area.

Pesticide Disposal

To avoid waste, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Dilutable Seed Treatment Products in Nonrefillable Plastic Containers

Rigid nonrefillable containers (equal to or less than 5 gallons)

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

(continued)

STORAGE AND DISPOSAL (continued)

Container Handling (continued)

Dilutable Seed Treatment Products in Nonrefillable Plastic Containers (continued)

Rigid nonrefillable containers that are too large to shake

(i.e., with capacities greater than 5 gallons or 50 lbs) Nonrefillable Container. DO NOT reuse or refill this container. After emptying product from container, either return container to BASF, or rinse and either recycle or dispose of the container as follows:

Bottom discharge Intermediate Bulk Container (IBC) (e.g. Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top discharge IBC, drums, kegs (e.g. Snyder 120 Next Gen, Bonar B120, drums, and kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

SEED TAG LABELING

The Federal Seed Act requires that the container of seed treated with **Poncho® 750 seed treatment** must be labeled with the following statements:

- This seed has been treated with **Poncho® 750 seed treatment (EPA Reg. No. 7969-XXX)**, which contains **57.25% clothianidin**.
- **DO NOT** use treated seed for food, feed, or oil production.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with **Poncho® 750 seed treatment**:

- **Store treated seeds away from food and feedstuffs.**
- **DO NOT allow children, pets, or livestock to have access to treated seed.**
- Wear long-sleeved shirt, long pants and chemical-resistant gloves made out of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils or Viton ≥ 14 mils when handling treated seed.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agromonic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Promptly cover or collect treated seed spilled during loading or planting.
- Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inches or greater to minimize exposure.
- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment washwater.
- Dispose of seed packaging or containers in accordance with local requirements.
- **DO NOT** use empty seed bags for any other purpose.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's specifications. Surplus seed or empty seed containers must be stored or disposed according to local federal regulations.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust, which can drift to blooming crops or weeds.

(continued)

SEED TAG LABELING (continued)

- This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Promptly cover or collect treated seed spilled during loading or planting.
- Rapeseed greens and seed grown or harvested from **Poncho® 750 seed treatment**-treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from **Poncho® 750 seed treatment**-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.
- In soybean, foliar sprays of clothianidin can be applied 45 days after planting **Poncho® 750 seed treatment** treated seeds for mid-to-late season foliar insects.
- For all other crops, **DO NOT** apply a neonicotinoid insecticide foliar spray after planting treated seed. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.
- **Cereal grains (except corn, rice, sorghum, wild rice) and potato**
 - Treated areas may be replanted with any crop specified on clothianidin label or any crop for which a tolerance exists for clothianidin, as soon as practical following the last application.
 - Areas planted with treated seed may be replanted immediately with canola, rapeseed, cereal grains (crop group 15 except rice), corn, cotton, potato, sorghum, sugar beet, and the following vegetable crops: *Brassica* (cole) leafy vegetables (crop group 5), bulb vegetables (crop group 3), cucurbit vegetables (crop group 9), fruiting vegetables (crop group 8), leafy green vegetables (crop group 4A), root vegetables (crop group 1).
 - These areas may also be replanted after 30 days with dry beans, dry peas, and soybean.
 - **DO NOT** plant any other crop in the treated area for at least one year after treated seeds are planted.
- **Cereal grains for grasshoppers**
 - Feeding damage to seedling plants under low-to-moderate insect pressure will be reduced when **Poncho® 750 seed treatment** is used. Treated seed may be planted as a 50 to 60 foot border around the edges of the crop field as a barrier to grasshoppers advancing into the interior of the field.
- **Field corn, popcorn**
 - Immediate plant-back: canola, cotton, rapeseed, cereal grains, field corn, popcorn, sweet corn, root and tuber vegetables, soybean
 - 30 day plant-back: dry beans, grasses, nongrass animal feeds
 - 8 month plant-back: sugarcane
 - 12 month plant-back: any crops without an earlier plant-back interval

(continued)

SEED TAG LABELING *(continued)*

- **Sweet corn, canola, rapeseed, sorghum**
 - Immediate plant-back: canola, cotton, rapeseed, cereal grains, field corn, popcorn, sweet corn, root and tuber vegetables, sorghum, soybean
 - 30 day plant-back: dry beans, grasses, nongrass animals feeds
 - 4 month plant-back: any crop without an earlier plant-back interval
- **Soybean**
 - Immediate plant-back: canola, rapeseed, cereal grains, field corn, popcorn, sweet corn, cotton, root and tuber vegetables, soybean
 - 30 day plant-back: dry beans, grasses, nongrass animal feeds
 - 8 month plant-back: sugarcane
 - 12 month plant-back: any crops without an earlier plant-back interval
- Foliar spray of clothianidin can be applied 45 days after planting **Poncho® 750 seed treatment**-treated soybean seeds for mid-to-late season foliar insects.
- Follow-up foliar sprays of a registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.
- **Restrictions:**
 - **DO NOT** broadcast plant seeds.
 - **DO NOT** apply more than 0.2 lb ai/A of clothianidin per year (including seed treatment, foliar, or in-furrow).
 - **DO NOT** use **Poncho® 750 seed treatment** in areas where rice/crawfish aquaculture practices are in place or in or near fish farms; shrimp, prawn, or crab ponds; or nursery operations.
 - Not for use on sorghum or rice seed in the state of California.
 - **Poncho® 750 seed treatment** is to be used only as a dry-seeded rice seed treatment.
 - **DO NOT** graze or feed soybean forage and hay to livestock.
- **Broccoli** - Treated at the rate of XX fluid ounces per 1,000 seeds will result in XX grams of clothianidin per 1,000 seeds.
- **Canola, rapeseed** - Treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.
- **Carrot** - Treated at the rate of XX fluid ounces per 1,000 seeds will result in XX grams of clothianidin per 1,000 seeds.
- **Corn (field corn, popcorn)** - Treated at the rate of XX mg ai/seed.
- **Corn for pest insect southern corn billbug** - Follow-up foliar sprays of a locally registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.
- **Corn (sweet)** - Treated at the rate of XX mg ai/seed.

(continued)

SEED TAG LABELING *(continued)*

- **Cotton** - Treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.
- **Leafy greens** - Treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX mg of clothianidin per seed.
- **Leek** - Treated at the rate of XX fluid ounces per 1,000 seeds will result in XX grams of clothianidin per 1,000 seeds.
- **Onion (bulb)** - Treated at the rate of XX fluid ounces per 1,000 seeds will result in XX grams of clothianidin per 1,000 seeds.
- **Onion (bunching)** - Treated at the rate of XX fluid ounces per 1,000 seeds will result in XX grams of clothianidin per 1,000 seeds.
- **Rice** - Treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX lbs of clothianidin per 100 lbs of seed.
- **Sorghum** - Treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.
- **Soybean** - Treated with 0.11 mg clothianidin/seed.
- **Sugar beet** - Treated at the rate of XX fluid ounces per 100,000 seeds will result in XX grams of clothianidin per 100,000 seeds.
- **Cereal grains (except corn, rice, sorghum, wild rice)** - Treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as the presence of other or untested or unapproved materials, seed of low quality or low vigor or low germination, use of the product in a manner inconsistent with its labeling, misapplication of the product, or weather conditions at planting or environmental conditions during seed storage, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

ST1108

Poncho is a registered trademark of BASF.

© 2024 BASF Corporation
All rights reserved.

007969-00xxx.20230630d.NVA 2023-04-0631-0114

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry