

EFFECTIVE WHERE OTHER BAITS FAIL

DID YOU KNOW?

Left alone for one year, one pair of rats produce approximately eighty (80) young.

One (1) rat will eat approximately 27 1/2 pounds of food in one (1) year.

WARFARIN BAITS ARE NOW RECOMMENDED BY GOVERNMENT AGENCIES, HEALTH DEPARTMENTS AND AGRICULTURAL COLLEGES.

RAT BAIT

Kills Mice Too

ACQUITTED
1-25-68
7997-4

HOW TO ESTIMATE THE NUMBER OF RATS

	Number of Rats
Rat signs, none seen	100
Rat signs, night only	500-1,000
Rat signs, day time	1,000-5,000

EFFECTIVE WHERE OTHER BAITS FAIL

DID YOU KNOW?

Left alone for one year, one pair of rats produce approximately eighty (80) young

One (1) rat will eat approximately \$2.00 worth of feed in one (1) year.

WARFARIN BAIT IS NOW RECOMMENDED BY GOVERNMENT AGENCIES, HEALTH DEPARTMENTS AND AGRICULTURAL COLLEGES.

RAT BAIT

Kills Mice Too

ACCENTED
1-25-68
7997-4

HOW TO ESTIMATE THE NUMBER OF RATS.

	Number of Rats
Rat signs, none seen	100
Rats seen, night only	500-1,000
Rats seen, day time	1,000-5,000

Always insist on the rat bait with the special lure found only in the J. W. Lawton formula.

U.S.D.A. Registration
No. 7997-4

3 Pounds

1-25-68

7997-4

THE CO-OP WAY

SEASONED TO A RAT'S TASTE

RAT BAIT

Kills Mice Too

CONTAINS
PROLIN
AN IMPROVED
WARFARIN PRODUCT

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS REQUIRED

Active Ingredients:	Warfarin 3-(Alpha-Acetylbenzyl)-4-hydroxycoumarin	.025%
	N ¹ -(2-quinoloxalyl) Sulfanilamide	.025%
Inert Ingredients:	Cereal Base	99.950%
	Total	100.000%

Manufactured for

FARMERS UNION GRAIN TERMINAL ASSOCIATION - ST. PAUL, MINNESOTA

**OUR
UNCONDITIONAL
GUARANTEE**

If you follow directions, rats and mice will take GTA RAT BAIT in the presence of any other food, in any place or your money will be refunded upon return of the unused portion.

GTA RAT BAIT contains as its active ingredient an anticoagulant chemical which if taken by humans or animals may result in serious blood hemorrhages. If baits are consumed by a human, give a tablespoonful of salt in a glass of water to induce vomiting and call a physician. In case of CAT BAIT, give EMERSON'S CAT BAIT TO CHILDREN. **AVOID** ANIMALS.

NOTE TO PHYSICIAN: Treat patient as for an overdose of dicumarol.

ANTIDOTE: Use Vitamin K.

TO OPEN,
PRESS
HERE

1-25 68

7997-4

Ready to use

Directions
SET ALL BAITS AT GROUND
OR FLOOR LEVEL

RATS: Suggested use: Set out in clear containers with clamp bands at least 8 to 10 baits of 4 ounces or more. An uninterrupted supply of bait should be maintained for at least 10 days. Place around or in each building in rat runways next to walls. **PROTECT FROM PETS AND CHILDREN.** Replenish the amount of bait daily until no more GTA RAT BAIT is eaten. The average rat eats around 1/2 cup of bait per day. **1 YEAR APPROXIMATE** consumption of bait is 1/2 cup per rat. Dispose of bait promptly. Containers should be kept clean and dry.

FOR ALL MICE
RETTA

Directions: Set out in clear containers with clamp bands at least 8 to 10 baits of 4 ounces or more. An uninterrupted supply of bait should be maintained for at least 10 days. Place around or in each building in rat runways next to walls. **PROTECT FROM PETS AND CHILDREN.** Replenish the amount of bait daily until no more RETTA is eaten. The average mouse eats around 1/2 cup of bait per day. **1 YEAR APPROXIMATE** consumption of bait is 1/2 cup per mouse. Dispose of bait promptly. Containers should be kept clean and dry.

**OUR
UNCONDITIONAL
GUARANTEE**

If you follow directions, rats and mice will take GTA RAT BAIT in the presence of any other food, in any place or your money will be refunded upon return of the unused portion.

GTA RAT BAIT contains as its active ingredient an anticoagulant chemical which if taken by humans or animals may result in serious blood hemorrhages. If baits are consumed by a human, give a tablespoonful of salt in a glass of water to induce vomiting and continue until vomit is clear. Call a physician at once. PLACE BAITS IN LOCATIONS NOT ACCESSIBLE TO CHILDREN, PETS OR DOMESTIC ANIMALS.

NOTE TO PHYSICIAN: Treat patient as for an overdose of dicumarol.

ANTIDOTE: Use Vitamin K.

**THIS BAIT IS THE
EXCLUSIVE GUNNISON
FORMULA**

PRESS
HERE

1-25-68

7997-4

Ready to use

**Directions
SET ALL BAITS AT GROUND
OR FLOOR LEVEL**

RATS: Suggested use: Set out in clean containers with clean hands at least 8 to 10 baits of 4 ounces or more. An uninterrupted supply of bait should be maintained for at least 10 days. Place around or in each building in rat runways next to walls. **PROTECT FROM PETS AND CHILDREN.** Increase the amount of bait daily until no more GTA RAT BAIT is eaten. Then leave two baits around each building **THE YEAR AROUND.** Contaminated or spoiled baits should be replaced promptly. Constant and continued feeding is what kills.

**FOR FAST RESULTS
REPLACE OLD BAIT.**

MICE: Tablespoon amounts (1 to 2 ounces) of bait should be placed at 8-12 foot intervals next to the wall and out of sight. Replace bait daily until no more is taken, or 15 days.