



(Contains RABON® Oral Larvicide)

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl	
dimethyl phosphate	0.18%
INERT INGREDIENTS	99.82%
Total	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

(Continued on back of tag.)

EPA Reg. No. 7997-11
U.S. Patent No. 3,102,842
Made in U.S.A.

EPA Est. No. 7997
Net Weight 50 lbs.
X-2738-6

Manufactured by
FARMERS UNION GRAIN TERMINAL ASSN.
General Offices
ST. PAUL, MINNESOTA 55165

ACCEPTED

AUG 31 1976

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7997-11

Flp

0 - 1 2 3 4 5 6 7 8 9

15

18

Feed ROL-20 to beef cattle weighing 400-1200 pounds at the rate of 1 pound per head per day blended into the total ration.

Start feeding early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

ROL-20 may be fed up to slaughter.

ROL-20 prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around feedlots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, ROL-20 should be used in conjunction with other good management and sanitation practices.

GUARANTEED ANALYSIS

Crude Protein, not less than	20.0%
(This includes not more than 6.5% equivalent crude protein from non-protein nitrogen).	
Crude Fat, not less than	2.5%
Crude Fiber, not more than	9.0%
Calcium (Ca), not less than	1.25%
Calcium (Ca), not more than	2.25%
Phosphorus (P), not less than	1.0%
Salt (NaCl), not less than	1.25%
Salt (NaCl), not more than	2.25%
Cobalt (Co), not less than	0.0004%
Iodine (I), not less than	0.001%
Potassium (K), not less than	1.5%
Sulfur (S), not less than	0.1%
Zinc (Zn), not less than	0.025%
Vitamin A, minimum USP units/lb.	30,000
Vitamin D3, minimum USP units/lb.	6,000

INGREDIENTS

Forage Products, Plant Protein Products, Processed Grain By-Products, Cane Molasses, Urea, Mono-ammonium Phosphate, Di-calcium Phosphate, Salt, Calcium Carbonate, Vitamin A Supplement, D-Activated Animal Sterol (source of vitamin D3), Bentonite, Manganese Oxide, Zinc Oxide, Iron Sulfate, Copper Oxide, Cobalt Carbonate, Magnesium Oxide, Calcium Sulfate, Potassium Chloride, Sulfur, Ethylene Diamine Dihydriodide, and Ethoxyquin (a preservative).

40

42

46

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

X-2738-6

1 2 3 4 5 6 7 8 9 10 11 12

30

29

28

27

26

25

24

23

22

21

20

19

18

17

16

15

14

13

12

11

10

9

8

7

6

5

4

3

ACCEPTED

AUG 31 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7997-11.

klp