



**(Contains RABON® Oral Larvicide)**

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenoxy) ethyl dimethyl phosphate	1.79%
INERT INGREDIENTS	98.21%
Total	100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
CAUTION**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**GUARANTEED ANALYSIS**

Calcium (Ca), not less than	12.0%
Calcium (Ca), not more than	14.0%
Phosphorus (P), not less than	12.0%
Magnesium (Mg), not less than	2.5%
Iodine (I), not less than	0.1%
Zinc (Zn), not less than	0.1%
Cobalt (Co), not less than	8.0%
Vitamin A, minimum USP units/lb	2,000,000
Vitamin D3, minimum USP units/lb	2,000,000

(Continued on back of tag.)

EPA Reg. No. 7997-12  
U.S. Patent No. 3,102,842  
Made in U.S.A.

EPA Est. No. 7997  
Net Weight 50 lbs.  
M 2579-6

**ACCEPTED**

JUL 13 1976

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 7997-12

Manufactured by  
**FARMERS UNION GRAIN TERMINAL ASSN.**  
General Offices  
ST. PAUL, MINNESOTA 55165

9  
8  
7  
6  
5  
4  
3  
2  
1  
0  
31

15  
18  
20  
23

**INGREDIENTS**

Di-calcium Phosphate, Calcium Carbonate, Iron Oxide, Manganous Oxide, Iron Sulfate, Zinc Oxide, Copper Oxide, Ethylene Diamine Dihydriodide, Sulfur, Cobalt Carbonate, Magnesium Oxide, Vitamin A Supplement, D-Activated Animal Sterol (source of vitamin D3), Corn Meal, Cane Molasses, Natural and Artificial Flavors Added.

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed ROL-179 Mineral at the rate of one ounce per 750 pounds of body weight per head per day by mixing it into the ground grain or complete daily ration.

Start feeding ROL-179 Mineral early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

ROL-179 Mineral may be fed up to stud mares and to lactating dairy cows.

ROL-179 Mineral prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, ROL-179 Mineral should be used in conjunction with other good management and sanitation practices.

53

55

40

42

46

**STORAGE AND DISPOSAL**

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

M-2579-6  
1 2 3 4 5 6 7 8 9 10 11 12

**ACCEPTED**  
JUL 13 1976  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7997-12

cuo