

ROL-38

(Contains RABON® Oral Larvicide)

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.18%
INERT INGREDIENTS	99.82%
Total	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

(Continued on back of tag.)

EPA Reg. No. 7997-13
U.S. Patent No. 3,102,842
Made in U.S.A.

EPA Est. No. 7997
Net Weight 50 lbs.
X-2739-6

Manufactured by
FARMERS UNION GRAIN TERMINAL ASSN.
General Offices
ST. PAUL, MINNESOTA 55165

ACCEPTED
AUG 31 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7997-13

FUP

6 8 7 9 5 4 3 2 1 0 31

15 30 29 28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 0 31

18 20 23 26 29 30

DAIRY CATTLE
Feed ROL-38 at the rate of 1 pound per animal per day to animals weighing between 400-1200 pounds by incorporating it into the daily ration. Cattle weighing between 1200-1800 pounds should be fed 1 1/2 pounds per day.

ROL-38 can be fed to dry cows, replacement animals and lactating cows.

BEEF CATTLE
Feed ROL-38 to feeder cattle weighing 400-1200 pounds at the rate of 1 pound per head per day blended into the total ration.

ROL-38 may be fed up to slaughter.

GENERAL RECOMMENDATIONS
Start feeding ROL-38 early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

ROL-38 prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but it is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around dry lots, calf pens, barns, feedlots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, ROL-38 should be used in conjunction with other good sanitation and management practices.


GUARANTEED ANALYSIS

Crude Protein, not less than	38.0%
(This includes not more than 13.0% equivalent crude protein from non-protein nitrogen).	
Crude Fat, not less than	1.0%
Crude Fiber, not more than	8.0%
Calcium (Ca), not less than	1.0%
Calcium (Ca), not more than	2.0%
Phosphorus (P), not less than	1.25%
Salt (NaCl), not less than	1.50%
Salt (NaCl), not more than	2.50%
Cobalt (Co), not less than	0.001%
Iodine (I), not less than	0.0025%
Vitamin A, minimum USP units/lb.	15,000
Vitamin D3, minimum USP units/lb.	3,000

INGREDIENTS
Plant Protein Products, Processed Grain By-Products, Forage Products, Cane Molasses, Urea, Di-calcium Phosphate, Salt, Calcium Carbonate, Bentonite, Vitamin A Supplement, D-Activated Animal Sterol (source of vitamin D3), Manganous Oxide, Iron Sulfate, Zinc Oxide, Copper Oxide, Cobalt Carbonate, Magnesium Oxide, Sulfur, Ethylene Diamine Dihydriodide, and Ethoxyquin (a preservative.)

STORAGE AND DISPOSAL
Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

X-2739-6
1 2 3 4 5 6 7 8 9 10 11 12



Flup