



FORMULA NO. 1

"EASY" CATTLE OIL

Contains 5% Toxaphene for use on the range and in the feedlot. Aids in the control of lice and for the reduction of hornflies. Improves the condition of hair coat on the animal

ACTIVE INGREDIENTS

*Toxaphene	5%
Refined Mineral Oil	95%
	<hr/>
	100%

*Technical Chlorinated Camphene (Chlorine Content—67% to 69%)
USDA REG. NO. 8006-11Net Gal.

CAUTION Keep Out of Reach of Children

See Antidote Statement and other required warning statements on other panels.

EASY CHEMICAL & MFG. CO., INC.

R.R. 1 SEWARD, NEBRASKA 68434 Phone 402-643-4950

DIRECTIONS FOR USE

"Easy" Cattle Oil is ready to use; do not dilute. For use in all types of cattle oilers and back-rubbing devices. In automatic oilers, refill when necessary and make available to from 50 to 100 head. With burlap oilers, use one gallon of Easy Cattle Oil to 20 feet of oiler; recharge when necessary.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CORRY'S

SLUG & SNAIL DEATH

General Applications

Corry's Slug and Snail Death is so easy to use. Requires no mixing, spraying, or special applicators. Just place small heaps about the size of a fifty-cent piece on the soil at frequent intervals in infested areas. These small piles of bait will then actually attract the slugs and snails in the immediate area and kill them. All likely places should be treated as these pests travel. They favor damp places around all vegetable plants such as beans, tomatoes, lettuce, celery, and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra lawns, ivy patches, and other ground cover where they obtain shelter by day. Use approximately one pound per 1750 square feet.

Corry's Slug and Snail Death applied in the above manner will withstand rain and sprinkling and remain effective, protecting your garden for several weeks.

Where slug and snail control is needed but the presence of children or pets is a concern, we suggest the broadcast method of application, as listed below.

*Special Applications

*To treat an area of particularly heavy infestation, broadcast lightly over the ground. When used in this manner the bait will give exceptional results but may have to be renewed more often.

*If slugs and snails invading from adjacent properties are a menace, a pencil-thin line of Corry's Slug and Snail Death the entire length of the infested area will form a barrier and result in excellent control.

For best results be sure area to be baited is damp. In very dry weather water well before baiting.

Bait should be renewed at least every two weeks—more often during

Original ENGLISH FORMULA

CORRY'S

SLUG & SNAIL DEATH

CUT AND
LIFT

Slugs and snails are two of our most destructive agricultural pests. Although their outward appearance is quite different they are essentially the same creature, except that the slug does not have the external shell of the snail.

Slugs and snails feed almost entirely at night, in the evening, or during dark days. They prefer moist, warm conditions and rarely venture from their hiding places during cold weather or during exceedingly dry or hot spells.

Slugs and snails hatch from eggs, laid in clusters, under debris, stones, or beneath the surface of the soil. Generally they will hatch out after 20 to 30 days of favorable weather conditions after which time they will continue to grow slowly, eating our most tender and succulent vegetation.

While these small slugs and snails are growing and eating they migrate toward better feeding and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snail problem. However, with the use of CORRY'S SLUG and SNAIL DEATH and a consistent baiting program, these pests can be reduced to where they no longer present a major threat to our gardens.