

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W. Washington, D.C. 20460

NOTICE OF PESTICIDE: X Registration Reregistration

EPA Reg. Number

8033-35

Date of

APR

5 2005

Term of Issuance:

Conditional

Name of Pesticide Product:

Acetamiprid (F5025) 0.35% Protein Bait Gel PMP

Name and Address of Registrant (include ZIP Code):

Nisso America Inc.

45 Broadway, Suite 2120

New York, NY 10006

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(a)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
 - 2. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 8033-35". b Add an electrical to shock in "do not use....where the possibility of a shock hazard exists" the end sentence of the Crack and Crevice Application section.
 - 3. You must provide storage stability and corrosion characteristics data within one year from the registration date of this product.

A copy of the label stamped "accepted with comments" is enclosed for your records.

Signature of Approving Official:

Akiva Abramovitch, Ph.D.

Chemist

Insecticide-Rodenticide Branch

Registration Division (7505C)


Date:

APR 5 2005

EPA Form 8570-6

ACCEPTED with COMMENTS

DRAFT EPA File Symbol 8033-GL Version March 29, 2005



In EPA Letter DatedoiRections for use:

5 2005 It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Acetamiprid (F5025) 0.35% Arateineral Insecticid toockroaches have been seen and where they hide. Place gel as spots or beads in cracks and crevices where

. . 99.65% 100.00%

EPA Reg. No. 8033-GL

EPA Est. No.

1 oz gel per tube (28.35 g)

Stop - Read the label before use. Keep out of reach of children.

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile

(If you do not understand the label find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling

Keep out of reach of children and domestic animals. Remove contaminated clothing and wash before reuse.

Allergy information: this product contains peanuts.

FIRST AID	
If swallowed	Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambu-lance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For MEDICAL and TRANSPORTATION Emergencies Only Call CHEMTREC at 1-800-424-9300

NOTE TO PHYSICIAN

There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

For Information Regarding the Use of this Product Call 1-800-321-1FMC (1362)

Nippon Soda Co., Ltd

2-1, 2-Chome Ohtemachi , Chiyoda-ku, Tokyo 100-8165 Japan

Apply as a spot and/or crack and crevice application to spaces within structures, their walls and ceilings (both inside and out), and the immediate adjacent surrounding grounds. Such structures include, but are not limited to, residential (e.g. houses, apartments), commercial, industrial, and institutional buildings (e.g. restaurants, supermarkets, offices, hospitals, nursing homes, schools, warehouses, hotels, motels and utilities).

SPOT OR CRACK AND CREVICE APPLICATIONS

Spot treatments: This product should be applied as spots or droplets of approximately 4 to 6 mm (1/4 -1/2 gram) in diameter. Approximately 10 droplets per 10 linear ft, is recommended for use under conditions with low to moderate insect infestations. Approximately 40 droplets per 10 linear ft. is recommended for more severe infestations or where the population occurs in inaccessible locations, such as in wall voids. Under most circumstances, more numerous, smaller bait placements will provide faster and more effective control than fewer larger placements.

Crack and Crevice Treatments: This product may also be applied as a bead or thin film where appropriate. Bait should be applied into cracks and crevices in which insects hide or through which they may enter the building. Such openings commonly occur at expansion joints, between different elements of constructions, between, behind or under various types of equipment such as cabinets, refrigerators, sinks, stoves, dishwashers, and through floors, doors, windows, walls and ceilings. These openings may lead to voids such as hollow walls, hollow equipment legs and bases, utility entrances, electrical conduit, junction and switch boxes, and wall and floor drain exits. Trapping techniques or spot flushings will aid in determining areas with insect activity.

Apply this product by placing the applicator tip directly into cracks and crevices and other openings or voids where insects may hide or harbor and deposit the bait as multiple spots or as a continuous bead. Care should be taken to avoid depositing the product onto open or exposed surfaces. During follow up visits, inspect bait placements and reapply as needed. If bait is completely consumed, reapply in the same location, as many insects (such as cockroaches) leave behind secretions and droppings to attract other insects to harborages and food sources. Do not use this product where a possibility of a shock hazard exists.

INDOOR PEST CONTROL

For indoor uses, make applications in a manner as previously described above under "Spot or Crack and Crevice Applications." Particular attention should be given to treating entry points and harborage areas. Place the applicator tip into cracks, crevices, voids, holes and other small openings where insects may hide. Apply small amounts of bait (up to 1/2 gram/spot) or as a bead or film at insect activity sites. Cleaning of all areas prior to use of this product will increase levels of control. Any foodstuffs infested with insect pests should be removed and destroyed. Applications to furniture should be restricted to those areas where contact to humans will not occur. Applications to textiles or fabrics may cause discoloration or staining

OUTDOOR PEST CONTROL

Apply this product as a spot, bead or thin film to outside surfaces of buildings, porches, patios, door and window openings, crawl spaces, air conditioning units, utility entrances through outside walls, under siding/molding, in garages and other adjacent areas where pests have been seen or found. To help prevent infestation, treat areas adjacent to structures where pests may be active and may find nesting sites or harborage. Applications should be made to substrates adjacent to buildings, the building foundations, wall voids, attic and soffit vents, expansion joints, around doors and windows, soffit areas, eaves, under stairways, around screens, on fences, trees and other areas conducive to infestation. Do not apply this product to areas that have been recently sprayed with insecticides, and do not spray insecticides over gel, as it may cause the bait to become repellent

DRAFT EPA File Symbol 8033-GL Version March 29, 2005

Conditions	of	Sale	and	l Liı	mita	tion	of	W	arran	ty and
Liability:										

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or conditions beyond the control or Nippon Soda Co., Ltd. or Seller. Ali such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Nippon Soda Co., Ltd. and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. Nippon Soda Co., Ltd. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or Nippon Soda Co., Ltd., and buyer assumes the risk of any such use.

In no event shall Nippon Soda Co., Ltd. or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF Nippon Soda Co., Ltd. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF Nippon Soda Co., Ltd. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Nisso is a trademark of Nippon Soda Co., Ltd.

Species/Level of Infestation	Application Rate (Per Linear ft.)					
Moderate infestations or reapplication for German or Brown-Banded cockroaches:	1-2 spots about the size of the gel syringe cap or 1 1/2"-3" bead					
Severe or heavy infestations for German or Brown-Banded cockroaches: Or Large cockroaches such as American, Smokey-Brown or Oriental:	2-4 spots about the size of the gel syringe cap or 3"-6" bead					

Recommended spot size (1/4-1/2 gram)



FOOD/FEED HANDLING ESTABLISHMENT APPLICATIONS

Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only. Apply with syringe applicator directly into cracks and crevices. Place applicator tip 1/2 inch into cracks, crevices, holes and other small openings where cockroaches may be a problem. Apply small amounts of gel (up to 1/2 gram) per spot. Do not apply bait to areas where food/feed, food utensils or food processing surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation and prepared food holding surfaces as bait may be removed by washing. Do not apply bait to surfaces where the temperature exceeds 130F as bait may run. Examples include, but are not limited to, portions of stoves, ovens, grills, fume hoods, heat lamps, coffee urns, steam tables, toasters, fryers, dishwashers and hot water pipes. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface. Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation. Apply bait in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerator, cabinets, sinks, closets, stoves and other equipment where cockroaches hide. During follow-up visits, inspect bait placements and reapply when necessary. Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Development of resistance cannot be predicted. Therefore, use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in a cool, dry area out of reach of children. **DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.