

8033-94

01/14/2009

1/9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 14 2009

John J. Wrubel
Regulatory Affairs Director
Nisso America Inc.
45 Broadway, suite 2120
New York, NY 10006

Subject: Label Amendment: Addition of ornamental and vegetable transplants
Tristar 30 SG
EPA Reg. No. 8033-94, 8033-23, 8033-36
Submission dated April 29, 2008
Decision No. 392945

Dear Mr. Wrubel:

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following comments:

- 1) On page 3 revise the statement "NOT FOR HOMEOWNER USE" to "**NOT INTENDED FOR HOMEOWNER USE**".
- 2) On page 4, under Ornamentals and Flowering Plants Use Restrictions, reduce the number of applications from "5" to "4" to comply with the maximum of 0.55 lb a.i. allowed per acre per year with maximum single application of 0.15 a.i. per acre.
- 3) On page 6 Under the Restrictions and precautions for vegetable transplants:
 - Modify the first bullet to read, "Do not make more than 1 application per crop **prior to transplant.**"
 - Delete the second restriction starting with "Do not reapply.....thresholds."
 - Specify that the total maximum seasonal use rate for each crop, including pre-transplant application can not exceed the maximum post-transplant season use rate for each respective commodity.
- 4) The following applies to all acetamiprid labels bearing vegetable crops (leafy vegetables, fruiting vegetables, cole crops, cucurbits, and onions and other bulb vegetables) such as 8033-23 and 8033-36:

Amend the label directions to specify that the total maximum seasonal use rate for each crop, including pre-transplant application can not exceed the maximum post-transplant season use rate for each respective commodity. For example on page 11 of EPA Reg. No. 8033-36, under the Restrictions and Precautions for Fruiting Vegetables, add "Do not exceed a total of 0.3 lb active ingredient acetamiprid (16ozs product) per acre per growing season **including the pre-transplant application (.15 lb active ingredient or 8ozs product)**".

In order to prevent applications to field vegetable crops above the maximum rate allowed, submit revised labels for products 8033-23 and 8033-36 within one month from the date of this letter. In addition the revised stamped label for EPA Reg. Nos. 8033-23 and 8033-36 must be provided to all distributors when the product EPA Reg. No. 8033-94 is released for shipment.

5) An Immunotoxicity study per CFR 40 Part 158, Test Guidelines (OPPTS 870.7800) must be submitted within 18 months from the date of this letter.

A stamped copy "accepted with comments" is enclosed for your records. Please submit two copies of your final printed label before you release the product for shipment. Your release for shipment of the product with the revised label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6 (e). If you have any questions please call me at (703) 308-8328.

Sincerely yours,



Akiva Abramovitch, Ph.D.

Chemist

Insecticide-Rodenticide Branch
Registration Division (7505P)

TriStar® 30 SG Insecticide

ACCEPTED
with COMMENTS
In EPA Letter Dated:

JAN 14 2009

GROUP	4A	INSECTICIDE
-------	----	-------------

ACTIVE INGREDIENT: Acetamiprid.....

INERT INGREDIENTS:.....

EPA Reg. No. 8033-94

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
8033-94

30% by wt.
70% by wt.

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label find someone to explain it to you in detail.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-424-9300
For PRODUCT USE Information Call 1-212-490-0350

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-424-9300. Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p>NOTE TO PHYSICIAN: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.</p>	

FOR ADDITIONAL PRECAUTIONARY STATEMENTS SEE INSIDE BOOKLET

CAUTION

HAZARDS TO HUMANS (and DOMESTIC ANIMALS)

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing. Keep out of reach of children and domestic animals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long sleeved shirt and long pants, socks, shoes, and gloves. Wear protective eyewear. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and chemical resistant headgear for overhead exposure.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treated area. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not contaminate water used for irrigation or domestic purposes.

SPRAY DRIFT MANAGEMENT: The interaction of many equipment factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

IMPORTANCE OF DROPLET SIZE FOR OUTDOOR USE: An important factor influencing drift is droplet size. Small droplets (< 150 to 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Spray should be released at the lowest possible height consistent with good pest control.

DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in any manner inconsistent with its labeling.
Read entire label before using this product.**

Ornamentals and vegetable transplant use only; not for woodlands or forest management.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, waterproof gloves and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated areas until dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

STORAGE

Do not store in or around the home. Store unused product in a cool, ventilated, dry, locked area. Do not allow prolonged storage in areas where temperatures frequently exceed 115° F (46° C) NEVER TRANSFER THIS PRODUCT TO ANOTHER CONTAINER FOR STORAGE.

PESTICIDE DISPOSAL

Contamination with this product will render water, food or feed unfit for human or animal consumption. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available.

COMPATIBILITY

TriStar 30 SG Insecticide, when diluted with an equal volume of water, is physically compatible with a wide range of commonly used spray products, but the full range of compatibilities under local conditions is not known. Therefore, it is essential that before using TriStar 30 SG Insecticide in any tank mixture, the compatibility of the mixture be established. Add a small amount of this product to an equal volume of water in a small container and then add the other pesticide or spray product and mix thoroughly. DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.

SPECIFIC USE DIRECTIONS

NOT FOR HOMEOWNER USE

FOR USE ON ORNAMENTAL PLANTS, FLOWERING PLANTS, AND VEGETABLE TRANSPLANTS GROWN OUTDOORS AND IN GREENHOUSES, SHADEHOUSES AND LATHHOUSES: bedding plants, flowers grown for cuttings, foliage plants, potted flowering plants, ornamental trees, listed vegetable transplants and non-bearing fruit and nut trees (Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application). Do not apply to bearing fruit trees.

For control of insect pests, apply broadcast spray to obtain thorough and uniform spray coverage of the plants. Choose a finished spray volume appropriate for the size of the plants and amount of foliage which will provide thorough coverage throughout the canopy. For optimum control, allow at least 6 hours before overhead irrigation of foliage. Do not allow public use of treated area during application.

Mix TriStar 30 SG Insecticide with sufficient water and apply as a foliar spray to obtain thorough and uniform spray coverage of the plants. Choose a finished spray volume appropriate for the size of the plants and amount of foliage which will provide thorough coverage throughout the canopy. Apply as soon as insects reach treatment thresholds. See resistance management section if multiple sprays are needed.

TriStar 30 SG Insecticide mixes quickly in water. This product has been found to be compatible with many commonly used surfactants, miticides and insecticides. Check physical compatibility using the correct proportion of products when combining products without prior history of use.

Note: Since plant varieties are numerous and constantly changing and may react differently at various sites, test the product and any tankmixes on a small scale before making large-scale applications if there is not local experience.

ORNAMENTAL AND FLOWERING PLANTS

PESTS	OUNCES of TriStar 30 SG Insecticide per 100 GALLONS*	COMMENTS
Aphids , such as Green Peach, Wooly, Melon, and Cotton aphid European pine sawfly Psyllids	1.3	Apply as a full coverage foliar spray with a non-ionic spreader-sticker adjuvant. When mixing with surfactant treat a small area first to make sure the surfactant does not cause phytotoxicity.
Tentiform leaf miner Mealybugs , such as Citrus, Obscure, Longtail, Pink Hibiscus, and Maderia mealybugs Leafhoppers , such as Glassy Wing Sharpshooter and Potato Leaf Hopper	2.7	Tank mixing with a surfactant may improve the control of mealybugs. Tank mixing with a surfactant or a pyrethroid may improve control of adult whiteflies. When mixing with surfactants, treat a small area first to make sure the surfactant does not cause phytotoxicity.
Caterpillars , such as Gypsy Moth, Tobacco Bud Worm, Fall Army Worm, Southern Army Worm, Cabbage Looper, and Diamondback Moth Hard and Soft Scales , such as Caribbean Black Scale, Pine Needle Scale, Green Shield Scale, San Jose Scale, Oyster Shell Scale, Tea Scale, Fletcher Scale, Florida Wax and Indian Wax Scales, Cottony Maple Scale, Euonymus Scale and Asian Cycad Scale Plant bugs, Adelgids Whiteflies , such as Greenhouse, Sweet Potato, Silverleaf, Banded, and Giant	2.7 – 5.3**	
Swede Midge	2.7 – 5.3**	Apply as a preventative spray to control the first generations if Swede Midge has been found in your area. Preventative applications will decrease the chance of quick population increases later in the season.
Fungus gnat larvae , Crane fly larvae	2.7 – 5.3**	Apply as a directed spray to thoroughly wet the upper ½ to 1 inch of soil media.
Citrus thrips Other thrips , such as Cotton, Palm, and Western Flower thrips Leaf Eating Beetles (adults), such as Japanese Beetle, European Chafer and Oriental Beetle**, Strawberry Weevils	4.0 – 8.0**	Tank mixing with a surfactant will improve control. When mixing with surfactant treat a small area first to make sure the surfactant does not cause phytotoxicity.
Leaf miners , such as Chrysanthemum and Citrus Leaf Miner	6.7 – 8.0**	Rotate or tank mix with Avid®, Conserve™, Pedestal™, Distance®, Enstar®, or Talus®. Tank mixing with a surfactant may improve control. When mixing with surfactant treat a small area first to make sure the surfactant does not cause phytotoxicity.

*Product can be applied with the water volume needed to provide thorough coverage.

**Use the higher rate when insect pressure is high.

USE RESTRICTIONS AND PRECAUTIONS FOR ORNAMENTAL AND FLOWERING PLANTS

- Do not make more than 5 applications per year.
- Do not reapply more than once every 7 days. To determine if application is necessary, monitor pest densities. Consult local extension experts for thresholds.
- Do not apply more than 8 ounces of TriStar 30 SG Insecticide per acre (0.15 lbs ai/A) in a single application.
- Do not apply more than 29.3 ounces of TriStar 30 SG Insecticide per acre (0.55 lb ai/A) per year.
- Do not apply to bearing fruit trees.

7/9

VEGETABLE TRANSPLANTS

VEGETABLE TRANSPLANTS	PESTS	OUNCES of TriStar 30 SG Insecticide per100 GALLONS*	COMMENTS
LEAFY VEGETABLES (within Crop Group 4) Amaranth, Arrugula, Cardoon, Celery, Chinese celery, Celtuce, Chervil, Chrysanthemum (edible leaved, garland), Corn Salad, Cress (garden, upland), Dandelion, Dock, Endive, Florence Fennel, Lettuce (head, leaf), Orach, Parsley, Purslane (garden, winter), Radicchio, Rhubarb, Spinach (leaf, vine, New Zealand), Swiss Chard FRUITING VEGETABLES (within Crop Group 8) Eggplant, Groundcherry, Pepino, Pepper (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato COLE CROPS (within Crop Group 5) Broccoli, Chinese Broccoli (gai lon), broccoli raab (rapini), Brussel Sprouts, Cabbage, Chinese Cabbage (bok choy, napa), Chinese Mustard Cabbage (gai choy), Cavalo broccolli Cauliflower, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens CUCURBITS (within Crop Group 9): Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible), <i>Mormordica</i> spp., Muskmelon (hybrid and/or cultivars of <i>Cucumis melo</i>), Pumpkin, Squash (summer and winter), Watermelon ONIONS AND OTHER BULB VEGETABLES (within Crop Group 3-07): Chives, fresh leaves; Chinese chives, fresh leaves; daylily bulbs, Elegans hosta, Fritillaria leaves and bulbs; bulb garlic; great headed bulb garlic, serpent bulb garlic; kurrat; lady's leek; leek; wild leek; lily bulb; Beltsville bunching onion; bulb onion; Chinese bulb onion; fresh onion; green onion; macrostem onion; pearl onion; potato bulb onion; treetops onion; Welsh onion tops; shallot bulb and fresh leaves; and cultivars, varieties, and/or hybrids of these.	Aphids , such as Green Peach, Woolly, Melon, and Cotton aphid Psyllids	1.3	Apply as a full coverage foliar spray with a non-ionic spreader-sticker adjuvant. When mixing with surfactant treat a small area first to make sure the surfactant does not cause phytotoxicity.
	Mealybugs , such as Citrus, Obscure, Longtail, Pink Hybiscus, and Maderia mealybugs Leafhoppers , such as Glassy Wing Sharpshooter and Potato Leaf Hopper	2.7	Tank mixing with a surfactant may improve the control of mealybugs Tank mixing with a surfactant or a pyrethroid may improve control of adult whiteflies. When mixing with surfactants, treat a small area first to make sure the surfactant does not cause phytotoxicity
	Caterpillars , such as Gypsy Moth, Tobacco Bud Worm, Fall Army Worm, Southern Army Worm, Cabbage Looper, and Diamondback Moth Hard and Soft Scales , such as Caribbean Black Scale, Pine Needle Scale, Green Shield Scale, San Jose Scale, Oyster Shell Scale, Tea Scale, Fletcher Scale, Florida Wax and Indian Wax Scales, Cottony Maple Scale, Euonymus Scale and Asian Cycad Scale Plant bugs , Whiteflies , such as Greenhouse, Sweet Potato, Silverleaf, Banded, and Giant	2.7 – 5.3**	
	Swede Midge	2.7 – 5.3**	Apply as a preventative spray to control the first generations if Swede Midge has been found in your area. Preventative applications will decrease the chance of quick population increases later in the season.
	Fungus gnat larvae, Crane fly larvae	2.7 – 5.3**	Apply as a directed spray to thoroughly wet the upper ½ to 1 inch of soil media
	Thrips , such as Citrus Cotton, Palm, and Western Flower thrips Leaf Eating Beetles (adults), such as Japanese Beetle, European Chafer and Oriental Beetle**, Strawberry Weevils	4.0 – 8.0**	Tank mixing with a surfactant will improve control. When mixing with surfactant treat a small area first to make sure the surfactant does not cause phytotoxicity.
	Leaf miners , such as Chrysanthemum and Citrus Leaf Miner	6.7 – 8.0**	Tank mixing with a surfactant may improve control. When mixing with surfactant treat a small area first to make sure the surfactant does not cause phytotoxicity.

*Product can be applied with the water volume needed to provide thorough coverage.

**Use the higher rate when insect pressure is high.

8/9

USE RESTRICTIONS AND PRECAUTIONS FOR VEGETABLE TRANSPLANTS:

- Do not make more than 1 application per crop.
- Do not reapply to successive crops more than once every 7 days. To determine if application is necessary, monitor pest densities. Consult local extension experts for thresholds.
- Do not apply more than 8 ounces of TriStar 30 SG Insecticide (0.15 lb a.i.) per acre per crop.
- Do not apply more than 0.55 lb a.i. per acre per year of any product containing acetamiprid on any outdoor field or in any greenhouse, shadehouse, or lathhouse.
- Do not harvest for food use within seven days following the last application (7 day PHI).

Use the following Chart to convert the amount of product per 100 gallons above into smaller spray volume units to accommodate smaller volume application equipment.

Conversion Guide for Small Volume (less than 100 gallon) Application Equipment – Teaspoons

Label Rates Ounces/100 Gallon	Teaspoons TriStar 30 SG per Volume			
	3 Gal Tank	5 Gal Tank	10 Gal Tank	25 Gal Tank
1.3	1/3	1/2	1	2 2/3
2.7	2/3	1	2	5 1/3
4.0	1	1 1/2	3	8
5.3	1 1/3	2	4 1/4	10 2/3
6.7	1 2/3	2 1/2	5 1/3	13 1/3
8.0	2	3	6 1/2	16

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Avoid treating these insects with consecutive applications of insecticides within the same class of chemistry. Resistance to other neonicotinoids, such as Merit and Marathon may result in resistance to TriStar 30 SG Insecticide. Therefore, to minimize the potential for neonicotinoid resistance, rotate with other classes of insecticides. Consult your agricultural advisor for resistance management strategies and recommended pest management practices for your area.

ENDANGERED SPECIES NOTICE: Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species.

Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Nippon Soda Co., Ltd. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: NIPPON SODA CO., LTD. MAKES NO OTHER WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Nippon Soda Co., Ltd. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. NIPPON SODA CO., LTD. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT TO THE EXTENT ALLOWABLE BY APPLICABLE LAW.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT NIPPON SODA CO., LTD.'S ELECTION, THE

9/a

REPLACEMENT OF THE PRODUCT.

NET CONTENTS: BATCH No.:

Nippon Soda Co., Ltd.
2-1, 2-Chome Ohtemachi
Chiyoda-ku, Tokyo 100-8165
Japan

TriStar® is a registered trademark of Nippon Soda Co., Ltd..

Avid® is a trademark of a Syngenta Group Company.
Conserve™ is a trademark of Dow AgroSciences LLC.
Distance® is a registered trademark of Valent U.S.A. Corporation.
Enstar® is a registered trademark of Wellmark International.
Merit® and Marathon® are registered trademarks of Bayer.
Pedestal™ is a trademark of Crompton Corporation.
Talus® is a registered trademark of Nichino America Inc.