



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 21, 2019

Kirsten B Knox
Regulatory Affairs Manager
Nippon Soda Co., Ltd.
c/o Nisso America Inc.
88 Pine street, 14th Floor
New York, NY 10005

Subject: PRIA Label Amendment – Amendment to Revise the Grape Application
Product Name: Miltrex 10 SC Fungicide
EPA Registration Number: 8033-103
Application Date: 08/01/2018
Decision Number: 543357

Dear Ms. Knox:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 8033-103
Decision No. 543357

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Manjula Unnikrishnan by phone at 703-347-8520, or via email at unnikrishnan.manjula@epa.gov .

Sincerely,

A handwritten signature in black ink, appearing to read "Erik Kraft". The signature is fluid and cursive, with the first name "Erik" being more prominent than the last name "Kraft".

Erik Kraft, Product Manager 24
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



Group **U6** Fungicide

MILTREX® 10 SC FUNGICIDE

For Agricultural Use Only

Active Ingredient:

Cyflufenamid;
(Z)-N-(α-cyclopropylmethoxyimino-2,3-difluoro-6-trifluoromethylbenzyl)-2-phenylacetamide.....10%

Other

Ingredients:.....90%

Total:.....100%

Contains 0.85 lbs. of cyflufenamid active ingredient per gallon

EPA Reg. No. 8033-103

EPA Est. No. _____

KEEP OUT OF REACH OF CHILDREN CAUTION

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label find someone to explain it to you in detail.)**

For Medical and Transportation Emergencies ONLY call CHEMTREC at 1-800-424-9300

FIRST AID	
If on Skin or Clothing:	<input type="checkbox"/> Take off contaminated clothing. <input type="checkbox"/> Rinse skin immediately with plenty of water for 15-20 minutes. <input type="checkbox"/> Call a poison control center or doctor for treatment advice.
If Swallowed:	<input type="checkbox"/> Call a poison control center or doctor immediately for treatment advice. <input type="checkbox"/> Have person sip a glass of water if able to swallow. <input type="checkbox"/> Do not induce vomiting unless told to by a poison control center or doctor. <input type="checkbox"/> Do not give anything by mouth to an unconscious person.
If in Eyes:	<input type="checkbox"/> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. <input type="checkbox"/> Call a poison control center or doctor for treatment advice.
If Inhaled:	<input type="checkbox"/> Move person to fresh air. <input type="checkbox"/> If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. <input type="checkbox"/> Call a poison control center or doctor for further treatment advice.
For MEDICAL EMERGENCIES call CHEMTREC at 1-800-424-9300.	
Have the product container or label with you when calling a poison center or doctor or going for treatment	
NOTE TO PHYSICIAN: There is no specific antidote. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.	

Net Contents: _____ fl.oz.

Batch No. _____

Nippon Soda Co., Ltd.
c/o Nisso America, Inc.
88 Pine Street, 14th fl.
New York, NY 10005

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves. Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests,

nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves.

COMPATIBILITY

MILTREX 10 SC FUNGICIDE, when diluted with an equal volume of water, is physically compatible with a wide range of commonly used spray products, but the full range of compatibilities under local conditions is not known. Therefore, it is essential that before using MILTREX 10 SC FUNGICIDE in any tank mixture the compatibility of the mixture be established. Add a small amount of this product to an equal volume of water in a small container and then add the other pesticide or spray product and mix thoroughly. **DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. FOR BEST RESULTS, SPRAY MIXTURES MUST BE USED IMMEDIATELY AFTER MIXING WITH ADEQUATE AGITATION.**

DIRECTIONS FOR AERIAL OR GROUND SPRAY APPLICATION

MILTREX 10 SC FUNGICIDE is a 10% suspension concentrate for the control of powdery mildew on the crops listed on this label. Thorough spray coverage is essential for optimal performance. Use adequate spray volume to ensure complete coverage of foliage.

Users must read, understand, and follow the label use rates and restrictions. Minimum label rates may be used under low disease pressure conditions while maximum label rates and shortened spray intervals are required under high disease pressure. For application, determine the number of acres to be treated, the specified label use rate, and the spray volume per acre. Prepare only the amount of spray solution that is necessary to spray the measured acres. Calibrate spray equipment prior to use.

Ground Application: For optimal disease control use thorough spray coverage. Good spray coverage is a function of spray pressure, spray volume per acre, nozzle type and spacing and application equipment speed. Calibrate spray equipment prior to use.

Aerial Application: Apply in a minimum of 5 GPA for row crops and 10 GPA for orchard crops. **Do not apply under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.**

Mandatory Spray Drift

Aerial Applications

- Do not release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the

field.

- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE
An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.
Controlling Droplet Size - Ground Boom
 - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
 - Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- Controlling Droplet Size - Aircraft
 - Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- RELEASE HEIGHT - Aircraft
Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

- WIND
Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- Boom-less Ground Applications:
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:
Take precautions to minimize spray drift.

MIXING INSTRUCTIONS

Mixing and Application Instructions for MILTREX 10 SC FUNGICIDE

MILTREX 10 SC FUNGICIDE is a liquid suspension concentrate formulation that readily disperses in water to form a spray mixture, which may be applied by ground or air.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

1. Plan ahead. Prepare only enough spray mixture as can be applied on the day of mixing.
2. Fill tank $\frac{1}{4}$ - $\frac{1}{2}$ full with the required amount of total spray volume of water.
3. Shake the product container well before using. Begin agitation and add product.
4. Continue to fill tank.
5. Allow mixing in tank for 2 minutes after filling or until thoroughly mixed before applying.
6. Maintain continuous agitation during mixing and application to assure uniform suspension. If mixture sits without agitation for extended periods, agitate the mixture for at least 10 minutes before use.
7. Equip spray system with a 50-mesh inline filter, which will protect nozzles that are typically used. Nozzles may also be equipped with 50-mesh nozzle filters or 25 to 50 mesh (equivalent) slotted nozzle filters.
8. MILTREX 10 SC FUNGICIDE may be unstable in water pH below 4 and above 9. If necessary, buffer water to obtain optimum pH range.

Special Instructions for Tank Mixing MILTREX 10 SC FUNGICIDE

When tank mixing MILTREX 10 SC FUNGICIDE with other products, introduce the products into the tank in the following order: (1) water soluble packets (2) wettable powders (3) water dispersible granules (4) flowable liquids (including MILTREX 10 SC FUNGICIDE) (5) emulsifiable concentrates and (6) adjuvants and/or oils. Always allow each product to fully disperse before adding the next product.

The use of adjuvants or additives may enhance the fungicide performance of MILTREX 10 SC FUNGICIDE under some conditions. Local environmental conditions may affect crop sensitivity. Since all possible tank mix combinations have not been examined, test the combination on a small section of the crop to be sprayed to ensure that injury will not occur as a result of application. Consult a Gowan Company representative, local agricultural authorities, or local extension service for more information and directions on adjuvants and additives.

APPLICATION INSTRUCTIONS

ROW CROPS

Apply a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground unless otherwise directed under crop specific directions. For best results, it is important to obtain thorough and uniform spray coverage of the plant. To clean the sprayer after use, drain and flush with water. Use rinsate on crop according to label instructions or dispose of in an approved manner (See STORAGE AND DISPOSAL).

ORCHARD AND VINE CROPS

To achieve optimum pest control, it is important to obtain thorough and uniform spray coverage. Choose a finished spray volume appropriate for the size of tree or vine and amount of foliage which will provide thorough coverage throughout the canopy. Also follow directions listed under crop specific directions. Aerial applications may not provide as thorough coverage as ground applications and subsequently disease control may be reduced.

To clean the sprayer after use, drain and flush with water. Use rinsate on crop according to label instructions or dispose of in an approved manner (See STORAGE AND DISPOSAL).

RESISTANCE MANAGEMENT RECOMMENDATIONS For resistance management, MILTREX 10 SC contains a Group U6 fungicide. Any fungal population may contain individuals naturally resistant to MILTREX 10 SC and other Group U6 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, consider:

- 1) Avoiding the consecutive use of MILTREX 10 SC or other Group U6 fungicides that have a similar target site of action on the same pathogens.
- 2) Using tank-mixtures or premixes with fungicides from different target site of action Groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- 3) Basing fungicide use on a comprehensive IPM program.
- 4) Monitoring treated fungal populations for loss of field efficacy.
- 5) Contacting your local extension specialist, certified crop advisor, and/or the manufacturer for fungicide resistance management and/or IPM recommendations for specific crops and resistant pathogens.

RESTRICTIONS

- Do not plant back any crop other than those on this label within 30 days following the last application.
- Do not apply this product through any type of irrigation system.

CHERRIES (within Crop Group 12-12A)

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz./acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
------	---------	---	----------------	--------------

<p>CHERRIES:</p> <p>Capulin, black, Nanking, sweet, tart, cultivars, varieties, and/or hybrids of these.</p>	<p>Powdery mildew</p>	<p>5.0 - 8.0 [0.033 - 0.053]</p>	<p>Begin application at first sign of disease development.</p>	<p>Do not make more than two (2) applications per year.</p> <p>The minimum retreatment interval is seven (7) days.</p> <p>The maximum application rate is 8.0 oz. of product (0.053 lb./ai) per acre.</p> <p>Do not exceed a total of 16.0 oz. of product (0.106 lbs. active ingredient) per acre per calendar year.</p> <p>Do not apply within six (6) days of harvest; (PHI = 6 days)</p>
---	-----------------------	--------------------------------------	--	---

CUCURBIT VEGETABLES (within Crop Group 9)

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz./acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
<p>CUCURBIT VEGETABLES including but not limited to Pumpkin, Squash (summer and winter), Watermelon, and other melons: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible), <i>Mormordica</i> spp., Muskmelon (hybrid and/or cultivars of <i>Cucumis melo</i> including true cantaloupe, cantaloupe, casaba, chreshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon).</p>	<p>Powdery mildew</p>	<p>1.7 - 3.4 [0.011- 0.022]</p>	<p>Begin application at first sign of disease development.</p> <p>For the control of other foliar cucurbit diseases, applications of registered fungicides must be made according to their label directions for use.</p>	<p>Do not make more than two (2) applications per year.</p> <p>The minimum retreatment interval is seven (7) days.</p> <p>The maximum application rate is 3.4 oz. of product (0.022 lbs. ai) per acre.</p> <p>Do not exceed a total of 6.8 oz. of product (0.044 lbs. active ingredient) per acre per calendar year.</p> <p>Applications may be made up to and including the day of harvest; (PHI = 0 days).</p>

FRUITING VEGETABLES (within Crop Group 8—10)

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz./acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
------	---------	---	----------------	--------------

<p>FRUITING VEGETABLES: Cocona, Eggplant (including African, Pea and Scarlet), Garden huckleberry, Goji berry, Ground cherry, Martynia, Naranjilla, Okra, Pepino, Peppers (including bell and non-bell), Roselle, Sunberry, Tomatillo, and Tomatoes (including Bush, Currant, and Tree), and cultivars, varieties and/or hybrids of these.</p>	<p>Powdery mildew</p>	<p>3.4 [0.022]</p>	<p>Begin application at first sign of disease development.</p>	<p>Do not make more than three (3) applications per year.</p> <p>The maximum application rate is 3.4 oz. of product (0.022 lbs. ai) per acre</p> <p>The minimum retreatment interval is fourteen (14) days.</p> <p>Do not exceed a total of 10.2 oz. of product (0.066 lbs. active ingredient) per acre per calendar year.</p> <p>Applications may be made up to and including the day of harvest.</p> <p>(PHI = 0 days)</p>
---	-----------------------	------------------------	--	--

**GRAPES (within Crop Group 13-07F)
and small fruit vine climbing crop subgroup except fuzzy kiwifruit**

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz./acre) [POUNDS ACTIVE]	SPRAY INTERVAL (DAYS)	USE DIRECTIONS	RESTRICTIONS
<p>GRAPES; Amur river grape; gooseberry; kiwifruit, hardy; maypop; schisandra berry; cultivars varieties, and/or hybrids of these</p>	<p>Powdery mildew</p>	<p>1.7 [0.011]</p>	<p>7 - 14</p>	<p>Begin application at first sign of disease development.</p> <p>For the control of other foliar diseases of grapes, applications of registered fungicides must be made according to their label directions for use.</p>	<p>Do not make more than two (2) applications per year.</p> <p>The maximum application rate is 1.7 oz. of product (0.011 lb. ai) per acre.</p> <p>The minimum retreatment interval is seven (7) days.</p> <p>Do not exceed a total of 3.4 oz. of product (0.022 lbs. active ingredient) per acre per year.</p> <p>Do not apply within three (3) days of harvest; (PHI = 3 days).</p>

		3.4 [0.022]	14 - 21		Do not make more than two (2) applications year. The maximum application rate is 3.4 oz. product (0.022 lb. ai) per acre. The minimum retreatment interval is 14 days. Do not exceed a total of 6.8 oz of product (0.044 lbs. active ingredient) per acre per calendar year. Do not apply within three (3) days of harvest; (PHI = 3 days).
		5.1 - 6.8 [0.034 - 0.044]	N/A	Use the high rate for longer residual control of powdery mildew. For the control of other foliar diseases of grapes, applications of registered fungicides must be made according to their label directions for use.	Do not make more than one (1) application year. The maximum application rate is 6.8 oz. of product (0.044 lb. ai) per acre. Do not exceed a total of 6.8 oz. of product (0.044 lbs. active ingredient) per acre per calendar year. Do not apply within seven (7) days of harvest; (PHI = 7 days)

HOPS

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz./acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
HOPS	Powdery mildew	5.0 - 8.0 [0.033 - 0.053]	Begin application at first sign of disease development.	Do not make more than two (2) applications per year. The minimum retreatment interval is seven (7) days. The maximum use rate is 8.0 oz. of product (0.053 lb. ai) per acre. Do not exceed a total of 16.0 oz of product (0.106 lbs. active ingredient) per acre per calendar year. Do not apply within six (6) days of harvest. (PHI = 6 days)

POME FRUIT (within Crop Group 11)

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz./acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
POME FRUIT: Apple, Pear, Crabapple, Loquat, Mayhaw, Quince, Oriental pear	Powdery mildew	2 applications at 3.4 [0.022] [and/or] 1 application at 6.8 [0.044]	Begin application at first sign of disease development. For the control of other foliar diseases of pome fruit, applications of registered fungicides must be made according to their label directions for use.	The minimum retreatment interval is fourteen (14) days. Do not exceed a total of 6.8 oz. of product (0.044 lbs. active ingredient) per acre per calendar year. [Do not make more than one (1) application at the 6.8 oz. product rate per calendar year] [and/or] [Do not make more than two (2) applications at the 3.4 oz. product rate per calendar year] Do not apply within 14 days of harvest; (PHI = 14 days).

STRAWBERRIES AND OTHER LOW GROWING BERRIES (within Crop Group 13-07G except cranberries)

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz./acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
STRAWBERRIES AND OTHER LOW GROWING BERRIES (except cranberries): Bearberry; Bilberry; Lowbush Blueberry; Cloudberry, Lingonberry; Muntries; Partridgeberry; and cultivars, varieties, and/or hybrids of these.	Powdery mildew	1.7 – 3.4 [0.011 – 0.022]	Begin application at first sign of disease development. For the control of other foliar diseases of strawberries and other low growing berries, applications of registered fungicides must be made according to their label directions for use.	Do not make more than two (2) applications per year. The minimum retreatment interval is fourteen (14) days. The maximum use rate is 3.4 oz. of product (0.022 lbs. ai) per acre. Do not exceed a total of 6.8 oz. of product (0.044 lbs. active ingredient) per acre per calendar year. Applications may be made up to and including the day of harvest; (PHI = 0 days).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store unused product in original container only, out of reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Non-refillable container (5 gallons or less). Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents in application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Non-refillable container (greater than five gallons). Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties, and Limitations of Liability.

CONDITIONS: The Directions for Use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Nippon Soda Co., Ltd. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Nippon Soda Co., Ltd. makes no other warranties, express or implied, or merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Nippon Soda Co., Ltd. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Nippon Soda Co., Ltd. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries, or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall not exceed the purchase price paid, or at Nippon Soda Co., Ltd's election, the replacement of product.

Miltrex® is a registered trademark of Nippon Soda Co., Ltd.



NIPPON SODA CO.,LTD.

SHIN-OHTEMACHI BLDG., 2-1, 2-CHOME OHTEMACHI
CHIYODA-KU, TOKYO 100-8165, JAPAN

Supplemental Label

ACCEPTED
05/21/2019
Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 8033-103

MILTREX 10 SC FUNGICIDE

This supplemental label expires on January 30, 2022 and must not be used or distributed after this date.

EPA Reg. No. 8033-103

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Miltrex 10 SC Fungicide before applying. Use of Miltrex 10 SC Fungicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container of Miltrex 10 SC Fungicide.

Grapes within Crop Group 13-07F

SITE	DISEASE	MILTREX 10 SC FUNGICIDE (oz/acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
GRAPES	Powdery mildew	3.4 [0.022]	Begin application at first sign of disease development. For the control of other foliar diseases of grapes, applications of registered fungicides must be made according to their label directions for use.	Do not make more than two (2) applications per calendar year. The maximum application rate is 3.4 oz. product (0.022 lb. ai) per acre. The minimum retreatment interval is 14 days. Do not exceed a total of 6.8 oz. of product (0.044 lb. active ingredient) per acre per calendar year. Do not apply within three (3) days of harvest; (PHI = 3 days)
		OR		
		5.1 - 6.8 [0.023 - 0.044]	Use the high rate during times of heavy disease pressure For the control of other foliar diseases, applications of registered fungicides must be made according to their label directions for use.	Do not make more than one (1) application per calendar year. The maximum application rate is 6.8 oz. of product (0.044 lb. ai) per acre. Do not exceed a total of 6.8 oz. of product (0.044 lb. active ingredient) per acre per calendar year. Do not apply within seven (7) days of harvest; (PHI = 7 days)