

Description

Single treatment will give knock-down of vegetation, plus long-lasting prevention of seedlings and regrowth. Especially effective on perennial grasses. Use for general mixtures of grasses. Specially potent on perennial grasses such as Johnson grass, Bermuda grass, Dallis grass, and quackgrass. Use WEED KILLER No.50 on non-crop areas, such as industrial sites, fence lines, storage yards, road sides and railroads. Use any time during the growth season. Weeds and grasses are easier to kill when they are young and growing rapidly. Uniform application over the entire area to be cleared of vegetation is essential.

IMPORTANT

This material is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops and other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlaid by roots of adjacent valuable growths. Careless application of this material, or washing by water run-off, to areas where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Do not drain or flush equipment on or near such areas. Do not contaminate domestic or irrigation water or waters used by fish and wildlife. Wash clothing and equipment after application. Keep animals off treated areas until rainfall has leached all the herbicide into the soil. Mechanical equipment should be thoroughly cleaned after use.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

WEED KILLER NO. 50

NON-SELECTIVE WEED AND GRASS KILLER
(ON INDUSTRIAL SITES OR SIMILAR
NON-CROP AREAS).

ACTIVE INGREDIENTS

| | |
|---|--------|
| *Sodium Meta Borate (Na ₂ B ₂ O ₄) | 4.29% |
| Sodium Chlorate (NaClO ₃) | 3.00% |
| Bromacil (5-bromo-3-sec-butyl-6-methyluracil) | 0.15% |
| INERT INGREDIENTS | 92.56% |
| Water | |
| *Boron Trioxide (B ₂ O ₃) equivalent 2.26% | |

Dec 18, 1975
847-38

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid contamination of feed and foodstuffs. Harmful if swallowed. May cause irritation of eyes, nose throat and skin. Avoid contact with skin and clothing. Do not store in direct contact with ground or concrete floors. Do not reuse container. Destroy when empty

MANUFACTURED BY:

POLY CHEM, INC.

P. O. BOX 100

NEW ORLEANS, LA. 70116

This pro
Use spr
to be ch
For spot
cation u

a.

b.

c.

Because
tion, sea
buyer th
the use
teed. It
mental
seller s
use of
the dire
not expr
and assu
ance wi
and age
or impli

WEED KILLER NO. 50

NON-SELECTIVE WEED AND GRASS KILLER
(ON INDUSTRIAL SITES OR SIMILAR
NON-CROP AREAS)

ACTIVE INGREDIENTS

*Sodium Meta Borate
($\text{Na}_2\text{B}_2\text{O}_4$)

4.29%

Sodium Chlorate
(NaClO_3)

3.00%

Bromacil (5-bromo-3-sec-butyl-6-methyluracil)

0.15%

INERT INGREDIENTS

Water

92.56%

*Boron Trioxide (B_2O_3) equivalent 2.26%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid contamination of feed and foodstuffs. Harmful if swallowed. May cause irritation of eyes, nose throat and skin. Avoid contact with skin and clothing. Do not store in direct contact with ground or concrete floors. Do not reuse container. Destroy when empty

MANUFACTURED BY:

POLY CHEM, INC.

P. O. BOX 1177

NEW ORLEANS, LA. 70117

Directions for use

This product is designed for spot killing of grass.

Use spraying equipment for thorough and uniform application over the entire area to be cleared of vegetation.

For spot treatment sprinkling can may be used. Use the following rates of application under most conditions.

- Annual weeds and grass such as sunflower, kochia, Russian thistle, pig-weed, crabgrass, downy brome — 1½ gallons per 100 sq. ft.
- Perennial weeds and grasses such as Johnsongrass, Bermuda grass, Dallis grass, quack grass, Canada thistle, field bindweed, bluegrass — 4½ gallons per 100 sq. ft.
- Mixed annual and perennial weeds and grasses — 4½ gallons per 100 sq. ft.

NOTICE

Because of the many varying conditions affecting use and application such as location, seasonal weather soil, composition and type of vegetation, manufacturer warns buyer that these and other conditions may impair or vary the results or effects of the use of this product. In any event, complete elimination of weeds is not guaranteed. It is advisable to consult your State Extension Service or Agricultural Experimental Station weed specialist on specific problems. Neither manufacturer nor seller shall be liable in respect to any injury or damage suffered by reason of the use of this product for a purpose not indicated by the label or when contrary to the directions or instructions hereon nor with respect to breach of any warranty not expressly specified herein. Buyer accepts this material subject to these terms, and assumes all risk of usage and handling, except when used or handled in accordance with this label. Neither manufacturer nor seller makes, and their employees and agents are not permitted to make, any representations or warranties, express or implied by law not specifically set forth herein.

EPA Reg. No. 8047-38

EPA Est. No. 8047-LA-1