

DE'S DRYATOME

specially processed diatomaceous earth for

GRAIN STORAGE INSECTICIDE

Packaged and Distributed by:

THE JOHN GALT COMPANY

SAN ANTONIO, TEXAS - 78209

Active Ingredients: Silicon Dioxide from Natural Diatomite 80%
Inert Ingredients 20%

When used as directed, DRYATOME is safe to use to protect Barley, Buck-wheat, Corn, Oats, Rice, Rye, Sorghum grain (Milo), Wheat, and mixture of these grains against the common insect pests of stored grain, including granary weevil, brain bugs, lesser grain borer, millers moth, rice weevil, Indian meal moth, flat grain beetle, saw tooth beetle and confused flour beetle.

DIRECTIONS: Use a DRYATOME screw feeder that can be adjusted to the amount of grain being mixed. The rate of DE to be applied is six (6) pounds per ton of grain. The machine holds 40 pounds of DE and can be timed to check the right amount of DRYATOME to be mixed directly with the grain.

WARNING Avoid breathing dust. Wear a respirator of the type approved by the U.S. Bureau of Mines for protection against this material. Use with adequate ventilation.

The milling, baking, and nutritive qualities of the grain are not affected by the treatment.

WARNING KEEP OUT OF THE REACH OF CHILDREN

ELEMENT APPROXIMATE PERCENT

Boron	0.06
Silicon	85.164 %
Aluminum	3.9 %
Manganese	0.002
Magnesium	1.0
Gallium	0.002
Iron	2.0
Calcium	.04 %
Vanadium	0.002
Copper	0.01
Sodium	0.0
Titanium	2.0
Zirconium	0.01
Strontium	0.01

ACCEPTED

Spec. 10, 1969

8103-1

[8108-1]

4

D E 5
D R Y A T O M E

specially processed diatomaceous earth for
GRAIN STORAGE INSECTICIDE

Packaged and Distributed by:
THE JOHN GALT COMPANY
SAN ANTONIO, TEXAS - 78209

Active Ingredients: Silicon Dioxide from Natural Diatomite 80%
Inert Ingredients 20%

When used as directed, DRYATOME is safe to use to protect Barley, Buckwheat, Corn, Oats, Rice, Rye, Sorghum grain (Milo), Wheat, and mixture of these grains against the common insect pests of stored grain, including granary weevil, bran bugs, lesser grain borer, millers moth, rice weevil, Indian meal moth, flat grain beetle, saw tooth beetle and confused flour beetle.

DIRECTIONS: Use a DRYATOME screw feeder that can be adjusted to the amount of grain being moved. The rate of DE to be applied is six (6) pounds per ton of grain. The machine holds 40 pounds of DE and can be timed to check the right amount of DRYATOME to be mixed directly with the grain.

WARNING Avoid breathing dust. Wear a respirator of the type approved by the U.S. Bureau of Mines for protection against this material. Use with adequate ventilation.

The milling, baking, and nutritive qualities of the grain are not affected by the treatment.

WARNING KEEP OUT OF THE REACH OF CHILDREN

ELEMENT	APPROXIMATE PERCENT
Boron	0.06
Silicon	85.164 *
Aluminum	3.9
Manganese	0.002
Magnesium	1.0
Gallium	0.002
Iron	2.0
Calcium	.84 *
Vanadium	0.002
Copper	0.01
Sodium	5.0
Titanium	2.0
Zirconium	0.01
Strontium	0.01
	100%

ACCEPTED

Jan 10, 1964

FEDERAL BUREAU OF INVESTIGATION
U.S. DEPARTMENT OF JUSTICE
LABORATORY NO. 8108-17

[1-8018]