

the above manner will withstand rain and sprinkling and remain effective, protecting your garden for several weeks.

Where slug and snail control is needed but the presence of children or pets is a concern, we suggest the broadcast method of application, as listed below.

*Special Applications

*To treat an area of particularly heavy infestation, broadcast lightly over the ground. When used in this manner the bait will give exceptional results but may have to be renewed more often.

*If slugs and snails invading from adjacent properties are a menace, a pencil-thin line of Corry's Slug and Snail Death the entire length of the infested area will form a barrier and result in excellent control.

For best results be sure area to be baited is damp. In very dry weather water well before baiting.

Bait should be renewed at least every two weeks—more often during heavy infestations.

Dead slugs and snails should be removed and destroyed.

WARNING!

Avoid contact with skin, eyes, and clothing. Harmful dust. Avoid breathing dust. May be fatal if swallowed. Avoid contamination of food, food-stuffs, water supplies, and ponds. Do not apply bait directly on fruits, vegetables or field crops — use only on ground around these plants. Bait may be attractive to dogs. Keep dogs off treated areas. Destroy container when empty. Wash hands after use.

This product may be toxic to wildlife. Birds feeding on treated areas may be killed.

CORRY'S SLUG & SNAIL DEATH



from their hiding places during cold weather or during exceedingly dry or hot spells.

Slugs and snails hatch from eggs, laid in clusters, under debris, stones, or beneath the surface of the soil. Generally they will hatch out after 20 to 30 days of favorable weather conditions after which time they will continue to grow slowly, eating our most tender and succulent vegetation.

While these small slugs and snails are growing and eating they migrate toward better feeding and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snail problem. However, with the use of CORRY'S SLUG and SNAIL DEATH and a consistent baiting program, these pests can be reduced to where they no longer present a major threat to our gardens.

MANUFACTURED BY
E. M. MATSON, JR. CO.
7808 8th AVE. SO.
SEATTLE, WASHINGTON 98108

Canadian Registration No. 997
P C P Act—Guarantee:
Effective for Purpose Claimed.
Metaldehyde 2%

EPA REG. No. 8119-1-AA

MADE IN USA

NET WT. 16 OZ. (1 LB.)

ACTIVE INGREDIENT, METALDEHYDE 2%. INERT INGREDIENTS 98%.
WARNING: KEEP OUT OF REACH OF CHILDREN. (SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.)

ACCEPTED
Jan 5, 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR LISTING TO POISON REGISTER-
ED UNDER NO. 8119-1

Comments

WARNI

Y'S

NAIL
H

ISH FORMULA

CORRY'S

SNAIL

TH

**CUT AND
LIFT**

Slugs and snails are two of our most destructive agricultural pests. Although their outward appearance is quite different they are essentially the same creature, except that the slug does not have the external shell of the snail.

Slugs and snails feed almost entirely at night, in the evening, or during dark days. They prefer moist, warm conditions and rarely venture from their hiding places during cold weather or during exceedingly dry or hot spells.

Slugs and snails hatch from eggs, laid in clusters, under debris, stones, or beneath the surface of the soil. Generally they will hatch out after 20 to 30 days of favorable weather conditions after which time they will continue to grow slowly, eating our most tender and succulent vegetation.

While these small slugs and snails are growing and eating they migrate toward better feeding and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snail problem. However, with the use of CORRY'S SLUG and SNAIL DEATH and a consistent baiting program, these pests can be reduced to where

Original ENGLISH FORMULA

CORRY'S

SLUG & SNAIL

DEATH



SNAIL TH



Generally they will hatch out after 20 to 30 days of favorable weather conditions after which time they will continue to grow slowly, eating our most tender and succulent vegetation.

While these small slugs and snails are growing and eating they migrate toward better feeding and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snail problem. However, with the use of CORRY'S SLUG and SNAIL DEATH and a consistent baiting program, these pests can be reduced to where they no longer present a major threat to our gardens.

MANUFACTURED BY
E. M. MATSON, JR. CO.
7808 8th AVE. SO.
SEATTLE, WASHINGTON 98108

Canadian Registration No. 997
P C P Act—Guarantee:
Effective for Purpose Claimed.
Metaldehyde. 2%

EPA REG. No. 8119-1-AA

MADE IN USA

NET WT. 16 OZ. (1 LB.)

INERT INGREDIENTS 98%.
DREN. (SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.)

Comments

SLUG & SNAIL DEATH



WARNING: ACTIVE INGREDIENT, METALDEHYDE 2%. INERT INGREDIENTS 98%.
KEEP OUT OF REACH OF CHILDREN. (SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.)

DEATH
SLUG & SNAIL
CORRY'S

ACCEPTED WITH COMMENTS

DIRECTIONS

GALCO BEEF CATTLE OIL is ready to use on beef cattle. Do not mix or dilute with oil, kerosene or distillate.

Soak backrubber roll until it is thoroughly wet but not dripping. Repeat as necessary to keep roll wet, usually at two week intervals. If oiler has manually operated valve adjust so that backrubber is kept saturated. Regular use will give good results against lice and will help reduce weight loss due to hornfly attacks.

CAUTION: Do not treat or allow calves under 3 months of age access to back rubbers.

GALCO BEEF CATTLE OIL

READY TO USE FOR BEEF CATTLE

U.S.D.A. Registration No. 8152-4

ACTIVE INGREDIENTS	100.00%
Toxaphene *	5.00%
Mineral Oil	95.00%

*Technical Chlorinated Camphene; Chlorine content
67-69%

Kills horn flies. Aids in the reduction of lice.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See right panel for additional cautions.

GALCO

Box "A" Sec. 3

Council Bluffs, Iowa 51501

CAUTION

Avoid using on small calves. Do not use on dairy cattle. Do not permit animal access to treatment within 28 days of slaughter. Avoid storage near feed or food products.

WARNING: May be fatal if swallowed. Avoid contact with eyes. Do not breathe vapors or get on skin or clothing. Wash thoroughly after using.

FISH AND WILDLIFE CAUTION: This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Apply this product only as specified on label.

CAUTION: Do not contaminate water by cleaning of equipment, or disposal of wastes.

ANTIDOTE: Internal -- Give tablespoon of salt in glass of warm water to induce vomiting. Repeat until vomit is clear. Call the doctor immediately.

NOTICE OF WARRANTY

GALCO warrants (its liability being limited to the purchase price of the product named on this label) that this product consists of the ingredients specified. Our recommendations for use of this product are based on tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. Buyer must assume all responsibility, including injury or damage, resulting from its use as such, or in combination with other materials.

Contents _____

