

1975 16
300

Mr. E.M. Matson, Jr.
E.M. Matson, Jr. Co.
7808 8th Avenue South
Seattle, WA 98108

AUG 15 1986

Dear Mr. Matson:

Subject: Revision of Precautionary Statement
Corry's Slug & Snail Death
EPA Registration No. 8119-1
Your Application Dated May 7, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

On the side panel of your label revise the sentence "Corry's Slug & Snail Death can be used to protect all fruits and vegetable plants" to read:

Corry's Slug & Snail Death can be used to protect all fruit and vegetable plants listed on this label.

This change must be made at your next printing of labeling for this product.

Sincerely yours,

Lawrence J. Schnaubelt ^{LJS}
Acting Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

90026:Schnaubelt:E-11;KENCO:8/5/86;8/15/86;SONJA:VO

CONCURRENCES

SYMBOL ▶							
SURNAME ▶							
DATE ▶							

CUT AND
LIFT

Original ENGLISH FORMULA

NOT AFFECTED BY RAIN



**IDEAL
FOR
VEGETABLE
GARDENS**

Slugs and snails are two of our most destructive agricultural pests. Although their outward appearance is quite different they are essentially the same creature, except that the slug does not have the external shell of the snail.

Slugs and snails feed almost entirely at night, in the evening, or during dark days. They prefer moist, warm conditions and rarely venture from their hiding places during cold weather or during exceedingly dry or hot spells.

Slugs and snails hatch from eggs, laid in clusters, under debris, stones, or beneath the surface of the soil. Generally they will hatch out after 20 to 30 days of favorable weather conditions after which time they will continue to grow slowly, eating our most tender and succulent vegetation.

While these small slugs and snails are growing and eating they migrate toward better feeding and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snail problem. However, with the use of CORRY'S SLUG & SNAIL DEATH and a consistent baiting program, these pests can be reduced to where they no longer present a major threat to our gardens.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store in a cool, dry place out of direct sunlight, in original container, and not accessible to children or pets. Avoid contamination of feed and foodstuffs. Do not reuse empty carton. Wrap carton and put in trash.

MANUFACTURED BY

E. M. MATSON, JR. CO.

7808 - 8th AVE. SO.

SEATTLE, WASHINGTON 98108

EPA REG. NO. 8119-1-AA

EPA EST. NO. 8119-WA-1

MADE IN U.S.A.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

AUG 15 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

8119-1

**DEATH
SLUG & SNAIL
CORRY'S**



GENERAL APPLICATIONS

CORRY'S SLUG & SNAIL DEATH is so easy to use. Requires no mixing, spraying, or special applicators. Just scatter lightly on the soil surface in infested areas. Apply heavier if infestation is severe. This bait is extremely attractive to slugs and snails and actually lures them from their hiding places and kills them. Reapply at least every two weeks — more often during heavy infestations or in areas heavily watered. All likely places should be treated as these pests travel. They favor damp places around all vegetable plants such as beans, tomatoes, lettuce, celery and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra lawns, citrus groves, ivy patches, and other ground cover where they obtain shelter by day. Use approximately one pound per 1750 square feet.

*SPECIAL APPLICATIONS

If slugs and snails invading from adjacent properties are a menace, a pencil-thin line of CORRY'S SLUG & SNAIL DEATH the entire length of the infested area will form a barrier and result in excellent control.

CORRY'S SLUG & SNAIL DEATH can be used to protect all fruit and vegetable plants. Apply only to soil surface. Do not apply to foliage or edible parts of the crop.

For best results be sure area to be baited is damp. In very dry weather, water well before baiting.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after use.

IMPORTANT: Keep away from children, dogs and pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

STATEMENT OF PRACTICAL TREATMENT
If swallowed: call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

This product may be toxic to birds and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water.

Origin



SLUG

IDEAL
FOR
VEGETABLE
GARDENS

