

READY TO USE FORMULA




ACCEL
 JAN 6 1987
 Under the
 Example
 for women
 1984

SLUG & SNAIL CONTROL

WARNING: Keep out of reach of children. This pesticide may be fatal to children, and dogs or other pets if eaten. Keep children and pets out of treated area. Follow directions and use with care.

Active Ingredient: Metaldehyde 4%
 Inert Ingredients: 96%

NET CONTENTS 1 QT. (32 FL. OZ.)



Carry's Liquid Slug & Snail Control is a new, convenient, ready-to-use liquid bait. It is extremely effective in attracting and killing both large and small slugs and snails. Easy to use, quick, clean and economical Carry's Liquid Slug & Snail Control resists rain and sprinkling and will remain effective in the garden for many days.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

TO APPLY: Raise spout and squeeze drops of bait 4-6 inches apart on soil in infested areas. All likely places should be treated as these pests travel. Slugs and snails favor damp places around vegetables, berries, ornamental plants, dichondra lawns, rockeries, greenhouses, compost heaps, ivy patches and other ground covers where they obtain shelter by day.

Normally apply bait every two weeks until infestation is under control. Heavy or prolonged rains will usually increase infestation and shorten effective period of bait. Use more frequently under these conditions. Use approximately one quart per 6160 square feet.

Slugs and snails feed almost entirely at night, in the evening, or during dark days. They prefer moist, warm conditions and rarely are active during exceedingly cold, hot or dry weather. Consider above when baiting. Preferred baiting time is in the evening and after warm rains.

***SPECIAL SITUATIONS:** *If slugs and snails are invading from adjacent properties squeeze line of drops of Carry's Liquid Slug & Snail Control along perimeter for excellent control.

*Carry's Liquid Slug & Snail Control can be used to protect the following fruit and vegetable plants. Apply to soil around plants and between rows. Do not apply to foliage or edible parts of crop. Protect asparagus, beans, cabbage, carrots, celery, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, radishes, spinach, squash, tomatoes, turnips, apples, avocados, blackberries, cherries, citrus, grapes, peaches, plums, and strawberries.

TIPS: When dispensing bait, occasionally tip bottle upright to be refilled with air. When bottle is almost empty add three tablespoons of water, shake vigorously, then continue to apply.

STORAGE: Store in original bottle, out of direct sunlight, in a cool, locked location. Avoid contamination of feed and foodstuffs.

DISPOSAL: Partially used bottle may be disposed of by securely wrapping in several layers of newspaper and discard in trash. Do not reuse empty bottle. Wrap bottle and put in trash.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: May be fatal if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes or clothing. Wash thoroughly after use. Do not allow children or pets to contact treated areas until they are dry. Do not apply to foliage or other plant parts.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. If on skin: Wash thoroughly with soap and water. If in eyes: Flush with plenty of water.

ENVIRONMENTAL HAZARDS

This product is toxic to birds and wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes.

Mfg. By: E. M. Matson, Jr. Company
7808 - 8th Avenue South
Seattle, Washington 98108

EPA Reg. No.
EPA Est. No.
Made in USA

