

Original ENGLISH FORMULA
NOT AFFECTED BY RAIN

CUT AND
LIFT

CORRY'S SLUG & SNAIL DEATH

SEE
ENCLOSURE

BLE
NS



DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

GENERAL APPLICATIONS
CORRY'S SLUG & SNAIL DEATH is so easy to use. Requires no mixing, spraying, or special applicators. Just scatter lightly on the soil surface in infested areas. Apply heavier if infestation is severe. This bait is extremely attractive to slugs and snails and actually lures them from their hiding places and kills them. Reapply at least every two weeks -- more often during heavy infestations or in areas heavily watered. All likely places should be treated as these pests travel. They favor damp places around vegetable plants, flower gardens, rockeries, hedges, dichondra lawns, ivy patches, and other ground cover where they obtain shelter by day. For best results, be sure area to be baited is damp. In dry weather, water well before baiting. Use approximately one pound per 1750 square feet.

SPECIAL APPLICATIONS
If slugs and snails invading from adjacent properties are a menace, a pencil-thin line of CORRY'S SLUG & SNAIL DEATH the entire length of the infested area will form a barrier and result in excellent control.

CORRY'S SLUG & SNAIL DEATH can be used to protect the following fruit and vegetable plants. Apply only to soil surface. Do not apply to the foliage or edible parts of the crop. Protect asparagus, beans, cabbage, carrot, celery, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, radishes, spinach, squash, tomatoes, turnips, apples, avocados, blackberries, cherries, citrus, grapes, peaches, plums, and strawberries.

STORAGE AND DISPOSAL
Store in cool, dry, locked location out of direct sunlight in original container. Do not allow access to children. Avoid contamination of feed and water. Do not reuse container. Discard cards in trash. Store in original container in several layers of newspaper and discard in trash.

Manufactured By
E. M. Watson, Jr. Co., Inc.
7808 - 8th Ave. South
Seattle, Washington 98108
EPA Reg. No. 8119-1-AA
EPA Est. No. 8119-WA-1
© 1992 E. M. Watson, Jr. Co., Inc.
Made in U.S.A.

KEEP OUT OF REACH OF CHILDREN
WARNING: THIS PESTICIDE MAY BE FATAL TO DOGS OR CATS IF EATEN. KEEP PETS OUT OF TREATED AREA. SEE ADDITIONAL PRECAUTIONARY STATEMENTS.
NET WT. 10 LBS.
Active Ingredient
Metaldehyde (2, 4, 6, 8-tetramethyl-1, 3, 5, 7-tetraoxacyclo-octane) 2%
Inert Ingredients 98%

JUL 27 1992

NOT AVAILABLE COPY

365 10201

Original ENGLISH FORMULA
NOT AFFECTED BY RAIN

CORRY'S

SLUG & SNAIL

DEATH

**IDEAL
FOR
VEGETABLE
GARDENS**



Slugs and snails are two of our most destructive agricultural pests. Although their appearance is quite different they are essentially the same creature, except that the snail does not have the external shell.

Slugs and snails feed almost entirely at night, in the evening, or during dark days. They prefer moist, warm conditions and crawl out from their hiding places during cool, damp weather or during exceedingly dry or hot weather.

Slugs and snails hatch from eggs in small clusters, under debris, stones, or other objects on the surface of the soil. Generally they hatch out after 20 to 30 days of favorable conditions after which time they begin to grow slowly, eating our most succulent vegetation.

While these small slugs and snails are feeding and eating they migrate toward better and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snail. However, with the use of CORRY'S SLUG & SNAIL DEATH and a consistent program, these pests can be reduced to such a degree they no longer present a major problem in gardens.

**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after use. Keep out of reach of children and domestic animals. This pesticide is toxic to dogs and other pets if eaten. Keep out of treated area.

IMPORTANT: Keep away from children and pets. Bait may be attractive to domestic animals. Do not apply to lawns or fine pets during application to avoid injury. Do not believe they are being treated. Store under conditions where there is no hazard.

STATEMENT OF PRACTICAL TREATMENT:
If swallowed: Call a physician or poison center. Drink 1 or 2 glasses of water to induce vomiting by touching back of tongue with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Wash thoroughly with soap and water. Flush with plenty of water.

ENVIRONMENTAL HAZARD:
Do not apply directly to water, swamps, bogs, marshes, and other areas that will not contaminate water when used in equipment washwaters.

KEEP OUT OF REACH OF CHILDREN

CAUTION: THIS PESTICIDE MAY BE FATAL TO DOGS ON OTHER PETS IF EATEN. KEEP PETS OUT OF TREATED AREA. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET WT.	9.1 LB.
Active Ingredient	
Metaldehyde (2, 4, 6, 8-tetramethyl-1, 3, 5, 7-tetraoxycyclo-octane)	2%
Inert Ingredients	98%



Additional STORAGE AND DISPOSAL statement for plastic bucket as per PR NOTICE 83-3 AND 84-1:

STORAGE: Store in cool, locked location out of direct sunlight, in original container, and not accessible to children or pets. Avoid contamination of feed and foodstuffs.

DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bucket. Rinse thoroughly before discarding in trash.

