

SLUGZSNAIL DEATH



KEEP OUT OF REACH OF CHILDREN

CAUTION: THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN. KEEP PETS OUT OF TREATED AREA. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET WT. 2 LBS. 4 OZ. (1.02 KG)

Active Ingredient Metaldehyde (2, 4, 6, 8-tetramethyl-1, 3, 5, 7-tetraoxycyclo-octane) Inert Ingredients

Slugs and snails are two of our most destructive garden pests. Although their outward appearance is quite different they are essentially the same creature, except that the sluo does not have the external shell of the snail.

Slugs and snails feed almost entirely at night, in the evening, or during dark days. They prefer moist, warm conditions and rarely venture from their hiding places during cold weather or during exceedingly dry or hot spells.

Slugs and snails hatch from eggs, laid in clusters, under debris, stones, or beneath the surface of the soil. Generally they will hatch out after 20 to 30 days of favorable weather conditions after which time they will continue to grow slowly, eating our most tender and succulent vegetation.

While these small slugs and snails are growing and eating they migrate toward better feeding and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snat problem. However, with the use of CORRY'S SLUG & SNAIL DEATH and a consistent baiting program, these pests can be reduced to where they no longer present a major threat to our gardens.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after use. Keep out of reach of children and domestic animals. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area.

IMPORTANT: Keep away from children, doos and pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

STATEMENT OF PRACTICAL TREATMENT If swallowed: Do not induce vomiting. Seek medical assistance immediately. If on skin: Wash, with plenty of spap and water. Get medical attention if irritation persists. If in eyes: Flush with plenty of water. Call a physician a irritation persists. It inhaled: Remove from forther exposure and treat symptomatically.

ENVIRUNMENTAL HAZARDS *

For terrestrial dises, do not apply directly to water, or to areas where surface-water is present or to intertidal areas below the mean high water mask Do not contaminate water when disposing of equipment washwaters.



CAUTION: THUS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN. KEEP PETS OUT OF TREATED AREA. SEE

SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

CUT AND LIFT

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL APPLICATIONS

CORRY'S SLUG AND SNAIL DEATH is so easy to use. Requires no mixing, spraying, or special applicators. Just scatter lightly on the soil surface in infested areas. Apply heavier if infestation is severe. This product is for use in home and garden areas. This bait is extremely attractive to slugs and snails and actually lures them from their hiding places and kills them. Reapply at least every two weeks - mr during heavy infestations or in areas heavily watered. All likely places impuld be treated as these pests travel. They favor damp places around vegetable plants, flower gardens, rockeries, hedges, dichondra lawns, ivy patches, and other ground cover where they obtain shelter by day. For best results be sure area to be baited is damp. In very dry weather, water well before baiting. Use approximately one pound per 1750 square feet.

*SPECIAL APPLICATIONS

*If slugs and snalls invading from adjacent properties are a menace, a pencil-thin line of CORRY'S SLUG & SNAIL DEATH the entire length of the infested area will form a barrier and result in excellent control.

*CORRY'S SLUG AND SNAIL DEATH can be used to protect the following fruit and vegetable plants. Apply only to soil surface. Do not to the foliage or edible parts of the Protect asparagus, beans, cabbage, carrots, celery, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, radishes, spinach, squash, tomatoes, turnips, apples, avocados, blackberries, cherries, citrus, grapes, peaches, plums, and strawberries.

STORAGE AND DISPOSAL

Store in cool, dry, locked location out of direct sunlight, in original container, and not accessible to children or pets. Avoid contamination of feed and foodstuffs. Do not reuse carton. Discard carton in trash. Securely wrap original container in several layers of newspaper and discard in trash.

Manufactured By E.M. Matson, Jr. Co., Inc. P.O. Box 1820 North Bend, WA. 98045 EPA Reg. No 8119-1 EPA Est. No. 8119-WA-1 © 1998 E.M. Matson, Jr. Co., Inc. Made in U.S.A.

NET WT. 2 LBS. 4 OZ. (1.02 KG)

Active Ingredient Metaldehyde (2, 4, 6, 8-tetramethyl-

1, 3, 5, 7-tetraoxycyclo-octane) Inert Ingredients 98%