

READY TO USE FORMULA

Ideal for Vegetable Gardens



SLUG & SNAIL CONTROL

Keep out of reach of children.

CAUTION: This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area. See back panel for additional precautionary statements.

Active Ingredient: Metaldehyde (2, 4, 6, 8-tetra-methyl-1, 3, 5, 7-tetraoxycyclo-octane) 4%
 Other Ingredients 96%

NET CONTENTS 32 FL. OZ. (946ml.)

3/2002

A C C E P T E D

MAR 20 2003

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 Registered under
 P. Reg. No 8119-2

Corry's Liquid Slug & Snail Control is a convenient, ready-to-use liquid bait. It is extremely effective in attracting and killing both large and small slugs and snails. Easy to use, quick, clean and economical **Corry's Liquid Slug & Snail Control** resists rain and sprinkling and will remain effective in the garden for many days.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO APPLY: Raise spout and squeeze a solid line of this product on the soil in infested areas. This product is for use in home and garden areas. All likely places should be treated as these pests travel. Slugs and snails favor damp places around vegetables, berries, home ornamental plants, dahlia, lawns, rockeries, home greenhouses, compost heaps, ivy patches and other ground covers where they obtain shelter by day.

Normally apply bait every two weeks until infestation is under control. Heavy or prolonged rains will usually increase infestation and shorten effective period of bait. Use more frequently under these conditions.

Slugs and snails feed almost entirely at night, in the evening, or during dark days. They prefer moist, warm conditions and rarely are active during exceedingly cold, hot or dry weather. Consider above when baiting. Preferred baiting time is in the evening and after warm rains.

***SPECIAL SITUATIONS:** If slugs and snails are invading from adjacent properties. Squeeze a solid line of **Corry's Slug & Snail Control** along the perimeter for excellent control.

***Corry's Liquid Slug and Snail Control** can be used to protect the following fruit and vegetable plants. Apply to soil around plants and between rows. Do not apply to foliage or edible parts of plants. Protect asparagus, beans, cabbage, carrots, celery, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, radishes, spinach, squash, tomatoes, turnips, apples, avocados, blackberries, cherries, citrus, grapes, peaches, plums, and strawberries.

TIPS: When dispensing bait, occasionally tip bottle upright to be refilled with air. If spout becomes clogged, lift spout and clear with toothpick.

STORAGE: Store in original bottle, in a cool, locked location protected from direct sunlight and freezing temperatures. Avoid contamination of feed and foodstuffs.

DISPOSAL: IF EMPTY: Do not reuse this container. Place in trash or offer for recycling if available. IF PARTLY FILLED: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Wash thoroughly after use. Keep out of reach of children and domestic animals. Do not allow children or pets to contact treated areas. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area. Do not apply to foliage or other plant parts.

FIRST AID

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product, (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center (NPIC) at 1-800-858-7378

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Mfd. for **E.M. MATSON, JR. CO., INC.** © 3/2002
P. O. BOX 1820
NORTH BEND, WASHINGTON 98045

EPA Reg. No. 8119-2 **MADE IN CANADA**
EPA Est. No. 033518-CN-001 U.S. PATENT NO. 5,017,620
PRINTED IN THE U.S.A. 11/2002

