



IDEAL FOR VEGETABLE GARDENS

ACCEPTED

OCT 1 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 8119-3
EPA Reg. No. 8119-3

LIQUID BUG SNAIL & SLUG BAIT



READY TO USE FORMULA

Keep out of reach of children.


CAUTION: This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area. Read and follow all directions. See back panel for additional precautionary statements.

Active Ingredients:

Metaldehyde (2,4,6,8-tetra-methyl-1,3,5,7-tetraoxocyclo-octane).....	2%
Carbaryl (1-Naphthyl N-methylcarbamate)	5%
Other Ingredients	93%

Net Contents 24 fl. oz. (710ml.)

MADE IN CANADA



Corry's Liquid Bug, Snail & Slug Bait is a convenient, ready-to-use liquid bait. It is extremely effective in the control of slugs, snails, earwigs, sowbugs, cutworms, crickets, grasshoppers, ants, armyworms, millipedes, pillbugs, and darkling beetles. Easy to use, quick, clean, and economical **Corry's Liquid Bug, Snail & Slug Bait** resists rain and sprinkling and remains effective in the garden for many days.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow product to dry in treated areas before reentering. Do not repeat applications more than once every seven days. Do not apply product more than four times per season. Days from product application to harvest noted below in parenthesis under **SPECIAL SITUATIONS**.

TO APPLY: Raise spout and squeeze drops of bait 4-6 inches apart on soil in area to be treated. This product is for use in home and garden areas. This bait is extremely attractive to slugs, snails and other listed insects and will lure them from their hiding places. All likely places should be treated as slugs, snails, and other listed insects travel. They favor damp, well shaded locations where they can obtain shelter by day. Be sure area to be treated is damp! In dry weather, water area well before applying bait. Preferred baiting time is in the evening as these pests feed mainly at night. Normally apply bait every two weeks until infestation is under control. Heavy or prolonged rain can shorten effective period of bait. Use more frequently under these conditions. Use approximately 24 oz. per 4620 square feet.

***SPECIAL SITUATIONS:** *If slugs, snails and other listed insects are invading from adjacent properties, a pencil-thin line the entire length of the infested area will form a barrier and result in excellent control.

***Corry's Liquid Bug, Snail & Slug Bait** can be used to protect home ornamental plants, flower gardens, dichondra and home turf lawns, and fruit and vegetable plants. When protecting fruit and vegetable plants, treat area around plants and between rows. Do not apply to foliage or edible parts of plants. Protect the following fruit and vegetable plants: asparagus (1), beans (3), cabbage (3), carrots (7), celery (15), cucumbers (3), lettuce (15), melons (3), peas (3), peppers (3), potatoes (7), radishes (7), spinach (15), squash (3), tomatoes (3), turnips (7), and strawberries (7).

TIPS: When dispensing bait, occasionally tip bottle upright to be refilled with air. When bottle is almost empty, add three tablespoons of water, shake vigorously, then continue to apply. If spout becomes clogged, lift spout and clear with toothpick.

STORAGE: Store in original bottle, in a cool, locked location protected from direct sunlight and freezing temperatures. Avoid contamination of feed and foodstuffs.

DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bottle. Rinse thoroughly before discarding in trash.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes eye irritation. May be harmful if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Wash thoroughly after use. Keep out of reach of children and domestic animals. Do not allow children or pets to contact treated areas. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area. Do not apply to foliage or other plant parts.

USER SAFETY REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. Do not allow children or pets to contact treated area until product has dried.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Do not induce vomiting. Seek medical assistance immediately. If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists. If in eyes: Flush with plenty of water. Call a physician if irritation persists. If inhaled: Remove from further exposure and treat symptomatically. **NOTE TO PHYSICIANS:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Mfd. by: **E.M. MATSON, JR. CO., INC © 2000**
P. O. BOX 1820
NORTH BEND, WASHINGTON 98045
09/99 **PRINTED IN THE U.S.A.**



EPA Reg. No. 8119-3 EPA Est. No. 1791-CN-1
U.S. PATENT NO. 5,017,620
MADE IN CANADA

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378