## **BEST AVAILABLE COPY**

CORRYS

IDENTED BY VIEW BY STATE OF AND ENERGY









Panamoralesive Alearan a Socialina

### Keepouvoineren ordilliten

CAUMON Tribipesidoemey of italio dogs of other pasification (ealen Responsison to hireated area Resoland followall directors Seesido and followall directors

ACTIVE INGREDIENTS: Metaldenyde (2,4,6,8 tetramethyl=1),9,5,7,7 tetraoxycyclo-octane). (Carbary (11Naphthyl),1-methylcarbamate). (NERTINGREDIENTS)

SEVINGER FOR SERVING INCIDENCE OF FRANCE POR POR CONTRACTOR PROPERTY PROCESS.

Net Weight 2 lbs. (908g)

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

aline (ili

CAUTION: Causes eye irritation. May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or proceed wash thoroughly after use. Keep out control children and domestic animals. This proceed may be fatal to dogs or other pets if eaten. Keep pets out of treated area.

IMPORTANT: Keep away from children, dogs or pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

#### USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long parts
   shoes plus socks and household latex
   or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately handling and before eating, smoking the toilet.

  Using the toilet.

  Do not allow children or pets to contact.

treated area until dust has settled.

STATEMENT OF PRACTICAL TREATMENT
If swallowed: Call a physician or polson control
center. Drink 2 or 3 glasses of water and
induce vomiting by touching back of throat
with finger. Do not induce vomiting or give
anything by mouth to an unconscious person.
If on skin: Wash thoroughly with soap and
water. If In eyes: Flush with plenty of fresh
water. NOTE TO PHYSICIANS: Carbaryl is a
moderate, reversible cholinesterase inhibitor.
Alropine is antidotal

ENVIRONMENTAL HAZARDS
This product is extremely toxic to aqualic and estuarine invertabrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Mfd. by: E.M. Matson, Jr. Co., Inc. P.O. Box 1820 North Bend, WA 98045

EPA Reg. No. 8119-5 EPA Est. No. 8119-WA-1

NO 1997 E.M. Matson, Jr. Co., Inc.



IDAN FORVEGERNEUF GYRDANS











neribliékonésekönyegetki

CAUTION, This pesticide may be letted to does or other pat atem keeproatsoriikoriiteatta jajes. Reackaronollowalkelite airementasvasuoliivasjoineliojessioineavasiomenis

Net Weight 2 lbs: (908g)

BEST AVAILABLE COPY



CORRY'S SLUG, SNAIL & INSECT KILLER is a convenient, ready-to-use bait. It is extremely effective in the control of slugs, snails, earwigs, sowbugs, cutworms, crickets, grasshoppers, ants, armyworms, millipedes, pillbugs, and darkling beetles. Easy to use, quick and economical CORRY'S SLUG, SNAIL & INSECT KILLER resists rain and sprinkling and remains effective in the garden for many days.

#### **DIRECTIONS FOR USE**

in acety the

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Wearing household latex or rubber gloves, apply product directly from original container on soil In area to be treated. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. Do not repeat application more than once a week.

TO APPLY
Scatter bait lightly on soil in area to be treated. Apply heavier if infestation is severe. This product is for use in home and garden areas. This bait is extremely attractive to slugs, snails, and other listed insects and will lure them from their hiding places. All likely places should be treated as slugs, snails, and other listed insects travel. They favor damp, well shaded locations where they can obtain shelter by day. Be sure area to be treated is damp. In dry weather, water well before applying bait. Preferred baiting time is in the evening as these pests feed mainly at night. Normally apply balt every two weeks until infestation is under control. Heavy or prolonged rain can shorten effective period of bait. Use more frequently under these conditions. Use approximately one pound per 1750 square feet.

#### \*SPECIAL SITUATIONS

If slugs, snails, and other listed insects are invading from adjacent properties, a pencil-thin line the entire length of the infested area will form a barrier and result in excellent control.

\*CORRY'S SLUG, SNAIL & INSECT KILLER can be used to protect home ornamental plants, flower gardens, dichondra and home turf lawns, and fruit and vegetable plants. When protecting fruit and vegetable plants treat area around plants and between rows. Do not apply to foliage or edible parts of plant. Protect the following fruit and vegetable plants: asparagus, beans, cabbage, carrots, celery, cucumbers, lettuce, melons, peas, peppers, potatoes, radishes, spinach, squash, lomatoes, turnips, apples, blackberries, cherries, citrus, grapes, peaches, plums, and strawberries.

STORAGE: Store in cool, dry, locked location out of direct sunlight, in original container, and not accessible to children or pets. Avoid contamination of feed and foodstuffs.

DISPOSAL: Do not reuse carton. Discard carton in trash. Securely wrap original container in several layers of newspaper and discard in Irash.

🔄 insecticide.

ಾ ನವನೆ Eodenticiido Act ied, for the perticide