

CAUTIONS

This product will kill or seriously injure many desirable forms of vegetation. Do not apply directly to flowers, fruits, vegetables, grapes, ornamentals, cotton or other desirable plants. Do not use when there is hazard from drifting mists. (Coarse sprays are less likely to drift.)

Vapors from this product may injure susceptible plants in the immediate vicinity.

Avoid contamination of water used for domestic purposes and irrigation purposes. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not store near fertilizers, seeds, insecticides or fungicides.

Avoid contamination of feed and foodstuffs.

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place. Do not burn.

Spray equipment used in applying this product should be thoroughly cleaned before using for any other purpose. Use repeated flushing with soap and warm water or suitable chemical cleaner. It is best to use a separate sprayer for application of insecticides and fungicides.

May cause skin irritation.

Harmful if swallowed.

Avoid contact with eyes, skin or clothing.

Avoid inhaling spray mist.

Do not use on light sandy soils.

Do NOT store with foodstuffs.

NET CONTENTS _____

M C P P Selective Lawn Weed Killer

For Control of Broadleaf Weeds on Ornamental Turf Areas

ACTIVE INGREDIENT:

Diethanolamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid) 32.8%

INERT INGREDIENTS:

67.2%

TOTAL 100.0%

*Equivalent to 22.0% 2-(2-methyl-4-chlorophenoxy) propionic acid or 2.0 lbs. MCPP acid per gallon.

CAUTION: KEEP OUT OF REACH OF CHILDREN. See Other Precautions On Side Panels

GENERAL INFORMATION: MCPP has proven to be a very useful herbicide for control of the surface-creeping or prostrate broadleaf weeds listed below. Although relatively slow in action, one of the advantages of Riverdale Weeds MCPP is that it is less injurious to fine turf grasses, than 2,4-D or silvex. Can be used on bent varieties, but use caution on Seaside, Arlington and Congressional bents which may be injured. Do not use on St. Augustine Grass.

WEEDS CONTROLLED

Common Chickweed (<i>Stellaria media</i>)	Ground Ivy (<i>Glechoma hederacea</i>)
Mouse-ear chickweed (<i>Cerastium Vulgatum</i>)	Stitchwort (<i>Stellaria graminea</i>)
Red Clover (<i>Trifolium pratense</i>)	Knotweed (<i>Polygonum aviculare</i>)
White Clover (<i>Trifolium repens</i>)	

DIRECTIONS FOR USE

APPLICATION RATE ON:	PER 1,000 SQ. FT.	PER ACRE
Established greens and fine turf lawns	1 1/2-2 fl. ozs.	4-5 pints
Other established turf and fairways	2 1/2-3 fl. ozs.	6-7 pints

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

SOLD BY

FRANK MILLER & SONS INC.
13831 S. EMERALD AVENUE
CHICAGO, ILLINOIS 60627

DIRECTIONS Continued

Better weed control, especially of knotweed has been obtained with weekly applications at half recommended rate. This is preferred especially on bent grasses and aprons when there has been no previous experience.

HOW TO APPLY: Apply as a water-mixed spray in 20 to 40 gals. water per acre (4 gals. per 1,000 sq. ft.). Apply with boom type sprayer to give thorough coverage. Smaller areas may be treated with a knapsack sprayer. Do not mow for several days before or after application. Withhold sprinkling for 24 hours after application.

WHEN TO APPLY: For knotweed apply in spring or early summer when weeds are young for best results. For other weeds listed, apply in spring or fall when weeds are growing actively.

CAUTION: Do not apply when turf is suffering from drought conditions or during very hot or very wet weather or damage to turf may result.

Local conditions may affect the use of this chemical. Consult State Agricultural Experimental Station or Extension Service weed and turf specialists for specific recommendations as to dosage and time of application and for control of other weeds.

E.P.A. REG. NO. 8123-44

108-44