

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store pesticide in a locked area, away from access by children. Store in original container only. Keep container closed when not in use. Do not reuse empty container.

PESTICIDE DISPOSAL - Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL - Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. After a thorough pre-cleaning of the surface, apply solution with mop or cloth. Then wipe, mop or scrub so as to thoroughly wet the surface with the sanitizing solution. Sanitize with 1 ounce Concentrate to 4 gallons of water. Disinfectant with 3 oz. of Concentrate to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis.

HOSPITALS

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms. After a thorough pre-cleaning of the surface, apply a solution of the Concentrate at 3½ ounces per 5 gallons of water. Then wipe, mop or scrub so as to thoroughly wet the surface with sanitizing solution. At this level the Concentrate will be effective against Pseudomonas aeruginosa PRD-10.

FOOD PROCESSING EQUIPMENT

For sanitization of food processing equipment and food utensils, remove gross food particles and soil by a pre-flush or pre-scrape, and if necessary, a pre-soak. Thoroughly wash the item with a good detergent or compatible cleaner, followed by a potable rinse before application of the sanitizing solution. After thoroughly cleaning and rinsing equipment, dilute 1 ounce of Concentrate per 4 gallons of water to provide 200 ppm active quaternary. Then wipe or scrub with the sanitizing solution so as to wet the surfaces thoroughly making sure to wet all surfaces completely for at least 60 seconds. At this level, no potable water rinse is required. Prepare a fresh solution daily or more frequently if the solution becomes diluted or soiled.

NOT FOR SALE
In Accordance With

12-17-88

DISINFECTANT - SANITIZER-DEODORIZER CONCENTRATE

(Must Be Diluted Before Using)

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl	
ethylbenzyl ammonium chlorides	5%
INERT INGREDIENTS:	90%
	<hr/>
	100%

E.P.A. REG. NO. 8123-47
E.P.A. EST. NO. 8123-IL-1

NET VOLUME _____

DANGER

KEEP OUT OF REACH OF CHILDREN

See "First Aid" and other precautionary statements on left panel

Sold By

FRANK MILLER & SONS, INC
13831 S. Emerald Avenue
Chicago, Illinois 60627

DAIRIES

To sanitize dairy equipment, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 oz. of Concentrate to 4 gallons of water (200 ppm), by wiping or scrubbing the equipment with the solution so as to wet the surfaces thoroughly. At this level, no potable water rinse is required. Follow recommendations of local Health Board.

COLD DISINFECTANT:

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ oz. of concentrate per gallon of water.

RESTAURANT & BAR RINSE:

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce of concentrate to 4 gallons of water. No terminal potable water rinse required.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

DANGER: Keep Out of Reach of Children. Corrosive Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician immediately.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

SR-11 (8/88)