[8123-61]

[illegible]

WARRANT[illegible]

[illegible] the Direct[illegible] EXPRESS WARRANTIES [illegible] PRODUCT [illegible] SOLD AS [illegible] NO OTHER WARRANTIES [illegible] MADE, EXPRESS [illegible] IMPLIED.

DAMAGES: B[illegible] sole exclusive remedy [illegible] damages for breach [illegible] warranty, or negligence shall be limited to direct damages not exceeding the purchase price [illegible] and shall [illegible] incidental or consequential damages. [illegible] of any claim must be reported within [illegible] days of [illegible] buyer or user, following [illegible] such damage.

CAUSE FROM NON-CONTROLLABLE CONDITIONS: [illegible] and risks created or assumed by buyer [illegible] over which seller has no control, buyer and [illegible] of purchase or use, [illegible] for any release [illegible] seller from [illegible] handling or use of the product [illegible] distributors [illegible] or uses. No person is [illegible] any statement contained herein.

DIRECTIONS FOR USE

CROP SPRAY CONCENTRATE is designed for use indoors as a cleaning spray or as a preharvest spray where other materials cannot be used because of residue restrictions. When applied according to accepted use, cautions and directions, CROP SPRAY CONCENTRATE may also be used in combination with other insecticides where resistant insects are a problem, or as an "flexritter" where insects are flushed out [illegible] and [illegible] with spray residues. Observe [illegible] restrictions and limitations on the labeling of [illegible] used in mixtures.

Apply the listed amount per [illegible] repeat as required to maintain effective control. [illegible] sufficient water for thorough coverage is used. [illegible] unless otherwise noted, for listed crops [illegible] age of upper and lower leaf surfaces is essential for good control. When used alone, CROP SPRAY CONCENTRATE may be used up to and including day of harvest.

ASPARAGUS: Asparagus Beetle - 2 to 6 oz.

BEANS: Aphids, Leafhoppers, White Fly, Mexican Bean Beetle, 12-Spotted Cucumber Beetle [illegible] to 6 oz.

BROCCOLI, CABBAGE, BRUSSEL SPROUTS, CAULIFLOWER: Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth, [illegible] Flea Beetle, Harlequin Bug, Imported Cabbageworm [illegible] 2.

CELERY: Green Peach Aphid, [illegible] - 2 to 6 oz.

CRANBERRIES: Fireworms, Leafroller [illegible] oz.

EGGPLANTS: Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid - 2 to 6 oz.

LETTUCE: Cabbage Looper, Green Peach Aphid, [illegible] Imported Cabbageworm [illegible] 2.

PEPPERS: Green Peach Aphid - 2 to [illegible]

POTATOES: Aphids, Colorado Potato Beetle, Flea Beetle, Leafhopper [illegible] to 6 oz.

CROP SPRAY CONCENTRATE

ACTIVE INGREDIENTS	
Pyrethrins	[illegible]
Piperonyl Butoxide, Technical	50.00%
Petroleum Distillates	14.00%
INERT INGREDIENTS	20.00%
TOTAL	100.00%

*Equivalent to 48.[illegible]% Butylcarbityl 6,7-Propylpiperonyl Ether and 12.0% related compounds.

CAUTION

Keep out of reach of children

HARMFUL IF SWALLOWED
CAUTION -- DO NOT USE IN UNDILUTED FORM

This product is toxic to fish. Keep out of any body of water. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT REUSE EMPTY CONTAINERS. DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE OR RETURN TO DRUM RECONDITIONER.

E.P.A. REG. NO. 8123-61
E.P.A. Est. 8123-IL-1

NET CONTENTS _____

[illegible]

ANTS: [illegible] Ants, [illegible] Webworm, [illegible]

BEDBUGS: [illegible] Ants, [illegible] Flea Beetle.

MOSQUITOES: [illegible] per 4 gals. [illegible] per [illegible] water [illegible] per [illegible] gallon [illegible]

DROSOPHILA: [illegible] CROP SPRAY CONCENTRATE with 1206 parts water (1 pint per 150 gallons or 1 tablespoonful with 4 gallons water). Thoroughly wet the emulsion of the spray forms. Make treatments as follows:

1. Apply liberally to tomatoes and fruits in baskets, in trucks and in plants. Use sprayers at a high pressure for applying at the rate of five or six pints of the diluted spray to a two-ton load of tomatoes. Direct the spray for maximum coverage of the baskets or hampers. It is important to spray between and beneath the containers. This spray not only kills the flies, but the emulsion loosens any dead flies so that they are readily washed from the fruit.
2. Spray the raw stock stacked in the yard.
3. Dip baskets in the diluted spray after dumping the fruit to kill adhering larvae and pupae.
4. After washing and cleaning up the [illegible] in the processing plant and just prior to bringing fruit into it, the entire space inside the cannery should be sprayed at a dilution of 1 part CROP SPRAY CONCENTRATE to 59 parts water (1 pint with 7 gallons 3 pints water or 2 tablespoonfuls with 3-1/4 pints water). At the rate of 1 gallon to 750 square feet (1 quart to 187 sq. ft.) direct spray on walls, ceilings, and floors, paying special attention to forcing the spray into all cracks and crevices for the control of ants, roaches, silverfish, earwigs, spiders and cheese mites.

For use as a space spray in the processing plants, this dilution will give excellent control of flying insects, i.e. fruit flies, house flies, hornets, grain moths, gnats, mosquitoes and skipper flies. Apply at the rate of one-half ounce per 1000 cubic feet of space. Do not spray while the plant is in operation, as dead flies may fall into containers of the product being processed. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

SPACE SPRAY: To control flying insects such as fruit flies, hornets, wasps, grain moths, gnats, mosquitoes and skipper flies, dilute one part CROP SPRAY CONCENTRATE with 59 parts water (2 tablespoons with 3-1/4 pints water). Use at the rate of one-half ounce per 1000 cubic feet of space. Direct the space treatment upward and whenever practical keep doors and windows closed for at least ten minutes after application. Do not remain in treated areas. Ventilate the area after treatment is completed.

VAPORIZER APPLICATORS: For non-food processing areas, where oil residues are not objectionable, CROP SPRAY CONCENTRATE can be diluted in dendrol base oil instead of water, at the rate of 1-59, and applied by means of any good vaporizer type of application equipment. Control such insects as fruit flies, hornets, wasps, grain moths, mosquitoes and skipper flies.

2-17-[illegible]

8123-61