

BEST DOCUMENT AVAILABLE

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY

TERMS OF SALE OR USE: On purchase or use of this product, buyer and user agree to the following conditions: **WARRANTY:** This product is warranted, as of the time of sale, to (1) conform to the ingredient statement on the label, and (2) be reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THIS PRODUCT IS SOLD AS IS, NO OTHER WARRANTY IS MADE, EXPRESS, OR IMPLIED.**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage. **EXCUSE FROM NON-CONTROL-LABLE CONDITIONS:** Because of certain critical conditions created or incurred by buyer or user, or over which seller has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release seller from, liability arising out of the handling or use of this product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Obtain medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Do not expose to extreme temperatures. Do not stack more than four drums high. Drums should be opened in well-ventilated areas. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep container closed when not in use. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CROP SPRAY CONCENTRATE is designed for use alone as a clean-up spray or as a pre-harvest spray where other materials cannot be used because of residue restrictions. When applied according to accepted use, cautions and directions, **CROP SPRAY CONCENTRATE** may also be used in combination with other insecticides where resistant insects are a problem, or as an "exciter", where insects are flushed out of hiding and into contact with spray residues. Observe all cautions and limitations on the labeling of all products used in mixtures.

ACTIVE INGREDIENTS:

Pyrethrins	6.00%
Piperonyl Butoxide, Technical *	60.00%
Petroleum Distillates	14.00%
INERT INGREDIENTS:	20.00%

TOTAL: 100.00%

*Equivalent to 48.0% (Butylcarbityl) (6-Propylpiperonyl) Ether and 12.0% related compounds.

**CAUTION
KEEP OUT OF REACH OF CHILDREN**

**HARMFUL IF SWALLOWED
CAUTION - DO NOT USE IN UNDILUTED FORM**

EPA REG. NO. 8123-61
EPA EST. NO. 8123-IL-1

NET CONTENTS:

SOLD BY

FRANK MILLER & SONS, INC.
13831 S. Emerald Avenue
Chicago, Illinois 60627

ACC
AUG 10 1984
8123-61

Apply the listed amount per acre and repeat as required for effective control. Use in sufficient water for the (usually 100 gallons) unless otherwise noted, for list of upper and lower leaf surfaces is essential for good use alone, **CROP SPRAY CONCENTRATE** may be used including day of harvest.

- ASPARAGUS:** Asparagus Beetle - 2 to 6 oz.
- BEANS:** Aphids, Leafhoppers, White Fly, Mexican 12-Spotted Cucumber Beetle - 2 to 6 oz.
- BROCCOLI, CABBAGE, BRUSSEL SPROUTS,** Aphids, Cabbage Looper, Cross-striped Cabbageworm, Moth Larvae, Flea Beetle, Harlequin Bug, Imported Stink Bug - 2 to 6 oz.
- CELERY:** Green Peach Aphid, Cabbage Looper, Leaf Miner
- CRANBERRIES:** Fireworms, Leafhopper 2 to 6 oz.
- EGGPLANTS:** Blister Beetle, Colorado Potato Beetle, Green Peach Aphid - 2 to 6 oz.
- LETTUCE:** Cabbage Looper, Green Peach Aphid, D. Larvae, Imported Cabbageworm 2 to 6 oz.
- PEPPERS:** Green Peach Aphid - 2 to 6 oz.
- POTATOES:** Aphids, Colorado Potato Beetle, Flea hopper - 2 to 6 oz.
- RADISH:** Aphids, Flea Beetle - 2 to 6 oz.
- SPINACH:** Aphids, Cabbage Looper, Webworm - 2 to 6 oz.
- TOMATOES:** Green Peach Aphid, Colorado Potato Beetle, Flea Beetle - 2 to 6 oz.
- MOSQUITOES:** May be used on croplands as an adult agricultural workers during harvesting. Use at 1 oz. per acre when flown on by fixed wing aircraft.
- DROSOPHILA:** Fruit Fly Control: Dilute at the rate of 1 part **CROP SPRAY CONCENTRATE** with 1200 parts water (1 pint or 1 tablespoonful with 4 gallons water. Thoroughly in the spray tank. Make treatments as follows:
 1. Apply liberally to tomatoes and fruits in baskets, plants. Use sprayers at a high pressure for apply five or six pints of the diluted spray to a two-ton container. Direct the spray for a maximum coverage of the basket. It is important to spray between and beneath the fruit. Spray not only kills the flies, but the emulsion in the flies so that they are readily washed from the fruit.
 2. Spray the raw stock stacked in the yard.
 3. Dip baskets in the diluted spray after dumping, adhering larvae and pupae.
 4. After washing and cleaning up the inside of the processing plant just prior to bringing fruit into it, the entire processing plant should be sprayed at a dilution of 1 part **CROP SPRAY CONCENTRATE** to 59 parts water (1 pint with 1 gallon water or 2 tablespoonful with 3-1/4 pints water) 1 gallon to 750 square feet (1 quart to 187 sq. ft.) walls, ceilings, and floors paying special attention to spray into all cracks and crevices for the control of silverfish, crickets, spiders and cheese mites.
- For use as a space spray in the processing plants, give excellent control of flying insects, i.e. fruit flies, house flies, gnats, mosquitoes and skipper. The rate of one-half ounce per 1,000 cubic feet of space while the plant is in operation, as dead flies may fall on the product being processed. Foods should be removed during treatment. All food processing surfaces should be thoroughly cleaned before using. **SPACE SPRAY:** To control flying insects such as flies, wasps, grain moths, gnats, mosquitoes and skipper, use 1 part **CROP SPRAY CONCENTRATE** with 59 parts water with 3-1/4 pints water. Use at the rate of one-half ounce per 1,000 cubic feet of space. Direct the spray treatment upon practical keep doors and windows closed for at least 15 minutes. Do not return to treated areas. Ventilate treatment is completed.
- VAPORIZER APPLICATORS:** For non-food processing all residues are not undesirable, **CROP SPRAY CONCENTRATE** may be diluted in deodorized base oil instead of water, and applied by means of any good vaporizer type of applicator such insects as fruit flies, house flies, gnats, mosquitoes, and skipper flies.

BEST DOCUMENT AVAILABLE

PACKAGES ONLY

If this product, buyer
WARRANTY: This product
to the ingredient
for the purposes set
WARRANTED THIS
WARRANTY IS MADE,

damages for breach
direct damages not
include incidental or
it be reported within
which buyer or user
OM NON-CONTROL-
il conditions created
as no control, buyer
me responsibility for
e handling or use of
son is authorized to
d herein.

TS
ANIMALS

aid breathing vapors
In case of contact,
in medical attention

ly to water. Shrimp
ended on this label.
Do not apply where
her conditions favor
acter by cleaning of
ct only as specified

product in a manner

AL
storage or disposal.
ices. Do not stack
ed in well-ventilated
d in overpack drums
st or sand and dis-
sed when not in use.
use of this product
to disposal facility.
tainer. Triple rinse
filling, or puncture
procedures approved

alone as a clean-up
lets cannot be used
ording to accepted
ENTRATE may also
re resistant insects
are flushed out of
re all cautions and
mixtures.

CROP SPRAY CONCENTRATE

ACTIVE INGREDIENTS:	
Pyrethrins	5.00%
Piperonyl Butoxide, Technical *	60.00%
Petroleum Distillates	14.00%
INERT INGREDIENTS:	
	20.00%
TOTAL	
	100.00%

*Equivalent to 48.0% (Butylcarbityl) (6-Propylpiperonyl) Ether and 12.0% related compounds.

CAUTION
KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED
CAUTION - DO NOT USE IN UNDILUTED FORM

EPA REG. NO. 8123-61
EPA EST. NO. 8123-IL-1

NET CONTENTS:

SOLD BY

FRANK MILLER & SONS, INC.
13831 S. Emerald Avenue
Chicago, Illinois 60627

ACC
AUG 10 1984
8123-61

Apply the listed amount per acre and repeat as required for effective control. Use in sufficient water for thorough coverage (usually 100 gallons) unless otherwise noted, for listed crops. Coverage of upper and lower leaf surfaces is essential for good control. When used alone, CROP SPRAY CONCENTRATE may be used up to and including day of harvest.

- ASPARAGUS: Asparagus Beetle - 2 to 6 oz.
 - BEANS: Aphids, Leafhoppers, White Fly, Mexican Bean Beetle, 12-Spotted Cucumber Beetle - 2 to 6 oz.
 - BROCCOLI, CABBAGE, BRUSSEL SPROUTS, CAULIFLOWER: Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug - 2 to 6 oz.
 - CELERY: Green Peach Aphid, Cabbage Looper, Leaf Tier - 2 to 6 oz.
 - CRANBERRIES: Fireworms, Leafhopper - 2 to 6 oz.
 - EGGPLANTS: Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid - 2 to 6 oz.
 - LETTUCE: Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm - 2 to 6 oz.
 - PEPPERS: Green Peach Aphid - 2 to 6 oz.
 - POTATOES: Aphids, Colorado Potato Beetle, Flea Beetle, Leafhopper - 2 to 6 oz.
 - RADISH: Aphids, Flea Beetle - 2 to 6 oz.
 - SPINACH: Aphids, Cabbage Looper, Webworm - 2 to 6 oz.
 - TOMATOES: Green Peach Aphid, Colorado Potato Beetle, Stink Bug, Flea Beetle - 2 to 6 oz.
 - MOSQUITOES: May be used on croplands as an adulticide to protect agricultural workers during harvesting. Use at 1 oz. per 4 gals. water per acre when flown on by fixed wing aircraft.
 - DROSOPHILA: Fruit Fly Control: Dilute at the rate of 1 part CROP SPRAY CONCENTRATE with 1200 parts water (1 pint per 150 gallons or 1 tablespoonful with 4 gallons water). Thoroughly mix the emulsion in the spray tank. Make treatments as follows:
 1. Apply liberally to tomatoes and fruits in baskets, on trucks and in plants. Use sprayers at a high pressure for applying at the rate of five or six pints of the diluted spray to a two-ton load of tomatoes.
 1. Direct the spray for a maximum coverage of the baskets or hampers. It is important to spray between and beneath the containers. This spray not only kills the flies, but the emulsion loosens any dead flies so that they are readily washed from the fruit.
 2. Spray the raw stock stacked in the yard.
 3. Dip baskets in the diluted spray after dumping the fruit to kill adhering larvae and pupae.
 4. After washing and cleaning up the inside of the processing plant and just prior to bringing fruit into it, the entire space inside the cannery should be sprayed at a dilution of 1 part CROP SPRAY CONCENTRATE to 59 parts water (1 pint with 7 gallons 3 pints water or 2 tablespoonful with 3-1/4 pints water). At the rate of 1 gallon to 750 square feet (1 quart to 187 sq. ft.) direct spray on walls, ceilings, and floors paying special attention to forcing the spray into all cracks and crevices for the control of ants, roaches, silverfish, crickets, spiders and cheese mites.
- For use as a space spray in the processing plants, this dilution will give excellent control of flying insects, i.e. fruit flies, house flies, hornets, grain moths, gnats, mosquitoes and skipper flies. Apply at the rate of one-half ounce per 1,000 cubic feet of space. Do not spray while the plant is in operation, as dead flies may fall into containers or the product being processed. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleared before using. When using the product in these areas, apply only when the facility is not in operation.
- SPACE SPRAY: To control flying insects such as fruit flies, hornets, wasps, grain moths, gnats, mosquitoes and skipper flies, dilute one part CROP SPRAY CONCENTRATE with 59 parts water (2 tablespoons with 3-1/4 pints water). Use at the rate of one-half ounce per 1,000 cubic feet of space. Direct the spray treatment upward and whenever practical keep doors and windows closed for at least ten minutes after application. Do not spray in treated areas. Ventilate the area after treatment is completed.
- VAPORIZER APPLICATORS: For non-food processing areas, where oil residues are not undesirable, CROP SPRAY CONCENTRATE can be diluted in deodorized base oil instead of water, at the rate of 1-59, and applied by means of any good vaporizer type of applicator to control such insects as fruit flies, hornets, wasps, gnats, grain moths, mosquitoes, and skipper flies.