

ACCEPTED

JAN 24 1974

UNDER THE FEDERAL PESTICIDE
FUNCTIONAL AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 8123-47

COLD DISINFECTION:

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ oz. of concentrate per gallon of water.

RESTAURANT & BAR RINSE:

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce of concentrate to 4 gallons of water. No terminal: potable water rinse required.

DANGER: Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Call a physician immediately.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard it.

**DISINFECTANT -
SANITIZER-DEODORIZER
CONCENTRATE**

(Must Be Diluted Before Using)

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides 5%

n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
ethylbenzyl ammonium chlorides 5%

INERT INGREDIENTS: 90%
100%

E.P.A. REG. NO. 8123-47

NET VOLUME _____

DANGER

KEEP OUT OF REACH OF CHILDREN

See "First Aid" and other precautionary statements on left panel.

SOLD BY

Frank Miller & Sons, Inc.
13831 So. Emerald Avenue
Chicago, Illinois 60627

Cameraman's Note

Poor Copy



USE DIRECTIONS:

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. After a thorough precleaning of the surface, apply solution with mop or cloth. Then wipe, mop or scrub so as to thoroughly wet the surface with the sanitizing solution. Sanitize with 1 ounce Concentrate to 4 gallons of water. Disinfect with 3 oz. of Concentrate to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis.

HOSPITALS

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms. After a thorough pre-cleaning of the surface, apply a solution of the Concentrate at 3½ ounces per 5 gallons of water. Then wipe, mop or scrub so as to thoroughly wet the surface with the sanitizing solution. At this level the Concentrate will be effective against Pseudomonas aeruginosa PRD-10.

FOOD PROCESSING EQUIPMENT

For sanitization of food processing equipment and food utensils, after thoroughly cleaning and rinsing equipment, dilute 1 ounce of Concentrate per 4 gallons of water to provide 200 ppm of active quaternary. Then wipe or scrub with the sanitizing solution so as to wet the surfaces thoroughly. At this level, no potable water rinse is required.

DAIRIES

To sanitize dairy equipment, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 oz. of Concentrate to 4 gallons of water (200 ppm), by wiping or scrubbing the equipment with the solution so as to wet the surfaces thoroughly. At this level, no potable water rinse is required. Follow recommendations of local Health Board.