

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC
ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flames.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not expose to extreme temperatures. Do not stack more than four drums high. Drums should be opened in well-ventilated areas. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep container closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

INSECTICIDE FOGGING SPRAY

**DESIGNED FOR USE IN EDIBLE PRODUCT AREAS
OF OFFICIAL ESTABLISHMENTS OPERATING UNDER
THE MEAT, POULTRY, SHELL EGG GRADING AND
EGG PRODUCTS INSPECTION PROGRAMS AND
IN OTHER FOOD PROCESSING PLANTS**

**Apply with Equipment Designed and Adjusted
to Deliver a True Aerosol Mist**

ACTIVE INGREDIENTS:	
Allethrin (allyl homolog of Cinerin 1)	1.25%
Related compounds	0.14%
* Piperonyl Butoxide, Technical	5.00%
Petroleum Distillate	93.61%

100.00%

* Equivalent to 4.0% (butylcarbitol) (6-propylpiperonyl) ether and 1.0% related compounds

**CAUTION
KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed.

Avoid contamination of feed and foodstuffs.

See side panels for additional precautionary statements.

E.P.A. REG. NO. 8123-88
E.P.A. EST. NO. 8123-IL-1

NET CONTENTS: _____

SOLD BY:

FRANK MILLER & SONS, INC.
13831 S. Emerald Avenue
Chicago, Illinois 60627

SPACE SPRAY: To kill accessible stages of house flies, including fruit flies, small flying moths, gnats, cheese skippers, and those common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, meal worms and grain mites: Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 60% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind obstructions. Vacate and close off treated area. Do not reenter for at least 15 minutes after application. Ventilate before reoccupying.

In mushroom production and processing: To kill mushroom flies or fungus gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

SPACE AND CONTACT SPRAY: To kill accessible, exposed stages of silverfish, crickets, roaches, clover mites, earwigs, and ants, first using equipment which can deliver a jet spray, direct spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly off ant trails. Do not allow spray to contact stored food. Then shutting all doors, windows and shutting off all ventilating systems proceed with space spraying directions as given above. Use a dosage rate of 1 ounce per 1000 cubic feet of space. Vacate and close off treated area. Do not reenter for at least 15 minutes after application. Ventilate before reoccupying. Repeat treatment as necessary.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

Authorized for use in meat and poultry processing plants by the USDA Animal and Plant Health Inspection Program.

ULV-22 (6/84)

