[illegible] ADDITIONAL STATEMENT
HAZARDS TO HUMANS AND DOMESTIC [illegible]

WARNING

HARMFUL OR FATAL IF SWALLOWED [illegible]
[illegible] AVOID CONTACT WITH SKIN OR CLOTHING
MAY BE ABSORBED THROUGH SKIN [illegible]

[illegible]

[illegible]

[illegible] Wash eyes [illegible]
[illegible] eyes, get medical [illegible]

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

[illegible] of Federal law [illegible] inconsistent with its [illegible]

STORAGE AND DISPOSAL

Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Pesticide that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

METAL CONTAINERS: Triple rinse and offer for recycling, reconditioning, or disposal in an approved landfill or bury in a safe place.

PLASTIC CONTAINERS: Do not re-use empty containers. Triple rinse and incinerate or dispose of in an approved landfill or bury in a safe place.

IMPORTANT — Injury to or loss of desirable trees or other plants may result from failure to observe the following: Do not apply, or drain or flush equipment near desirable trees or other plants or on areas where the spray cut may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent spray drift to desirable plants. Do not contaminate irrigation water. Avoid contact with fertilizers, insecticides, fungicides and seeds. Thoroughly decontaminate the equipment from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens. Clean these parts separately.

GENERAL INFORMATION: This product is an effective [illegible] for use on non-crop [illegible] and many annual and perennial [illegible] [illegible] Effects [illegible] the application and the chemical [illegible] the root zone of the weeds [illegible] rate of uptake and duration of effect will vary [illegible] type, [illegible]

NOTICE OF WARRANTY

[illegible]

A [illegible]

SELLER MAKES NO WARRANTY [illegible]
[illegible] WARRANTIES [illegible]
[illegible] WARRANTY [illegible]

[illegible]

WEED KILLER

WATER [illegible]
[illegible]

[illegible]

[illegible]

KEEP OUT OF REACH OF CHILDREN

WARNING

See side panel for additional precautionary statements.

E.P.A. REG NO 8 [illegible]
E.P.A. EST NO 8[illegible]

NET CONTENTS

[illegible]

FRANK MILLER & SONS, INC.
13831 SOUTH MERAL
HIGHLAND, IL 62027

[illegible]

[illegible] the [illegible] results will be improved if the vegetation is present, results will be improved if the vegetation is removed before treatment. Do not apply when ground is frozen.

[illegible] the equipment to determine quantity [illegible] only cover measured area to be treated. [illegible] the proper amount of this product and mix into [illegible] water. After this product has been thoroughly mixed in the spray tank, agitation of the spray solution is not required.

Use a fixed-boom power sprayer properly calibrated to a [illegible] speed and rate of delivery. For application with a hand sprayer, use at least 200 gals. spray per acre. For treating small areas, a tank type hand sprayer or sprinkling can may be used. For basal (soil) treatment of brush, this product may be applied undiluted with an exact delivery hand gun applicator.

WEED CONTROL: To control most weeds for an extended period of time in non-cropland areas such as RAILROAD, HIGHWAY and PIPELINE RIGHT OF WAYS, PETROLEUM TANK FARMS, LUMBERYARDS, STORAGE AREAS, and INDUSTRIAL PLANT SITES:

Apply 12 to 24 gals. this product per acre to control ANNUAL WEEDS and GRASSES such as foxtail, ryegrass, wild oats, crabgrass, cheatgrass, bromegrass, ragweed, lambsquarters, purslane, [illegible], or hardgrass and turkey mullein. When applied just prior to or after emergence of annuals, rates as low as 6 gals. per acre control many annual weeds and grasses in low rainfall areas and give short term control in higher rainfall areas.

Apply 24 gals. this product per acre to control PERENNIAL WEEDS and GRASSES such as smooth brome, bahiagrass, bluegrass, redtop, purpletop, quackgrass, broomsedge, aster, dandelion, dog fennel, goldenrod, plantain and wild carrot. In areas with low or seasonal rainfall, rates as low as 18 gals. per acre control many perennial weeds and grasses.

Apply 46 to 96 gals. this product per acre to control JOHNSONGRASS, use at the same rate for OTHER HARD-TO-KILL PERENNIAL WEEDS and GRASSES such as bermudagrass, [illegible] grass, nutsedge, vaseygrass, saltgrass, bouncingbet, dogbane, bracken fern and horsetail. Where limited rainfall (usually less than 4 inches) occurs during the active growth period, such as some areas of the West, this product will not provide satisfactory control of hard-to-kill, deep rooted perennial weeds such as johnsongrass.

Use the higher levels of the dosage range on adsorptive soils (usually those high in organic matter or carbon).

Retreatment—Apply 6 to 24 gals. this product per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.

For Small Areas—24 fl. oz. of this product per 250 sq. ft. is approximately 3[illegible] gals. per acre.

BRUSH CONTROL: To control undesirable woody plants on rangelands and pastures such as RAILROAD RIGHT OF WAYS, STORAGE AREAS, INDUSTRIAL PLANT SITES and DRAINAGE DITCHES:

Apply as foliage or basal (soil) treatment, for use on [illegible] ditches, apply as a basal (soil) treatment only.

Note: For effective brush control and prevention of damage to the new vegetation, do not apply to brush standing in water. Do not water from treated ditches for irrigation purposes. Do not apply to ditches near or on right of ways or other areas where desirable timber or other desirable trees or shrubs are located that extend to the treated areas.

Broadcast Treatment—Apply 18 to 40 gals. per acre to control [illegible], sweet gum and pine; apply 40 to 96 gals. per acre to control or shrub species such as American elm, winged elm, hackberry, sumac and cottonwood. Use the higher levels of the dosage ranges on adsorptive soils (those high in organic matter or carbon).

Basal (Soil) Treatment—[illegible] woody plants such as cottonwood, hackberry, [illegible], [illegible], sweet gum, wild cherry, [illegible], American elm and sumac.

Undiluted—A[illegible] this product undiluted with an exact delivery hand gun applicator, this equipment delivers a thin stream [illegible] when triggered. Apply at the rate of 30 to 80 [illegible] per stem [illegible] to 4 [illegible] diameter [illegible] the base of the brush [illegible] When [illegible] stems with more than one [illegible] of solution is needed per stem, apply on the [illegible] to the stem. **Note**: This product is combustible [illegible] smoke [illegible] applying the [illegible] product [illegible] same.

[illegible]