

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS:

CAUTION

Avoid contact with skin, eyes or clothing.
 Avoid breathing spray mist.
 Wash thoroughly after handling.

ENVIRONMENTAL HAZARD:

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

USE PRECAUTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store more than four drums in high. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep container closed when not in use. Store at temperatures above 32° F. Do not store near seeds, fertilizers, insecticides, or fungicides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at your EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

This product is designed to control dandelions and many other broadleaf weeds susceptible to 2,4-D and MCPP, plus supply the necessary primary nutrients to help provide a healthy turf.

READ THE FOLLOWING DIRECTIONS CAREFULLY BEFORE USE

The individual ingredients in this formulation have proven to be very useful in the control of the broadleaf weeds, however, when combined, 2,4-D and MCPP will control more weeds more effectively, than if either ingredient was used separately. In some areas, Dichondra, bent grasses, or carpet buffalo may be injured. Do not use on St. Augustine or cent grass turf.

WEEDS CONTROLLED:

Ground Ivy	Buckhorn	Heavenly
Stitchwort	Curled Dock	Lamb's Quarter
Knotweed	Wild Garlic	Trifoliate
Clover	Wild Onion	Pigweed
Chickweed	Red Sorrel	Shepherd's Purse
Plantains	Ground Ivy	Purslane
Dandelion	Ragweed	Many others

This product will kill or seriously injure desirable forms of vegetation. Do not apply directly to or permit spray drift to susceptible plants as this product may injure flowers, fruits, vegetables, grapes, cotton, ornamentals, or other desirable plants.

FERTILIZER
10-6-4
(WITH BROADLEAF WEED KILLER)

ACTIVE INGREDIENTS: Herbicide

Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid	0.8%
Dimethylamine salt of 2-Methyl-4-chlorophenoxypropionic acid	1.6%
INERT INGREDIENTS:	98.20%
Equivalent of 2,4-Dichlorophenoxyacetic acid	0.06 lb.
2,4-Dichlorophenoxypropionic acid isomer specific by AOAC Method No. 6.001-51	0.06 lb.
and 0.66% 2-(2-methyl-4-chlorophenoxy)propionic acid, MCPP acid per gallon	0.06 lb.

Plant Food Analysis:

Total Nitrogen (N)	10%
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	4%

Potential Acidity 410 lbs. Calcium Carbonate Equivalent per ton.

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
STATEMENT OF PRACTICAL TREATMENT
 In case of contact, wash skin with soap and water. If in eyes, flush with water for 15 minutes and get medical attention.
 See Side Panels for Additional Precautionary Statements.

EPA REG. NO. 8123-10C
 EPA EST. NO. 8123-IL-1

NET CONTENTS _____

SOLD BY

FRANK MILLER & SONS, INC.
 13831 S. Emerald Avenue
 Chicago, Illinois 60627

To effectively control nuisance broadleaf weeds in turf, and to supply the required primary nutrients to the turf, dilute 2-3 gallons of this product in 3-5 gallons of water to treat 5000 sq. ft. For larger areas, dilute 15-20 gallons of this product in 30-50 gallons of water to treat one acre. Spray the turf area evenly, avoiding spray contact with all desirable broadleaf vegetation. Do not reseed for at least two weeks after application. Do not apply to a newly seeded lawn until after 3 mowings.

When to apply - For best results apply when weeds are young and actively growing. Do not water for at least 24 hours after application. If rain occurs during 24 hours after application, herbicidal action may be reduced, fertilizer action will not be effected. Delay mowing for 48 hours after use.

Where to apply - Use on lawns only. Do not use for gardens or on ground covers such as pachysandra or blue myrtle. Do not apply directly to or allow vapors to contact flowers, vegetables, fruits, shrubs, or trees.

NOTICE

Follow directions carefully. Timing and method of application, weather conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given here-with.

LF-22 (5'84)

BATCH NO. _____

RECEIVED
 JUN 27 1984
 EPA REG. NO. 8123-10C