

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

May be fatal if swallowed, inhaled, or absorbed through skin. Causes skin irritation. Do not get material on skin, eyes or clothing. Symptoms of injury may be delayed. Wear face shield, rubber gloves and rubber apron when handling concentrate. Do not breathe spray mist in order to avoid nasal, throat and respiratory tract irritation. When spraying or when contacting sprayed vegetation wet with spray, dew or rain, wear water-proof footwear and clothing. Do not contaminate feed or drinking water. Do not store or transport with feed or food.

ENVIRONMENTAL HAZARDS

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. Do not apply directly to water except as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

For containers triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

NON-CROP OR NON-PLANTED AREAS: To Kill Above Ground Weed Growth Around Buildings and Other Installations, Ornamental Gardens, Walkways, Paths, Beneath Greenhouse Benches, Fence Lines, Golf Courses, Parkways and Driveways for Control of Undesirable Broad-leaved and Grassy Weeds - 1 gal. per 20 gals. water (6 1/2 oz. per gal. water). Apply for full coverage and thorough weed contact. Apply preferably to young growth and up to seeding of weeds. Repeat applications necessary to maintain control, particularly on grasses and established weeds. Avoid spray contact with foliage or food crops or ornamental plants. Do not water treated areas for 24 to 36 hours after application.

AQUATIC USE DIRECTIONS: For application only to ponds, lakes, and drainage ditches where there is little or no outflow of water and which are totally under the control of the product's user. Do not use the treated water for animal consumption, swimming, spraying, irrigation, or domestic purposes for 14 days after treatment. Do not apply to aquatic sites in Florida.

Treatment of dense weed areas can result in oxygen loss from decomposition of dead weeds. This loss can cause fish suffocation; therefore, treat only 1/3 to 1/2 of the dense weed areas at a time and wait 14 days between treatments. DO NOT APPLY TO MUDDY WATER. Avoid creating muddy water during application. Do not apply under conditions of high wind and wave action. Do not apply where plants are covered with mud deposits. Consult your State Fish and Game Agencies before applying this product to public waters.

WEED SPECIES	GALS./SURFACE ACRE
SUBMERSED WEEDS (Infesting Still Ponds, Lakes, Ditches, or Portions thereof):	
Bladderwort (Utricularia spp.)	10 - 20
Coontail (Ceratophyllum demersum)	20
Elodea (Elodea spp.)	20
Naiad (Najas spp.)	10
Pondweeds (Potamogeton spp.)	20
*Except Richardson Pondweed (D. richardsonii) in Minnesota	
Watermilfoil (Hydrrophyllum spp.)	10 - 20

CONTACT VEGETATION CONTROLLER
FOR AREAS WHERE VEGETATION IS TO BE CONTROLLED WITHOUT SOIL STERILIZATION
NON-SELECTIVE - KILLS ON CONTACT
NON-RESIDUAL - INACTIVATED ON CONTACT WITH SOIL
DUAL-PURPOSE - FOR LAND & WATER VEGETATION

ACTIVE INGREDIENT:

Diquat dibromide [6,7-dihydrodipyrido (1,2-a:2',1'-c) pyrazinodinium dibromide] 4.35%

INERT INGREDIENTS: 95.65%

Contains 0.20 lbs. Diquat Cation per gal. or 0.373 lbs. Salt per gal. This product contains a herbicidal chemical that when used as directed will kill most all forms of plant growth without damage to the soil. Absorption and herbicidal action is usually quite rapid with effect visible in a few days. This product is inactivated on contact with soil.

Read entire label. Use strictly in accordance with label warnings and directions, and in conformity with Federal and State Regulations.

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENT OF PRACTICAL TREATMENT IF ON SKIN: Wash immediately with water, remove clothing and wash skin where necessary.

IF IN EYES: Wash thoroughly with water and get medical attention.

IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by placing fingers in back of throat. Never give anything by mouth to an unconscious person.

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

E.P.A. REG. NO. 8123-
E.P.A. EST. NO. 8123-IL-1

NET CONTENTS:

SOLD BY:

DIRECTIONS CONTINUED

APPLICATION: In early season where submersed weed growth has not reached the surface apply CONTACT VEGETATION CONTROLLER by pouring directly from the container into the water while moving slowly over the water surface in a boat. Distribute evenly over infested areas in strips 40 feet apart. In mixed weed populations, use the high rate of application as indicated by weeds present.

In late season or where submersed weed growth has reached the water surface throughout the area to be treated apply CONTACT VEGETATION CONTROLLER by pouring as above or injecting a diluted (10 - 20 to 1) solution below the water surface in strips no more than 20 feet apart. Use the high rate of application as indicated by the weeds present. Where algae are present along with the submersed weeds, pretreatment with Copper Sulfate at recommended rates is advised for best results with CONTACT VEGETATION CONTROLLER.

FLOATING WEEDS	GALS./SURFACE ACRE
Pennywort (Hydrocotyle umbellata)	5 - 7 1/2
Selvinia (Selvinia selindfolia)	5 - 7 1/2
Waterhyacinth (Eichhornia crassipes)	5 - 7 1/2
Waterlotus (Najas stratiotes)	5 - 7 1/2

APPLICATION: 150 to 200 gallons of water plus 1 pint non-ionic surfactant should be applied as an overall spray for control of the above weeds.

The high rates of CONTACT VEGETATION CONTROLLER should be used for late season application.

Duckweed (Lemna spp.) 10

APPLICATION: For control apply CONTACT VEGETATION CONTROLLER as an all-over spray in 50 to 150 gallons of water per acre. Care should be taken to thoroughly cover all plants on water and on damp marginal areas. Reinfestation of duckweed occurs readily from untreated areas. Retreatment with CONTACT VEGETATION CONTROLLER may be necessary to obtain season long control.

WEED SPECIES GALS./100 GALS. WATER
MARGINAL WEEDS (Infesting the Edges and Other Portions of Ponds, Lakes and Ditches)

Cattails (Typha spp.) 10

APPLICATION: For top kill, apply CONTACT VEGETATION CONTROLLER in 100 gallons water plus 1 pint non-ionic surfactant for full coverage and thorough weed control. Repeat treatment as necessary to control re-growth. For best results apply before flowering.

ALGAE: CONTACT VEGETATION CONTROLLER will control certain filamentous algae: Spirogyra spp. and Pithophora spp. Apply 0.5 - 1.5 ppm. w. CONTACT VEGETATION CONTROLLER for control. Application methods as for submersed weeds.

Formula For Parts Per Million (PPM) Recommendations
Gallons to use - p.p.m. recommended x acreage of water x average depth of water x 14.
For Example - 1/2 p.p.m. recommendations; two acre pond; 4 feet average depth; 1/2 x 2 x 4 x 14 equal 56 gallons required.

CONDITIONS OF SALE:

1. Seller warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Seller neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks or liability (except those assumed by Seller under 1 above) resulting from handling, storage and use of this material.

ONLY FOR SALE TO, USE, AND STORAGE BY SERVICEPERSONS.

8103-102

- 2 -

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Richard F. Mountfort *RFM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosures