

10F1
Pm 25
8/23-201

FERTILIZER 15-5-7

WITH CRABGRASS PREVENTER

ACTIVE INGREDIENT:	
N-butyl-N-ethyl-s,s,e-trifluoro	00.78%
2,6-dinitro-p-toluidine	99.22%
INERT INGREDIENTS:	
TOTAL:	100.00%

Contains 9.78 lb. benflin per 100 lbs.

GUARANTEED ANALYSIS

Total Nitrogen (N)	15.00%
15.0% Ammoniacal Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	5.00%
Soluble Potash (K ₂ O)	7.00%

Sources of Plant Nutrients: Urea, Ammonium Phosphate, Muriate of Potash

EPA REG. NO. 8123-
EPA EST. NO. 8123-IL-1

NET WEIGHT: _____

BATCH NO. _____

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water, for eyes flush with water for 15 minutes and get medical attention.

STORE IN A DRY PLACE

Sold By
FRANK MILLER & SONS, INC.
13821 S. Emerald Avenue
Chicago, Illinois 60627

ACCEPTED

AUG - 4 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 8123-201

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Do not contaminate feed or feedstuffs. Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this dust only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling.

STORAGE AND DISPOSAL

Do not contaminate water, feed, or food by storage or disposal.
STORAGE: Do not stack more than four containers high. Leaking or damaged containers should be placed in overpack drums for disposal. Spills should be collected and disposed of in a sanitary landfill. Keep container closed when not in use. Store in a dry place. Do not store near pesticides or seeds.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

Use this product only for recommended purposes and at recommended dosage.

This product contains a quality fertilizer and a selective pre-emergence herbicide which controls annual bluegrass (Poa annua), smooth and hairy crabgrass, green and yellow foxtail, barnyardgrass and goosegrass (silver crabgrass or crow-foot) by killing plants as seeds germinate. It does not control established weeds.

Use only on established turfs listed below which are tolerant to the herbicide in this product when applied as directed. See PRECAUTIONS.

- | | |
|-----------------------------|--------------------|
| Bentgrass (perennial) | Bermudagrass |
| Ryegrass (perennial) | Centipedegrass |
| Fescue (except fine-leaved) | St. Augustinegrass |
| Bahiagrass | Zoysiagrass |

WHEN TO APPLY: Apply to established turf before the germination of annual weeds listed above. The general times of application are covered below. The precise time of application for a specific area will vary significantly because of climate and individual cultural practices. Consult State Agricultural Experiment Station Weed Specialists or Local Turf Authorities for any additional information regarding specific weed germination dates in your area.

SOUTHERN (warm season) GRASSES:

- For pre-emergence control of annual bluegrass (Poa annua), apply in late summer to early fall, prior to annual bluegrass seed germination. Additional applications at the same rate can be made 4 to 6 months later in early spring, prior to soil warming and subsequent germination. In areas heavily infested with annual bluegrass, its control will temporarily result in thin turf areas.
- The fertilizer in this product plus proper irrigation will encourage the existing turfgrass to fill in the thin spots previously occupied by the annual bluegrass.
- For pre-emergence control of crabgrass, foxtail, barnyardgrass, and goosegrass, apply in late winter or early spring prior to annual weed grass germination. A second application may be made 4 to 6 months after initial treatment for continued control of annual weeds which may germinate into the summer or early fall.

AMOUNT TO APPLY: Use 8.8 lbs. per 1,000 sq. ft. or 384 lbs. per acre (equivalent to 3 lbs. of active ingredient per acre).

NORTHERN (cool season) GRASSES:

- For pre-emergence control of annual bluegrass (Poa annua), apply in late summer or early fall (mid-July to mid-September) or in early spring prior to annual bluegrass germination.
- For pre-emergence control of crabgrass, foxtail, barnyardgrass and goosegrass, apply in early spring prior to annual weed grass germination.

AMOUNT TO APPLY: Use 5.9 lbs. per 1,000 sq. ft. or 256 lbs. per acre (equivalent to 2 lbs. of active ingredient per acre.)

HOW TO APPLY: Use a fertilizer spreader, avoiding types which would apply product in narrow, concentrated bands. Apply evenly over the turf without skips or misses. A more uniform application can be made by spreading half the required amount of product over the area to be treated and then applying the remaining half at right angles, or crosswise.

RESEEDING: Chemicals which effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 12 to 16 weeks after application. When reseeding, it is necessary that proper irrigation be maintained. To insure satisfactory reseeding results after using this product, the amount of seed used should be increased.

PRECAUTIONS: Do not apply in spring to turfgrass planted the previous fall. Do not apply to bentgrass putting greens or Dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, or Centipedegrass until these turfs are well established. If verticutting is done, do not apply for 2 to 3 weeks. When applied as recommended, this product will thin fine-leaved festucos.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.