

(Front Panel)

WIDE SPECTRUM

WEED AND FEED

READY-TO-USE

FOR USE WITH A HOSE-END SPRAYER ON GRASS AND ORNAMENTAL TURF
NO MIXING

TREATS 5,000 to 7,500 SQ. FT.†

CONTAINS 2,4-D, MECOPROP, DICHLORPROP AND A 15-0-0 FERTILIZER

1064
PM 23
8123-204

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxy Acetic Acid*	3.26%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	3.28%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	3.22%

INERT INGREDIENTS	90.24%
TOTAL	100.00%

Isomer Specific by AOAC Method

*2,4-Dichlorophenoxyacetic Acid equivalent	2.7%	.26 lbs/gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid equivalent	2.7%	.26 lbs/gal.
***2-(2,4-Dichlorophenoxy)propionic Acid equivalent	2.7%	.26 lbs/gal.

GUARANTEED ANALYSTS:

Total Nitrogen (N)	15%
15% Urea Nitrogen	

TRIAMINE IS A TRADE MARK OF RIVERDALE CHEMICAL COMPANY
† Read directions for different grasses

Patent Applied For.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

MADE IN U.S.A.
MAY 10 1987
8123-204


EPA REG. NO. 8123-
EPA EST. NO. 8123-IL-1

NET CONTENTS: _____

NET WEIGHT: _____

Sold By

FRANK MILLER & SONS, INC.
13831 SOUTH EMERALD AVENUE
CHICAGO, ILLINOIS 60626



(Right Side Panel)

STORAGE: Keep from freezing. To be stored in original container and placed in a locked area inaccessible to children. Do not store near food or feedstuff.

PESTICIDE DISPOSAL: Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

NOTICE

Seller warrants that this product conforms to the ingredient statement of the label. Since the conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, seller makes no claims other than those stated on this label.

(Left Side Panel)

COVERAGE MEASURING GUIDE

(Not to scale)

1000 Sq. Ft.

1875 Sq. Ft.

2000 Sq. Ft.

3750 Sq. Ft.


3000 Sq. Ft.

6525 Sq. Ft.

4000 Sq. Ft.

5000 Sq. Ft.

7500 Sq. Ft.



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Wash thoroughly with soap and water after handling.

IF SWALLOWED: Get medical attention.

IF IN EYES: Immediately flush eyes with plenty of water.

IF ON SKIN: Wash with plenty of soap and water.

If irritation develops, get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABELING. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

5,000 Sq. Ft. Coverage for Bahia, Bermuda, Blue Grass, Fescue, Rye and Zoysia Grasses

7,500 Sq. Ft. Coverage for (spring and fall application only) St. Augustine, Centipede and Bent Grasses. Some temporary injury may occur with these grasses.

Do not exceed recommended rate because damage to lawn may occur. Do not apply to Dichondra, Lippia and Clover lawns. Do not use this product for controlling weeds in vegetable gardens, flower beds or around shrubs or ornamental plants.

WEEDS CONTROLLED

Black Medic
 Buckhorn
 Buttercup
 Canada Thistle
 Carpet Weeds
 Chickweed
 Chicory
 Clover
 Cocklebur
 Dandelion
 Dog Fennel
 English Daisy
 Florida Pursley
 Frenchweed
 Ground Ivy
 Heal-All
 Heartloaf Drymary
 Henbit

Jimson Weed
 Knotweed
 Lambsquarter
 Mallow
 Morning Glory
 Mustard
 Oxalis
 Pennywort
 Pepperweed
 Pig Weed
 Plantain
 Poison Ivy
 Poison Oak
 Purslane
 Rag Weed
 Red Clover
 Red Sorrel
 Sheep Sorrel

Speedwell
 Spurge
 Spur Weed
 Stitchwort
 Toadflax
 Wild Aster
 Wild Carrot
 Wild Garlic
 Wild Geranium
 Wild Lettuce
 Wild Onion
 Wild Radish
 Wood Sorrel
 Yarrow, and other
 broadleaf weeds

WHEN TO USE: Spray anytime during the growing season when weeds are actively growing (except as noted for St. Augustine, Centipede and Bent Grasses). If lawn is extremely dry, water 1 to 2 days before spraying. Do not apply when temperatures are expected to exceed 85° F., or if rain is expected to occur within 24 hours following application. On new lawns wait until grass has reached a height of 2 inches before spraying. Also wait 3 weeks before putting in new grass seed or stolons.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard to control broadleaf weeds may require a second treatment in 3-4 weeks.

(HOSE-END SPRAYER INSTRUCTIONS)

HOW TO APPLY: Use any hose-end sprayer that is designed for use on lawns and makes a minimum of 15 gallons of finished spray, such sprayers are Ortho's Orthomatic Lawn Sprayer, Ortho's and Black Leaf's Lawn Sprayers (dilution ratio of 60 to 1). The following are adjustable/dial sprayers:

MANUFACTURER	SETTING
Super K-Gro	20
Melnor #94 & #95	11
Melnor #96 & #97	13
Gilmour #362	4
Gilmour #486	12

Special note; Some adjustable sprayers do not hold 32 oz. The above settings are based on using 32 oz. per 5,000 or 7,500 sq. ft. The entire contents of this bottle must be used to make the proper recommended application rate.

Determine area to be sprayed. Apply at a rate of 32 oz. (1 quart) per 5,000 square feet or 7,500 for spring and fall application on St. Augustine, Centipede or Bent Grasses. Fill sprayer jar with appropriate amount of this product. Do not dilute this product with water when filling sprayer jar. The water will automatically mix with this product during spraying. Attach sprayer to garden hose while holding sprayer at waist level, point in direction away from face and body. Slowly walk back and forth at a steady pace. Overlap slightly for uniform coverage. Do not exceed recommended rate because damage to lawn may occur.

(DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS)

HOW TO APPLY: Shake well before applying. Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. Turn on the water at the tap. Place finger over hole on right side of nozzle and the Weed and Feed 15-0-0 will mix automatically at the proper ratio. Point in direction away from face and body. Slowly walk back and forth at steady pace. Overlap slightly for uniform coverage. Do not water fertilized lawn for at least 24 hours.

Marks on left side panel of bottle indicate square feet covered.

