

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS:

STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc. or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

K-O-K CLEANSER COMPANY
861 Camden Avenue
Columbus, OH 43201

EPA REG. NO. 8154-6 EPA EST. NO.
NET CONTENTS

8154-6
FEB 27 1981

**SODIUM
HYPOCHLORITE
CONCENTRATE
SOLUTION**

**"FOR SWIMMING POOL CHLORINATION
AND SANITIZING."**

ACTIVE INGREDIENT
SODIUM HYPOCHLORITE 9.2%
INERT INGREDIENTS 90.8%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

31

**DIRECTIONS FOR
GENERAL CLASSIFICATION**

IT IS A VIOLATION OF FEDERAL LAW TO
USE THIS PRODUCT IN A MANNER INCONSISTENT WITH

DIRECTIONS FOR SWIMMING POOL CHLORINATION
For a new pool or for spring start-up, superchlorinate with this sodium hypochlorite solution for each 2500 gallons of water, add 5 ppm available chlorine by weight.
For Pool Maintenance:
A) Adjust pool water pH to 7.6 - 7.8 range and maintain. B) Add manually or by a feeder device this sodium hypochlorite solution to maintain an available chlorine residual of 0.6 to 1.0 ppm available chlorine by weight. Frequency of application will depend upon temperature and bather load. Use test kit to make certain the pH and chlorine levels are maintained.

DIRECTIONS FOR DISINFECTION OF HOME WELL WATER
Dilute this sodium hypochlorite solution in the ratio of 1 part solution to 9 parts softened water. Mix thoroughly and begin feeding of solution with a hypochlorinator or a free available chlorine residual of at least 1.0 ppm throughout the distribution system, as determined by a spot sample of water contacted at a faucet. The National Sanitation Foundation Drinking Water Manual, 1972, is a good reference for this purpose.

DIRECTIONS FOR SANITIZING FOOD PROCESSING EQUIPMENT
Clean equipment in the normal manner. Just before use, wash with this sodium hypochlorite solution containing 100 ppm available chlorine for a minimum of 10 minutes. Rinse thoroughly with water. The National Sanitation Foundation Drinking Water Manual, 1972, is a good reference for this purpose.

NOTE: This product should not be used on food contact surfaces. It is not a food grade product. It is a strong oxidizing agent and should be stored in a cool, dry place.

STORAGE AND DISPOSAL
Store in original container. Do not use if container is empty. Corrosive. In case of spill, food and equipment should be thoroughly washed and rinsed with water. Placing in trash collection or burying in soil is not recommended. This product should be stored in a cool, dry place.

FEB 27 1981

**R SWIMMING POOL CHLORINATION
AND SANITIZING."**

**KEEP OUT OF REACH OF CHILDREN
DANGER**

See additional precautions on side panel

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING

DIRECTIONS FOR SWIMMING POOL CHLORINATION
 1. Turn the pump on for 30 minutes. Add 1 1/2 oz. of this solution per 10,000 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.
 For Pool Maintenance:

B) Add gradually to a feeder device the sodium hypochlorite solution at a rate to maintain a free chlorine residual of 1.0 to 1.3 ppm. One half pint (18 fl. oz.) of this sodium hypochlorite solution for each 6500 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend on temperature and number of swimmers. Use test kit to make certain the pH and free chlorine residual are in the proper range.

[illegible]

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR SERVING EQUIPMENT:
Clean equipment for the small machine just before use, e.g., use a detergent which is a pH 11.5-12.0. Rinse thoroughly with clean water. If the equipment is made of stainless steel, it can be in contact with a solution of 1:1000 bleach (sodium hypochlorite). Do not use in the same solution as equipment used for preparing food. Wash and soak in water for 10 minutes. Rinse thoroughly with clean water. Sterilize in peracetic acid solution or by steam sterilization. Do not use in the same solution as equipment used for preparing food.

STORAGE AND DISPOSAL:

Corrosive, may cause severe skin irritation on broken skin. Causes eye damage. Do not get on clothing. Wear goggles or face shield when handling this product. Wash off clothing vapors. Vacate poorly ventilated area. Do not return until odors have dissipated.

D CHEMICAL HAZARDS:

manufactured by

1-6 EPA EST. NO.

CONTENTS

31

12