

PM 31

8155-12

8-18-87

1042

AUG 18 1987

Canberra Corporation
3610 Holland-Sylvania Road
Toledo, OH 43615

Attention: Mr. John Wiegand

Gentlemen:

Subject: Sanitizer Viricidel Husky 803 S/V Disinfectant
EPA Registration No. 8155-12
Your Submission Dated April 4, 1987
EPA Received Date May 13, 1987

The amendment label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable provided that you:

1. Make the labeling change listed below before you release the product for shipment bearing the amended labeling:
 - Delete the words "And Other" from the subheading "For Hospital, Institutional, Industrial, School, Dairy, and other Farm Use" because these words are ambiguous and because there are no directions for other (than dairy) farm use on the label.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

John H. Lee
Product Manager (31)
Disinfectants Branch
Registration Division (TS-767C)

Enclosure

94693:I:Lee:I-12:KEND:6/29/87:7/9/87:ron:VO:EK:lisa

CONCURRENCES

SYMBOL	ORIGINATOR							
SURNAME								
DATE								


OFFICIAL FILE COPY

ACCEPTED
with COMMENTS
in EPA Letter Dated:

AUG 18 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

8155-12



TO OPEN CHILD RESISTANT
CAP: Use fingernail or coin to
pull back and push down rear-
most ridge on spout. Push
spout down following use.



Surfaces in Food Contact Locations. For sanitizing food processing equipment in meat and poultry plants, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard non-porous surfaces. No Potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. in 4 gallons of water (150ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush or mechanical spray device. No Potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz. 4 gallon solution. No Potable water rinse is required.

At 1 oz. 4 gallons the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1978 Recommendations of the U.S. Public Health Services in waters up to 800 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*. The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of 1 oz. in 4 gallons of warm water. No Potable water rinse is required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

Precautionary Statements
Hazard to Humans and Domestic Animals
DANGER

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not breathe the spray mist. Avoid contamination of food.

Storage and Disposal

- Do not contaminate water, food, or feed by storage or disposal.
- Do not store on side.
- Avoid creasing or impacting of side walls.

Manufactured by:
CANBERRA CORPORATION
3610 HOLLAND-SYLVANIA RD.
TOLEDO, OHIO 43615

EPA Registration No. 8155-12 EPA Establishment No. 8155-OH-1.

PESTICIDE DISPOSAL
Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of rinse is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL
Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.
Metal Container: Triple rinse (or equivalent), then offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. (If container is 1 gallon or less, use this container disposal statement.)

Do not reuse empty container (bottle, can, bucket). Wrap container and put in trash.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions. For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails, and other hard non-porous surfaces.

Add 3.5 oz. of this product to 4.5 gallons water. Apply to previously cleaned hard surfaces. At this use level, it is effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* in the presence of 5% blood serum when evaluated by the AOAC Use-Dilution test.

Virucidal Performance. At a 3.5 oz. 4.5 gallon use level this product was evaluated in the presence of 5% serum and found to be effective against the following viruses: Herpes Simplex Types 1 and 2, Vaccinia, and Influenza A₂ (Hong Kong) on inanimate hard non-porous environmental surfaces.

Disinfection in Institutions, Industry and Schools. Add 2 oz. of this product to 4.5 gallons of water.

Fungicidal Performance. At 0.5 oz. to 2.25 gallons of water use level, it is an effective fungicide against *Trichophyton mentagrophytes* (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities, utilizing the AOAC Fungicidal Test.

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

*SANITIZER/VIRUCIDAL

HUSKY 803

*S/V DISINFECTANT

DISINFECTANT-SANITIZER-FUNGICIDE-VIRUCIDE*

-DEODORIZER WITH

ORGANIC SOIL TOLERANCE FOR HOSPITAL, INSTITUTIONAL, INDUSTRIAL, SCHOOL, DAIRY AND OTHER FARM USE

ACTIVE INGREDIENTS

Octyl decyl dimethyl ammonium chloride	2 250%
Didecyl dimethyl ammonium chloride	1 125%
Dioctyl dimethyl ammonium chloride	1 125%

Alkyl (C ₁₂ , 50%, C ₁₄ 40%, C ₁₆ 10%) dimethyl benzyl ammonium chloride	3 000%
INERT INGREDIENTS	92 500%
	100 000%

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DANGER...KEEP OUT OF REACH OF CHILDREN

Net Contents: 128 Fl. Oz. (1 U.S. Gal.) 3.78 Litres

safe