

MAR 15 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 8192-20

Net Contents: 50 lb
USDA Registration No. 8192-20

Preforam[®]

(C-6989) 15 G HERBICIDE

PREFORAN is a selective herbicide for preemergence weed control in soybeans. Do not use treated plants for feed or forage within 60 days of application.

Active Ingredient:

p-nitrophenyl α,α,α -trifluoro-2-nitro-	
p-tolyl ether	15%
Inert Ingredients	85%
	Total 100%

CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS. May be harmful if swallowed. May cause eye irritation. Avoid contact of dust to eyes. Wear safety goggles when handling. In case of contact flush with water and get medical attention. Avoid contact with the skin. Wear clean clothes and wash clothing before reusing. Wash hands with soap before eating, drinking, or smoking. Do not contaminate food or feed. Do not reuse container. Destroy empty containers by burning. Do not inhale smoke from burning containers.

Toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of waste and containers. Apply this product only as specified on the label.

NOTICE

I
B
A

C
I
B
A

(C-6989)

15 G HERBICIDE

PREFORAN is a selective herbicide for preemergence weed control in soybeans. Do not use treated plants for feed or forage within 60 days of application.

Active Ingredient:

p-nitrophenyl α,α,α -trifluoro-2-nitro-
p-tolyl ether

15%

Inert Ingredients

85%

Total 100%

CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS. May be harmful if swallowed. May cause eye irritation. Avoid contact of dust to eyes. Wear safety goggles when handling. In case of contact flush with water and get medical attention. Avoid contact with the skin. Wear clean clothes and wash clothing before reusing. Wash hands with soap before eating, drinking, or smoking. Do not contaminate food or feed. Do not reuse container. Destroy empty containers by burning. Do not inhale smoke from burning containers.

Toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers. Apply this product only as specified on this label.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. In no case shall CIBA or the seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA and the seller offer this product, and the buyer and user accept it, subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of CIBA.

CIBA AGROCHEMICAL COMPANY
Division of CIBA-GEIGY Corporation
556 Morris Avenue, Summit, N.J. 07901

Preforan^(C-6989) 15 G HERBICIDE

GENERAL INFORMATION

PREFORAN (C-6989) 15 G Herbicide is a granular product effective for the control of certain annual grass and broadleaf weeds when applied preemergence to soybeans. PREFORAN may be applied as a broadcast or as a band treatment to the surface of the soil at planting or at any time preemergence to the crops and weeds.

The seedbed should be well prepared and as free as possible from trash and clods. Best results are obtained when granules lie undisturbed on an even flat surface not subject to shifting by wind or rain.

Do not soil incorporate PREFORAN (C-6989) 15 G Herbicide.

Use only one application to the crop during any one year. Do not follow the treated crop within the same year with any other crops except soybeans.

The granular size of this product is 30/60 mesh.

WEED CONTROL

PREFORAN is effective for controlling certain annual grass and broadleaf weeds in soybeans, such as:

Annual Grass Weeds

Brachiaria
Crabgrass
Giant Foxtail
Goosegrass
Green Foxtail
Yellow Foxtail

Annual Broadleaf Weeds

Carpetweed
Common Ragweed
Jimsonweed
Lambsquarters
Pigweeds
Purslane
Smartweeds
Wild Mustard

APPLICATION DIRECTIONS

Broadcast Treatment — Apply 25 to 30 pounds of PREFORAN 15 G per acre to the surface of the soil with conventional granular applicator. PREFORAN may be applied at the time of planting soybeans, or at any time preemergence to the crop and weeds.

Band Treatment — A band treatment of PREFORAN is made in the same manner as a broadcast treatment, except proportionately less herbicide is used. For example, for 14-inch band on 42-inch row use 1/3 of the broadcast rate. A minimum band width of 12-14 inches is recommended, which will reduce the need for close cultivation. The following formula may be used to calculate the amount of PREFORAN per acre for band treatment:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Rate per acre for broadcast treatment} = \text{Amount needed for band treatment}$$

Do not cover treated band with soil while cultivating.

MAKE CERTAIN GRANULAR APPLICATOR HAS BEEN THOROUGHLY CLEANED BEFORE USING THIS PRODUCT.

Manufactured for

CIBA AGROCHEMICAL COMPANY Division of CIBA GEIGY Corporation, 116 Morris Avenue, Summit, New Jersey, 07901

016

11111