

PROPOSED LABEL

MAN

Y

PURPOSE  
TRIPLE FUNGICIDE  
TREATMENT



of Germinating Seeds and Seedlings of

BEANS, BEANS, AND PEAS

Seed Corn Maggots, Wireworms,  
Seed Decay and Seedling Blights

~~SPECIMEN~~

Up Out of the Reach of Children

(see cautionary statements on back panel)

T

CHIPMAN 3-WAY DIRECTIONS

One-ounce Measuring Scoop Enclosed

Product Information:

3-WAY TRIPLE PURPOSE SEED TREATMENT is a specially formulated combination of diazinon for protection against seed corn beetles and seed corn maggots and lindane for protection against wireworms and captan for protection against fungus diseases of seeds and seedlings. Treatment supplies extra amount of captan which is beneficial, especially under stress conditions.

Recommendations:

Planter box application: Use on seed by mixing with the seed in the planter box at a rate of 3 ozs./bushel.

Size of Planter Box	No. of 1 oz. Scoops/Planter Box
1/2 bushel	1 1/2 heaping scoops (3 oz./bus.)
1 bushel	3 heaping scoops (3 oz./bus.)

Use only recommended rates. Lower amounts may not give the desired control. Excessive amounts may cause seed injury.

12 ounce container - treats 4 bushels.

Notice to Buyer:

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

E.P.A. Registration No. 8203-2

BEANS, BEANS, PEAS, AND SORGHUM

amping Off, Seed Decay, Seedling  
lso against Seed Corn Beetle  
ans), Seed Corn Maggots (corn,  
beans, and peas), and Wireworms  
(beans, beans, and sorghum)

ACTIVE INGREDIENTS:

+Captan .....	33.5%
++0,0-diethyl 0-(2-isopropyl-6-methyl- 4-pyrimidinyl) phosphorothioate.....	11.0%
Lindane (Gamma Isomer of Benzene Hexachloride).....	15.6%
INERT INGREDIENTS .....	38.9%
	100.0%

CONTAINS GRAPHITE

+N-trichloromethylthio-4-cyclohexene-1,  
2-dicarboximide  
++DIAZINON<sup>(R)</sup> (TM CIBA-GEIGY Corporation)

8203-2

2-24-74

1043

## CHIPMAN 3-WAY INFORMATION

## CAUTION:

MAY BE HARMFUL IF SWALLOWED OR  
ABSORBED THROUGH SKIN

Avoid breathing dust or spray mist.  
Avoid contact with skin, eyes and clothing.  
Wash thoroughly with soap and water after  
handling.  
Avoid contamination of feed and foodstuffs.

First Aid: In case of contact, immediately flush eyes or skin  
with plenty of water for at least 15 minutes while removing  
contaminated clothing and shoes. Call a physician. Wash  
clothing and decontaminate shoes before reuse.

Attention: This product is toxic to fish and wildlife. Do not  
contaminate any water by cleaning of equipment or disposal of  
waste. Treated seed on the soil surface will be hazardous to  
birds and other wildlife.

Do not use treated seeds for food, feed, or oil purposes.

Do not reuse container. Destroy when empty.

Container handling and storage: Protect from excessive heat.  
Store in cool dry place.

## NOTE:

Do not use on seed already treated with an insecticide other  
than Malathion or Methoxychlor (for control of stored grain insects).  
Seed within one month of treatment. Regularly inspect planter  
plates and clean as required. If the seed treatment increases the  
friction between seeds, sufficient to reduce the rate of delivery  
through a grain drill, 1 ounce of graphite per bushel of seed  
should be added as a drill lubricant. Adjust the seed drill to  
give the required rate of seed delivery with the treated seed.



1073

ACCEPTED

FEB 21 1974

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
ED UNDER NO. 1203-1-1 SUBJECT  
TO ATTACHED COMMENTS.