

CHIPMAN 3-WAY INFORMATION

CAUTION: May be harmful if swallowed or absorbed through skin.

Avoid breathing dust or spray mist.

Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

Avoid contamination of feed and foodstuffs.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician. Wash clothing and decontaminate shoes before reuse. Do not reuse container. Destroy when empty. Do not use treated seeds for food, feed or oil purposes.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish, birds and other wildlife. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER HANDLING AND STORAGE: Protect from excessive heat. Store in cool dry place.

NOTICE TO BUYER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use. Seller disclaims all other warranties, expressed or implied, including any warranty of FITNESS OR MERCHANTABILITY. Seller shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product and Seller's sole liability and Buyer's exclusive remedy shall be limited to the refund of the purchase price.

CHIPMAN CHEMICALS LIMITED
Stoney Creek, Ontario

Contents Made in Canada

Container Made in U.S.A.

CHIPMAN 3-WAY

TRIPLE PURPOSE INSECTICIDE FUNGICIDE SEED TREATMENT

For the Protection of Germinating Seeds and Seedlings of
CORN, SORGHUM, SOYBEANS, BEANS AND PEAS
against Damping Off, Seed Decay, Seedling Blights; also against
Seed Corn Beetle (corn, beans), Seed Corn Maggots (corn,
soybeans, beans and peas), and Wireworms (corn, soybeans,
beans and sorghum)

ACTIVE INGREDIENTS:

| | |
|---|--------|
| Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide) | 33.5% |
| + O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate | 11.0% |
| Lindane (Gamma Isomer of Benzene Hexachloride) | 16.6% |
| INERT INGREDIENTS | 38.9% |
| CONTAINS GRAPHITE | 100.0% |

+ DIAZINON (TM Ciba-Geigy Chemical Corp.)

CAUTION: Keep Out of the Reach of Children

(See additional precautionary statements on side panel)

12 OZ. NET WEIGHT

CHIPMAN 3-WAY DIRECTIONS

One-ounce Measuring Scoop Enclosed

PRODUCT INFORMATION:

3-WAY TRIPLE PURPOSE SEED TREATMENT is a specially formulated combination of Diazinon for protection against seed corn beetles and seed corn maggots and lindane for protection against wireworms and captan for protection against fungus diseases of seeds and seedlings. Treatment supplies extra amount of captan which is beneficial, especially under stress conditions.

RECOMMENDATIONS:

Planter box application: Use CHIPMAN 3-WAY at a rate of 3 ozs./bushel (3 heaping one oz. scoops/bushel). Mix thoroughly with seed in the planter box as follows — Fill the planter box one-half full of seed then treat with one-half amount of product and mix well by stirring with a stick or paddle until color is imparted to all seed. Add remaining seed to the planter box and cover with balance of product. Mix thoroughly as above.

Use only at the recommended rate. Lower amounts may not give the desired control. Excessive amounts may cause seed injury.

12 ounce container — treats 4 bushels.

NOTE:

Do not use on seed already treated with an insecticide other than malathion or methoxychlor (for control of stored grain insects). Seed within one month of treatment. Regularly inspect planter plates and clean as required. If the seed treatment increases the friction between seeds sufficient to reduce the rate of delivery through a grain drill, 1 ounce of graphite per bushel of seed should be added as a drill lubricant. Adjust the seed drill to give the required rate of seed delivery with the treated seed.

EPA Registration No. 8203-2

RECEIVED

8203-2
JAN 29 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.

8203-2

1-29-75

1041