Read Directions – Heed Cautions WARNING!

May be fatal or harmful if swallowed. May be absorbed through skin. Protect from excessive heat. Store in cool, dry place.

Do not breathe dust, vapor or fumes. Do not get in eyes or on skin or clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. Do not contaminate feed or food. Destroy empty container — crush, bury in non-crop lands away from water supplies. Never re-use. If swallowed, give raw white of egg or milk. Produce vomiting with either 1 tablespoon of salt or mustard in a glass of warm water, or warm soapy water. Call a physician. Do not use treated seeds for food, feed or oil purposes.

ENVIRONMENTAL CAUTIONS:

This product is toxic to fish, birds, and other wildlife. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Do not allow treated seeds to be exposed on soil surface. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, disposal of wastes. Dispose of excess treated seeds or wastes by burial. Apply this product only as specified on this label.

Note:

Do not use on seed already treated with an insecticide other than Malathion or Methoxychlor (for control of stored grain insects). Seed within one month of treatment. Regularly inspect planter plates and clean as required. If the seed treatment increases the friction between seeds, sufficient to reduce the rate of delivery through a grain drill, I ounce of graphite per bushel of seed should be added as a drill lubricant. Adjust the seed drill to give the required rate of seed delivery with the treated seed.

CHIPMAN CHEMICALS LIMITED

Stoney Creek, Ontario

Container Made in U.S.A.

Contents Made in Canada

For the Protection of Germinating Seeds and Seedlings of

CORN, SOYBEANS, BEANS AND PEAS against Seed Corn Beetles, Seed Corn Maggots, Damping Off, Seed Decay and Seedling Blights

Active Ingredients:

Captan (N-[(trichloromethyl)thio]-4-cyclohexene1,2-dicarboximide) 37.5%

*O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl)
phosphorothioate 25.0%
Inert Ingredients 37.5%

CONTAINS GRAPHITE 100.0%

*Diazinon (TM Ciba-Geigy Chemical Corp.)

Warning! Keep Out of the Reach of Children

(See additional precautionary statements on back panel)

8 OZ. NET WEIGHT

Product Information:

CHIPMAN I. F. DUAL PURPOSE SEED TREATMENT is a specially formulated Planter Box combination of diazinon for protection against seed corn beetles and seed corn maggots and captan for protection against fungus diseases of seeds and seedlings.

Recommendations:

Planter box application: Use on seed by mixing with the seed in the planter box at a rate of 2 ozs./bushel.

Size of Planter Box

No. of 1 oz. Scoops/Planter Box

Size of Planter Box

½ bushel

1 bushel

8 ounce container – treats 4 bushels

No. of 1 oz. Scoops/Planter Box

1 heaping scoop (2 oz./bus.)

2 heaping scoops (2 oz./bus.)

For seed corn beetles – While the above rate will provide some protection against seed corn beetles, preliminary results indicate that 4 to 6 oz./bu. (4 to 6 scoops) may be more effective for protection against the seed corn beetles.

Note:

When starting with an empty planter box, place enough treated seed in the planter box to cover the bottom. Treat this seed in a separate container that can be destroyed. Add the remaining seed to the planter box and treat by thoroughly stirring with a paddle so that the seed is coated with seed treatment.

Notice to Buyer:

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Registration No. 8203-10 Am 9046 Rev. 10-75