Read Directions

STOP

Heed Cautions

GRANOX P-F-M

PEARUT AND SOYBEAR STED FUNGICIDE

A Combination of Fungicides

For the Control of Sold-borne and Soil-bolne Discusos Including Dauping off, Seed Decay Organisms and Seedling Blight. Also contains Holybdenum.

Active	Ingredients:		
•	Maneb (Hanga	mous Ethylonobis (dithiocarbamate))richloromethylthio)-4-cyclohuxeau,	30.007
	Captan' (II-(1	richloromethyithio)-4-cyclohuxenu,	90 600
	2-d1	carboximide)	30. 000
Inert	Ingred ie nts*		40.00%
			100.00%
*Added	Ingredient:	Sodium Molybdate	1.00%

C A U T I O N - KEEP OUT OF REACH OF CHILDREN

HARMFUL IT SMALLOWED

Avoid inhaling dust. Wear respirator. Avoid prolonged or repeated contact with skin. Wash with scip and water after handling. Avoid contamination of feed and food.

Keep container tightly closed when not in use.

Store seed in a ventilated place away from foodstuffs.

Avoid contamination of feed and foodstuffs.

Do not allew to become wet during storage. If maneb becomes wet in storage, its fungicidal effectiveness may be reduced and it may also produce vapors which may be flammable. Keep away from flames or sparks.

Toxic to fish. Do not contaminate any body of water by cleaning of equipment and disposal of wastes and containers.

Container should be disposed of by crushing and burying with wastes.

IMPORTANA: Treated seed should not be used for huran consumption nor as animal food.

Treated seed should be labelled as follows: TREATED SEED -- DO NAT USE FOR FOOR, FEED OR OIL PURIOSES

PRODUCT INFO MITTON

GRANOX P-F-M is a special seed treatment formulation of a combination of fungicides which, when used at the correct dosage rate, protects peanuts and soybeans against seed and soil borne diseases. GRANOX P-F-M provides essential molybdenum for plant growth whose soils are found to be difficient in available molybdenum.

DIRECTIONS FOR USE

Pernuts: - Apply GRAMOX P-F-M to peanut seed at 5 to 7 oz. per 100 Ms. GRAMOX P-F-M may be used in any dry treating equipment such as co-croial mechanical treaters, home made rotary dress or for planter box application. Thoroughly mix treatments with seed in the planter box. Do not use slurry treatments on pranuts.

Soyleans: - Apply GRANDA P-F-M to soylean seed at 2 to 3 ozs, per bushel. GRAMOX P-F-M may be applied as a planter box application, or by the use of any dry treating equipment or for slurry treatment. (for soyberns only)

FOR SLURRY TREATMENT

Number of Pounds/CRANOX P-F-M per Gallen of Slurry

Fluid oze of slurry per bushel of soyheans	4	8	
GRANOX P-F-11 oz/bus.			
2 3	4 1bs. 6 1bs.	2 1bs. 3 1bs.	

Note: Seed treated by the slurry method should not be stacked until the seed is dry. Store in a well ventilated area.

Seed must be well cured and cleaned before treating.

Avoid very deep secding and exceptionally early sowing under poor growing conditions for maximum benefits. Use only recommended rates. Learn amounts may not give the desired control. Higher a punts may cause seed injury.

Thoroughly mix treatment with the seed in the planter box.

Wear a properly designed dust mask and goggles and change filters frequently while treating seed and handling treated seed.

Regularly inspect planter plates of seed drill and clean as required. If coating of planter plate occurs (i.e. under conditions of high hunidity), add small amounts of bone meal or graphite to facilitate uniform seeding.

NOTICE

Seller's guarantee shall be limited to the terms of the label and subject therete the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Registration No. 8203-13
50 Pounds Net Weight

CHIPPAN CHEMICALS LIMITED
Hamilton Ontario

VUVUV