

NET WEIGHT 35 lbs.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

WARNING

Treat outdoors or in a well ventilated place. May be fatal or harmful if swallowed. May be absorbed through skin. Do not breathe dust, vapor or fumes. Do not get in eyes or on skin or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. Do not contaminate feed or food. Do not use treated seed for food, feed or oil purposes.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Seed commercially treated with AGROX D-C-S SEED TREATMENT must be labelled in accord with all applicable requirements of the Federal Seed Act.

FIRST AID

If swallowed: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin: Remove contaminated clothing and wash affected areas with soap and water.

If in eyes: Flush eyes for at least 15 minutes with water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

PRODUCT INFORMATION

AGROX D-C-S SEED TREATMENT is a specialty formulated combination of DIAZINON for planter box application or for slurry treatment for protection against seed corn maggots and seed corn beetles, and captan for protection against fungus diseases of seeds and seedlings.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

For Seed Corn Maggots, Corn Peas, Soybeans, Beans, all types (including snap and field) - Apply 3 ozs per bushel of seed.

For Slurry Machines: adjust to apply 4 to 8 fl. oz. of slurry per bushel of seed and use 6 to 3 lbs. respectively of seed treatment per gallon of slurry. If machine

AGROX[®] D-C-S

INSECTICIDE FUNGICIDE SEED TREATMENT

For the Protection of Germinating Seeds and Seedlings
of CORN, SOYBEANS, BEANS, AND PEAS against
SEED CORN BEETLES, SEED CORN MAGGOTS,
DAMPING-OFF, SEED DECAY AND
SEEDLING BLIGHTS

ACTIVE INGREDIENTS

Captan (N-(1-chloromethyl)thio-4-cyclohexene-1,2-dicarboximide) 28.37
Related derivatives 1.63

*O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 16.60

INERT INGREDIENTS 53.40

TOTAL 100.00

Contains Graphite

*DIAZINON (Registered trade mark of Ciba-Geigy Chemical Corporation)

KEEP OUT OF REACH OF CHILDREN

WARNING

See back panel for additional precautionary statements

EPA Est 8203-CN 1
EPA Reg No 8203-16

AGROX is a registered trade mark of Chipman Inc., Canada

applies more or less slurry, adjust the amount of seed treatment so that 3 ounces are used per bushel of seed or -

For Hand Mixing: for every bushel of seed mix 3 ounces of seed treatment per 8 fl. oz. of water. Keeping mixture agitated and seed treatment in suspension in the water, pour over the seed and mix until uniformly coated. Allow to dry before seeding or bagging.

For Seed Corn Beetles: while the above rate of AGROX D-C-S SEED TREATMENT (3 ounces per bushel) may provide some protection against seed corn beetles, preliminary results indicate that 4 to 5 oz. bushel may be more effective for protection against the seed corn beetles.

NOTE: Do not use on seed already treated with an insecticide other than Malathion or Methoxychlor (for control of stored grain insects). Seed within one month of treatment. Regularly inspect planter plates and clean as required. If the seed treatment increases the friction between seeds, sufficient to reduce the rate of delivery through a grain drill, 1 ounce of graphite per bushel of seed should be added as a drill lubricant. Adjust the seed drill to give the required rate of seed delivery with the treated seed.

STORAGE AND DISPOSAL

Storage - Protect from excessive heat. Store in a cool, dry place away from water, food and feed.

Disposal - Do not re-use empty container. Do not contaminate water, food, or feed by disposal. Open dumping is prohibited. Pesticide that cannot be used should be disposed in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of empty container in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

NOTICE TO BUYER

Sealer's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

1080

M. J. Chipman

CHIPMAN INC.

P.O. Box 401


Stoney Creek, Ontario L8G 3Z1

Canada

CONTAINER AND CONTENTS

MADE IN CANADA

5,000 SQ. FT.
(50'x' 100')
30 LBS.
NET WT.



Arthroban™



TRIPLE ACTION 4
FORMULA FOR ESTABLISHED LAWNS



CRABGRASS
PREVENTER



INSECT KILLER
FERTILIZER



WARNING