

NET CONTENT
5 gallons



ACCEPTED
JUN 1982
Under the Pesticide
Furnish
of 0111
Total
EPA Reg. No. 8203-23

2,4-D AMINE

Liquid
Weed Killer

ACTIVE INGREDIENT

*Dimethylamine salt of 2,4-D dichlorophenoxyacetic acid . . . 49.26%

INERT INGREDIENTS 48.67%

TOTAL 100.00%

*Equivalent to 2,4-D dichlorophenoxyacetic acid 40.91%

This product contains 4 lb of acid equivalent per gallon

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY INFORMATION

EPA Est 8203-CN-1

EPA Reg. No. 8203-23

CHIPMAN INC.
Stoney Creek, Ontario
L8G 3Z1
Canada.

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

DANGER

Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or a face shield and rubber gloves when handling. Harmful if swallowed

STATEMENT OF PRACTICAL TREATMENT

In case of contact with skin, wash with soap and water. For eyes, flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply when weather conditions favour drift from target area

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

Mix thoroughly with clean water and provide continuous gentle agitation during spraying.

Do not spray during periods of high winds. Do not apply 2,4-D to broadleaf crops such as peas, beans, sugar beets, tomatoes or flowers, garden vegetables, trees or shrubs, and do not allow mist or spray to drift onto them. Coarse sprays are less likely to drift.

Do not mix with liquid fertilizers containing nitrogen. **Wheat, Barley and Rye:** Treat only from 3-leaf to early shot-blade stage. If weeds warrant spraying prior to 3-leaf stage, MCPA results in less crop injury than 2,4-D. Do not use on underseeded crops. Do not graze or feed forage from treated fields within 2 weeks after treatment.

Winter Wheat and Fall Rye: Treat in spring when crop commences growth until early shot-blade stage. Do not graze or feed forage from treated fields within 2 weeks after treatment.

Oats: Treat only at the 2-leaf stage or from the 6-leaf to early shot-blade stage. 2,4-D should only be used where MCPA resistant weeds are present since oats are more tolerant to MCPA. Do not graze or feed forage from treated fields within 2 weeks after treatment. SEE NOTE below table of suggested rates.

CORN (field, sweet or popcorn): Apply when corn is 6-8 inches high. When corn is higher than 6-8 inches use drop nozzles directing spray at base of plants up until corn is 16 inches high. Apply at 1 pint per acre in 5 to 25 gallons of water. Do not graze or feed treated corn fodder for 7 days following application.

SUGGESTED RATES TO USE ON CEREAL CROPS (NOT SEEDS TO LEGUMES)

WEED GROWTH	Stage of Weed Kind of Weather	Fluid Ounces		Acres Treated per Gal.
		Chipman 2,4-D Amine per Acre	Ounces 2,4-D Acid per Acre	
Susceptible, (easier to kill) weeds: Blue bur, Cockle bur, Common Plantain, Dandelion (in lawns), Flaxweed, Kochia, Lambs Quarters, Mustards, Pockly Lettuce, Ragweeds, Russian Pigweed, Shepherds Purse, Stinkweed.	Seedling (2-4 leaves) Growing rapidly.	9½	5	13½
	Weeds in bud. Dry or Cool Weather. Heavy infestation.	15¼	8	8½
More resistant (harder to kill) weeds: Chicory, Blue Lettuce*, Common Chickweed, Curled Dock, Dandelions (in fields) Dog and Tansy Meadows, Field Bindweed*, Goat's Beard, Leafy Spurge*, Pepper Grass, Red Root Pigweed, Russian Thistle, Smartweed*, Tumble Weed, Canada* and Sow Thistle, Wild Buckwheat*, Tartary Buckwheat*.	Seedling (2-4 leaves) Growing rapidly.	(17)	(10)	7½
	Weeds in bud. Dry or Cool Weather. Heavy infestation.	(23)	(12)	5½

NOTE: Apply in 5-25 gallons of water per acre. The higher rates (in brackets) may cause injury which will frequently be offset by a higher relative yield from a less weedy crop. On Oats use up to 1 pint per acre. 1 to 1¼ pints may be used on a very weedy crop, with probable injury. *Top growth control for these weeds

PASTURES: If no legumes are present, use 1½-2¾ pints (12-24 ounces 2,4-D acid) per acre (½-1 fl oz per 1000 sq ft) early in summer when weeds are in bud or in the fall. For 2,4-D resistant weeds, use Brushkiller. Repeat treatments may be necessary. Do not apply when grass is in early boot to milk stage where grass seed production is desired. Do not apply to newly seeded areas or after heading begins. Do not graze dairy cattle in treated areas for 14 days after application. Remove meat animals from freshly treated areas for 7 days before slaughter. Withdrawal is not needed if 2 weeks or more has elapsed since the treatment was applied. Do not cut treated grass for hay within 30 days after application.

FAIRWAYS AND LAWNS: Use 1¼ pints per acre. For small areas 2½ teaspoonfuls in 1 gallon water treats 400 square feet. Do not apply to newly seeded areas and do not apply to bentgrass or St. Augustine grass. For lawns and fairways, treat a small area first to test the effect on clover.

ROADSIDES, DRAINAGE DITCH BANKS AND WASTELAND: Use 3½ pints per acre in June, or 5½ pints per acre in July. Use low pressure and apply at least 60 gals. water per acre, using at least 180 gals. water per acre on dense weeds and wild carrot.

Woody Growth: Such as Western Snowberry, Willows, Poplars in pastures, drainage ditch banks, roadsides use 1½ to 3¾ pints per acre (16-32 ounces 2,4-D Acid) in 12 or more gals. water when plants are in full leaf. For mixed woody growth, resistant to 2,4-D, use Brushkiller.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage - Do not store near other pesticides or seeds. **AVOID FREEZING:** If subjected to freezing, warm at least 40°F and mix thoroughly before using. **Pesticide Disposal:** Pesticide or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal: Triple rinse and offer for recycling or reconditioning, or dispose in a sanitary landfill, or by other approved State and local procedures.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.