

NET WT. 2 OZ

AGRICULTURAL
USE ONLY

ACCEPTED

200
NOV 23 1981

Use only as directed. Do not use on food crops.
This product is registered under EPA Reg. No. 8203-28



AGROX®
D-L PLUS

INSECTICIDE-FUNGICIDE
SEED TREATMENT

For the Protection of Germinating Seeds and Seedlings of
CORN, SORGHUM, SOYBEANS, BEANS AND PEAS
against Damping Off, Seed Decay, Seedling Blights; also
against Seed Corn Beetle (corn, beans), Seed Corn
Maggots (corn, soybeans, beans and peas), and
Wireworms (corn, soybeans, beans and sorghum).

ACTIVE INGREDIENTS	%
Captan (N-[(1-chloromethyl)thio]-4-cyclohexene-1,2-dicarboximide)	14.19
Related derivatives	0.81
*O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)phosphorothioate	15.00
Lindane (gamma isomer of benzene hexachloride)	25.00
INERT INGREDIENTS	45.00
TOTAL	100.00

*DIAZINON (Registered trademark of Ciba-Geigy Chemical Corporation)

**KEEP OUT OF REACH OF CHILDREN
WARNING**

See back panel for additional precautionary statements

EPA Est B203-CN-1

EPA Reg. No. 8203-28

AGROX is a registered trade mark of Chipman Inc., Canada

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING

Treat outdoors or in a well ventilated place. May be harmful if swallowed or absorbed through skin. Avoid breathing dust. Wear a properly designed dust mask, goggles and gloves while treating seed. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash clothing and decontaminate shoes before reuse. Avoid contamination of feed and foodstuffs. Do not use treated seeds for food, feed or oil purposes. Apply this product only as specified on the label.

FIRST AID: If swallowed - Call a physician and or the Poison Control Centre immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated. If on skin - Remove contaminated clothing and wash affected areas with soap and water. If in eyes - Flush eyes for at least 15 minutes with water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of water.

PRODUCT INFORMATION

AGROX D-L Plus is specially formulated for use on seed previously treated with Captan or Thiram. Do not use on seed previously treated with an insecticide other than malathion or methoxychlor. AGROX D-L Plus is a combination of DIAZINON for protection against seed corn beetles and seed corn maggots and Captan for protection against wireworms and seedling blights. Treatment supplies additional Captan fungicide to the fungicide already on the seed from storage or other treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Planter box application: Use AGROX D-L Plus at a rate of 2 or 5 bushels (one pouch per bushel). Mix thoroughly with seed in the planter box as follows:

(1) For Conventional Plate Planters and John Deere MAX EMBLSE® Plateless Planter - Pour one bushel of seed into each planter box. Pour enough AGROX D-L Plus on top of the seed to treat one bushel. Mix thoroughly with an axe handle or paddle, until all the seed is uniformly coated.

(2) For All International Harvester Air Planters - Pour 5 bushels of seed into the planter box. Pour enough AGROX D-L Plus on top of the seed to treat 5 bushels. Mix thoroughly with an axe handle or paddle, until all seed is uniformly covered.

Use only at the recommended rate. Lower amounts may not give desired control. Excessive amounts may cause seed injury.

NOTE: Do not use on seed already treated with an insecticide other than malathion or methoxychlor (for control of stored grain insects). Regularly inspect planter plates and clean as required. If the seed treatment increases the friction between seeds sufficient to reduce the rate of delivery through a grain drill, 1 ounce of graphite per bushel of seed should be added as a drill lubricant. Adjust the seed drill to give the required rate of seed delivery with the treated seed.

Clean up for Electric Eye Population Monitors: At the end of each day of use, clean out the electric eye monitors with brushes provided by the planter manufacturer according to their directions.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Protect from excessive heat. Store in a cool dry place away from water, food and feed.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of in a sanitary manner or by other State and local procedures.

NOTICE TO BUYER

Seller's guarantee should be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

661

Manufactured by
Chipman Inc.
Stoney Creek, Ontario, Canada
L8G 3Z1

CONTENTS AND CONTAINER
MADE IN CANADA