DRAFT

NET MEIGHT 10 16.

CRANOER CHE

SOYBEAN SEED TREATMENT

PUNCICIDE

This product contains 9.5% molybdenum from sodium molybdate

ACTIVE INGREDIENTS

Ŧ

Capten (N-[(trichloromethyl)thio]-4-cyclohexane-1,2-dicarboximida) 47

47.3

Related derivatives

ACCEPTED
with COMMENTS
In EPA Letter Dated:

2.7

INERT INCREDIENTS

50.0

TOTAL

DCT 1 4 1382

100.0

Under the Federal Insecticide, Fungicide, and Redenticide Act as amended, for the posticide represent under EPA Ren. No. 8209-302

KREP OUT OF REACH OF CHILDREN

CAUTION

See mide panel for additional precautionary information

EPA Est 8203-CN-1

EPA Reg. No. 8203

GRANOX is a registered trade mark of Chipman Inc., Canada

PRECAUTIONARY STATEMENTS MAZARDS TO BUNANS AND DOMESTIC ANIMALS

CAUTION

May cause eye irritation. Harmful if swallowed. Do not get in eyes.

Avoid contact with skin. Avoid inhalation of dust. Wear a penticide
respirator jointly approved by the Mining Enforcement and Safety

Administration (formerly the U.S. Bureau of Mines) and by the National
Institute for Occupational Safety and Health under provisions of 30 CFR

Part II. Wash with soap and water after handling. Avoid contamination of
feed, food and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

If ewallowed, induce vomiting by drinking 1 or 2 large glasses of water and then touch back of throat with fingers.

If in eyes, flush with clean water for at least 15 minutes and then get medical excention.

ENVIRONMENTAL HAZARDS

Toxic to fish. Do not contaminate any body of water by cleaning of equipment and disposal of wastes and containers. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from budies of water.

PRODUCT INFORMATION

GRANOX CHM, when used at the correct rate, procests seeds from soil and seed bords diseases that cause damping-off, seed decay, and seedling blights. GRANOX CHM provides essential molybdenum for plant growth where soils are found to be deficient in available molybdenum.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

Soybeans: Apply GRANOX CHM to soybean seed at 2 oz per bushel. GRANOX CHM may be applied as a planter box application, or by the use of any dry treating equipment.

Planter box application: Use GRANOX CHM at a rate of 2 oz/bushel (one scoop per bushel). Him thoroughly with seed in the planter box as follows:

For Conventional Piste Planters and MAX-XMERGE® Plateless Planters by

John Deers - Pour one bushel of seed into each planter box. Pour enough

GRANOX CHM on top of the seed to treat one bushel. Stir thoroughly with

an axe handle (purchased and used for this purpose only), wooden lath or

stick, until all the seed is uniformly coated. (MAX-EMERGE® is a reg.

T.M. of Deers and Co.)

MOTE: Avoid very deep seeding and exceptionally early sowing under poor growing conditions for maximum benefits. Use only recommended rates. Lower amounts may not give the desired control. Regularly inspect planter plates of seed drill and clean as required. If coating of planter plate occurs (i.e. under conditions of high humidity), add small amounts of graphite to facilitate uniform seeding.

Clean-up for Electric Sye Population Monitors: At the end of each day of use, clean out the electric-eye monitors with brushes provided by the planter manufacturers, according to their directions.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use. Store treated seed in a ventilated place away from water, food and feed.

PERTICIDE DISPOSAL: Pesticide spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable state or local procedures.

CONTAINER DISPOSAL: Completely empty container and tap side and bottom to loosen clinging particles. Then dispose of in a sanitary landfill or by incideration if allowed by State and Local authorities.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms of the label and Subject thereto, buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

0982

CHIPMAN INC.

Stoney Creek, Ontario, Canada

L8G 321

CONTENTS AND CONTAINER FADE IN CANADA

OCT | 4 1982

8403~32

Grandy CHH S Fungicide ... Seed Treatment

ICI American, Inc. Agricultural Division Wilmington, DE 19897 18 19758

This product is conditionally registered in accordance with FIFMA accordance with FIFMA

- i. Submit and/or cite all data required for registration/reregistration f your product under FIFRA sec. 3(q)(5) when the Agency requires all registrants of similar products to submit such data.
- A. Add the phrase "EPA Registration No. 8303-32" to your label sefore you release the product for elippent.
- 3. Submit five (5) copies of your final printed labeling before you release the product for shipsent. Refer to the A-79 Maclosure for a further description of final printed labeling.

if these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. i(e). Your release for enignent of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Henry H. Jacoby Product Hanager (21)
Fungicide-Herbicide Stanch
Registration Division (73-767)

Edelogaros