

Net Contents: 10 lb.

5 lb.

2 oz.

AGROSOL^R

Systemic Soybean Seed Treatment Fungicide

Contains MERTECTR

This product contains 9.5% molybdenum from sodium molybdate

ACTIVE INGREDIENTS:

	%
Captan (N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide)	23.65
Related derivatives	1.35
Thiabendazole (2-(4-thiazolyl) benzimidazole)	0.68

INERT INGREDIENTS:

TOTAL

ACCEPTED
with COMMENTS
in EPA Letter Dated:

DEC 14 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

8203-40

74.32

100.00

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for additional precautionary information

EPA Est 8203-CN-1

EPA Reg. No. 8203-

AGROSOL is a registered trademark of Chipman Inc., Canada

MERTECTR is a registered trademark of Merck and Co., Inc.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Harmful if swallowed. Corrosive; causes eye damage. Do not get in eyes. Avoid inhalation of dust. Stand upwind while treating or handling treated seed. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration and by the National Institute for Occupational Safety and Health under provisions of 30 CFR Part 11. Avoid contact with skin by wearing rubber gloves while treating or handling treated seed. Avoid contact with eyes by wearing safety goggles while treating or handling treated seed. Wash with soap and water after handling and before eating, drinking or smoking. Avoid contamination of feed, food and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, induce vomiting by drinking 1 or 2 large glasses of water and then touch back of throat with finger. Repeat until vomit fluid is clear. Get medical attention. If in eyes, flush with clean water for at least 15 minutes and then get medical attention. If on skin, wash with soap and water.

ENVIRONMENTAL HAZARDS: Toxic to fish. Do not apply directly to water or wetlands. Do not contaminate any body of water by cleaning of equipment and disposal of wastes and containers. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water.

PRODUCT INFORMATION: AGROSOL, when used at the correct rate, protects seeds from soil and seed borne diseases that cause damping-off, seed decay, seedling blights and pod and stem blight (*Phenopsis* spp.) during the germination and early seedling stages. AGROSOL provides essential molybdenum for plant growth where soils are found to be deficient in available molybdenum.

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

Apply AGROSOL to soybean seed at the rate of 2 oz. per bushel (one scoop per bushel). AGROSOL may be applied as a planter box application, or by the use of any dry treating equipment.

Planter box application - Use AGROSOL at a rate of 2 oz. per bushel (one scoop per bushel). Mix thoroughly with seed in the planter box as follows:

For Conventional Plate Planters and MAX-EMERGER Plateless Planters by John Deere - Pour one bushel of seed into each planter box. Pour enough AGROSOL on top of the seed to treat one bushel. Stir thoroughly with an axe handle (purchased and used for this purpose only) until all the seed is uniformly coated. (MAX-EMERGER is a reg. T.M. of Deere and Co.).

NOTE: Avoid very deep seeding and exceptionally early sowing under poor growing conditions for maximum benefits. Use only recommended rates. Lower amounts may not give the desired control. Regularly inspect planter plates of seed drill and clean as required. If coating of planter plate occurs (i.e. under conditions of high humidity), add small amounts of graphite to facilitate uniform seeding.

Clean-up for Electric Eye Population Monitors: At the end of each day of use, clean out the electric-eye monitors with brushes provided by the planter manufacturers, according to their directions.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed on site or at an approved waste or disposal facility.

CONTAINER DISPOSAL: Completely empty container and tap side and bottom to loosen clinging particles. Empty residue into application equipment then dispose of in a sanitary landfill or by incineration if allowed by state and local authorities.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms of the label and subject thereto, buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

0984

CHIPMAN INC.

Stoney Creek, Ontario, Canada

L8G 3Z1

CONTENTS AND CONTAINER MADE IN CANADA

10-01-00

10-01-00