

8203-40

PM 21

1065

Net Contents: 5 lb.

10 lb.

AGROSOL™

Systemic Soybean Seed Treatment Fungicide

Contains MERTECT®

This product contains 9.5% molybdenum from sodium molybdate

ACTIVE INGREDIENTS	By Wt.
*Captan	24.44%
Related Derivatives	0.56%
**Thiabendazole	0.68%
INERT INGREDIENTS	74.32%
*N-Trichloromethylthio-4-cyclohexene-1,2-dithiocarbamate	
**2-(4-thiazolyl) benzimidazole	
TOTAL	100.00%

This product contains the toxic inert Rhodamine B

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol.

If in eyes, flush with plenty of water. Call a physician.

If inhaled, remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin, wash with plenty of soap and water.

See side panel for additional precautionary information

CHIPMAN, A Business Unit of C-I-L Inc.

Stoney Creek, Ontario, Canada L8G 3Z1

EPA Est 8203-CN-1

EPA Reg. No. 8203-40

AGROSOL is a trademark of C-I-L Inc., Canada

MERTECT is a registered trademark of Merck & Co., Inc.

Measuring Scoop Enclosed

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled.
May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Wash with soap and water after handling and before eating, drinking or smoking. Avoid contamination of feed, food and foodstuffs. Do not graze or cut soybean forage for feed.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate any body of water by cleaning of equipment and disposal of wastes and containers. Exposed treated seeds may be hazardous to birds and other wildlife. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil. Dispose of all excess treated seeds by burial away from bodies of water.

ENDANGERED SPECIES RESTRICTIONS

Before using this pesticide on soybeans in the counties listed below, you must obtain the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED

SPECIES for the county in which the product is to be used. The bulletin is available from your County Extension Agent, State Fish and Game Office, or your pesticide dealer. Use of this product in a manner inconsistent with the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES is a violation of Federal laws.

ALABAMA - Lauderdale, Limestone and Madison : ARIZONA - Graham,
Maricopa, Mohave, Pima, Pinal, and Santa Cruz : ARKANSAS - Benton and
Polk : MISSISSIPPI - Claiborne, Copiah and Hinds : MISSOURI -
Barry, Benton, Camden, Christian, Dallas, Greene, Hickory, Jasper,
Lawrence, Miller, Newton, Osage, Polk, St. Clair, Stone and Webster :
NEVADA - Clark : NEW MEXICO - Chaves, DeBaca and Eddy : OHIO -
Pickaway OKLAHOMA - Delaware, McCurtain and Pushmataha : OREGON
- Lake : TENNESSEE - Lawrence and Wayne : TEXAS - Bastrop,
Burleson, Comal, Harris, Hays, Jeff Davis, Pecos and Reeves : UTAH -
Utah and Washington.

PRODUCT INFORMATION


AGROSOL, when used at the correct rate, protects seeds from soil and seed
borne diseases that cause damping-off, seed decay, seedling blights and
pod and stem blight (Phomopsis spp.) during the germination and early
seedling stages. AGROSOL provides essential molybdenum for plant growth
where soils are found to be deficient in available molybdenum.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labelling.

Planter box application - Use AGROSOL at a rate of 2 oz. per bushel
(one scoop per bushel). Mix thoroughly with seed in the planter box as
follows:

For Conventional Plate Planters and MAX-EMERGE^R Plateless Planters by
John Deere - Pour one bushel of seed into each planter box. Pour enough
AGROSOL on top of the seed to treat one bushel. Stir thoroughly with an
axe handle (purchased and used for this purpose only) until all the seed
is uniformly coated. (MAX-EMERGE^R is a reg. T.M. of Deere and Co.).



NOTE: Avoid very deep seeding and exceptionally early sowing under poor growing conditions for maximum benefits. Use only recommended rates. Lower amounts may not give the desired control. Regularly inspect planter plates of seed drill and clean as required. If coating of planter plate occurs (i.e. under conditions of high humidity), add small amounts of graphite to facilitate uniform seeding.

Clean-up for Electric Eye Population Monitors: At the end of each day of use, clean out the electric-eye monitors with brushes provided by the planter manufacturers, according to their directions.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

A dust mask must be worn when mixing and loading this product.

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous.

Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty container and tap side and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

WARRANTY STATEMENT

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

CONTENTS AND CONTAINER MADE IN CANADA

