

Flea Shield™

DIRECTIONS To control fleas on dogs, remove at a distance of 12 inches from the dog, the plastic button. Be sure arrow points toward the dog. Hold at a convenient distance from the dog and spray liberally from all angles. Brush the plastic button with the dog's hair with the hand and a low clothes brush. Continue to spray the dog's head. Keep the dog back. Repeat this procedure every 10 to 15 days. For flea control on cats, apply Flea Shield to the cat's fur. Repeat every 10 to 15 days. For flea control on horses, apply Flea Shield to the horse's coat. Repeat every 10 to 15 days. For flea control on birds, apply Flea Shield to the bird's feathers. Repeat every 10 to 15 days. For flea control on small mammals, apply Flea Shield to the animal's fur. Repeat every 10 to 15 days. For flea control on reptiles, apply Flea Shield to the animal's skin. Repeat every 10 to 15 days. For flea control on amphibians, apply Flea Shield to the animal's skin. Repeat every 10 to 15 days. For flea control on insects, apply Flea Shield to the insect. Repeat every 10 to 15 days. For flea control on arachnids, apply Flea Shield to the arachnid. Repeat every 10 to 15 days. For flea control on mites, apply Flea Shield to the mite. Repeat every 10 to 15 days. For flea control on ticks, apply Flea Shield to the tick. Repeat every 10 to 15 days. For flea control on lice, apply Flea Shield to the louse. Repeat every 10 to 15 days. For flea control on bed bugs, apply Flea Shield to the bed bug. Repeat every 10 to 15 days. For flea control on cockroaches, apply Flea Shield to the cockroach. Repeat every 10 to 15 days. For flea control on termites, apply Flea Shield to the termite. Repeat every 10 to 15 days. For flea control on ants, apply Flea Shield to the ant. Repeat every 10 to 15 days. For flea control on bees, apply Flea Shield to the bee. Repeat every 10 to 15 days. For flea control on wasps, apply Flea Shield to the wasp. Repeat every 10 to 15 days. For flea control on flies, apply Flea Shield to the fly. Repeat every 10 to 15 days. For flea control on mosquitoes, apply Flea Shield to the mosquito. Repeat every 10 to 15 days. For flea control on sand fleas, apply Flea Shield to the sand flea. Repeat every 10 to 15 days. For flea control on biting flies, apply Flea Shield to the biting fly. Repeat every 10 to 15 days. For flea control on house flies, apply Flea Shield to the house fly. Repeat every 10 to 15 days. For flea control on fruit flies, apply Flea Shield to the fruit fly. Repeat every 10 to 15 days. For flea control on gnats, apply Flea Shield to the gnat. Repeat every 10 to 15 days. For flea control on midges, apply Flea Shield to the midge. Repeat every 10 to 15 days. For flea control on mosquitoes, apply Flea Shield to the mosquito. Repeat every 10 to 15 days. For flea control on sand fleas, apply Flea Shield to the sand flea. Repeat every 10 to 15 days. For flea control on biting flies, apply Flea Shield to the biting fly. Repeat every 10 to 15 days. For flea control on house flies, apply Flea Shield to the house fly. Repeat every 10 to 15 days. For flea control on fruit flies, apply Flea Shield to the fruit fly. Repeat every 10 to 15 days. For flea control on gnats, apply Flea Shield to the gnat. Repeat every 10 to 15 days. For flea control on midges, apply Flea Shield to the midge. Repeat every 10 to 15 days.

WARNING: Use only as directed. Harmful if swallowed. Irritation to the eyes, nose, throat, and skin may occur. Avoid contact with eyes, nose, throat, and skin. Wash hands with soap and water after handling. Do not spray toward your pet's face or onto its snout. Do not apply to broken, irritated or sensitive skin. If an irritation develops, discontinue use. Avoid contamination of food, utensils, and food preparation areas. Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator. KEEP OUT OF REACH OF CHILDREN.

ACTIVE INGREDIENT 2-propoxyphenyl methyl carbamate 100.0%

INERT INGREDIENTS 4-tert-butylphenol 34.75%
 1,1,1-trichloro-2,2,2-trifluoroethane 100.0%

Is a cholinesterase inhibitor, atropine sulfate antidote.
 # 1013 EPA Reg. No. 8004 CA 001 07

Dist. by **Lambert Kay**
 Div. of Carter-Wallace, Inc., Irvine, Ca. 92705



Flea Shield™

**FOR DOGS
 KILLS FLEAS AND TICKS**



**WARNING:
 KEEP OUT
 OF REACH
 OF CHILDREN**

SEE BACK
 PANEL FOR
 ADDITIONAL
 WARNING

NET WT. 7 OZ.

ACCEPTED
 JUL 19 1974
 UNDER
 FURNISHING
 FOR THE
 U.S. ARMY
 MEDICAL CENTER

Customer <i>MARLBORO PACKAGING</i>			Design <i>LAMBERT KAY FLEA SHIELD FOR DOGS</i>		
Eng. 22165	S.O. 263-1912-749P	Size 202 x 509	C.E. 5687x6.741	B.P. 1-8	Date 5-6-74
Colors <i>red, blue, yellow, brown, & black on w/c.</i>			Remarks <i>colors ident. to: 263-1524-73</i>		
Prepared by <i>[illegible]</i> Checked by <i>[illegible]</i> Approved by <i>[illegible]</i> Submitted by <i>[illegible]</i> Date <i>[illegible]</i>			Prepared by <i>[illegible]</i> Checked by <i>[illegible]</i> Approved by <i>[illegible]</i> Submitted by <i>[illegible]</i> Date <i>[illegible]</i>		